

UNIVERSITY OF NOTRE DAME
DEPARTMENT OF AEROSPACE AND MECHANICAL ENGINEERING

Professor Atassi

AME-562

INVERSION OF THE CAUCHY TYPE INTEGRAL

Consider the integral

$$\frac{1}{\pi} \oint_a^b \frac{\varphi(\tau)}{\tau - t} d\tau = f(t) \quad (1)$$

where $b > a$ and $\varphi(t)$ satisfies Holder's condition on the interval $[a, b]$. The inverse problem can be stated as follows: "Given the function $f(t)$ on the interval $[a, b]$, find the density function $\varphi(t)$." The inverse problem is *not* unique. It has the following solutions:

1. Solution bound at a and unbound at b

$$\varphi(t) = -\frac{1}{\pi} \sqrt{\frac{t-a}{b-t}} \oint_a^b \sqrt{\frac{b-\tau}{\tau-a}} \frac{f(\tau)}{\tau-t} d\tau \quad (2)$$

2. Solution bound at b and unbound at a

$$\varphi(t) = -\frac{1}{\pi} \sqrt{\frac{b-t}{t-a}} \oint_a^b \sqrt{\frac{\tau-a}{b-\tau}} \frac{f(\tau)}{\tau-t} d\tau \quad (3)$$

3. Solution unbound at both ends a and b

$$\varphi(t) = -\frac{1}{\pi} \frac{1}{\sqrt{(b-t)(t-a)}} \left[\oint_a^b \frac{\sqrt{(\tau-a)(b-\tau)} f(\tau)}{(\tau-t)} d\tau + C \right] \quad (4)$$

where c is an arbitrary constant.

4. Solution bound at both ends a and b

In general, the inverse solution may not exist. However, if the function $f(t)$ satisfies the condition

$$\oint_a^b \frac{f(\tau)}{\sqrt{(\tau-a)(b-\tau)}} d\tau \quad (5)$$

then, we have

$$\varphi(t) = -\frac{1}{\pi} \sqrt{(b-t)(t-a)} \oint_a^b \frac{f(\tau)}{\sqrt{(\tau-a)(b-\tau)}} \frac{d\tau}{(\tau-t)} \quad (6)$$