

CONTENTS

FIGURES	v
TABLES.....	x
ACKNOWLEDGEMENTS	xiii
CHAPTER 1: INTRODUCTION TO FAST BEAM-FOIL SPECTROSCOPY	1
CHAPTER 2: MULTIPLY EXCITED STATES IN OXYGEN, FLUORINE AND NEON	5
CHAPTER 3: THEORY	11
3.1 The Multi-Configuration Hartree-Fock (MCHF) Method with the QED and Higher-order Relativistic Corrections.....	11
3.1.1 <i>Multi-Configuration Hartree-Fock Method</i>	11
3.1.2 <i>Higher-Order Relativistic Contributions and QED Effects</i>	14
3.2 The Multi-Configuration Dirac-Fock (MCDF) Method	16
3.3 Multi-Configuration Hartree-Fock (MCHF) Calculations and Results on Quintet States in Beryllium-like C III, N IV, O V, F VI and Ne VII.....	19
3.4 The Multi-Configuration Hartree-Fock (MCHF) Calculations for Sextet States in Boron-like O IV, F V and Ne VI	29
3.4.1 <i>MCHF calculations for the sextet states $1s2s2p^3\ ^6S^o$, $1s2s2p^23s\ ^6P$ and $1s2p^33s\ ^6S^o$ of boron-like O IV, F V and Ne VI.....</i>	29
3.4.2 <i>MCHF calculations for the $1s2s2p^23l\ ^6L-1s2p^33l\ ^6L'$, $l=p, d$, electric-dipole transitions of boron-like O IV, F V and Ne VI.....</i>	33

3.5 The Multi-Configuration Dirac-Fock (MCDF) Calculations for Sextet States in Boron-like O IV, FV and Ne VI.	34
CHAPTER 4: FAST BEAM-FOIL EXPERIMENT.....	36
4.1 The Beam-Foil Light Source.....	36
4.2 Excitation Mechanisms.....	38
4.3 Spectrometers and Detectors.....	46
4.4 Targets.....	51
4.5 Experimental Method.....	51
4.6 Experimental Problems.....	54
CHAPTER 5: RESULTS FOR SEXTET TRANSITIONS IN O IV, F V AND NE VI.....	57
5.1 The $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ Transitions in O IV, F V and Ne VI.	57
5.1.1 <i>The $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ Transitions in O IV.</i>	57
5.1.2 <i>The $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ Transitions in F V.</i>	60
5.1.3 <i>The $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ Transitions in Ne VI.</i>	62
5.1.4 <i>Comparison of the $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ Transitions with Theory along Isoelectronic Sequence</i>	63
5.1.5 <i>Summary</i>	69
5.2 The $1s2s2p^23p\ ^6L^o-1s2p^33p\ ^6P, L=S, P, D$ Transitions in O IV, F V and Ne VI.....	71
5.2.1 <i>The $1s2s2p^23p\ ^6L^o-1s2p^33p\ ^6P$ Transitions in O IV.</i>	73
5.2.2 <i>The $1s2s2p^23p\ ^6L^o-1s2p^33p\ ^6P$ Transitions in F V.</i>	75
5.2.3 <i>The $1s2s2p^23p\ ^6L^o-1s2p^33p\ ^6P$ Transitions in Ne VI.</i>	76
5.2.4 <i>Comparison of the $1s2s2p^23p\ ^6L^o-1s2p^33p\ ^6P$ Transitions with Theory along Isoelectronic Sequence</i>	77
5.2.5 <i>Summary</i>	85

5.3 The $1s2s2p^23d^6L - 1s2p^33d^6D^o$, L=P, D, F Transitions in O IV, F V and Ne VI.....	86
5.3.1 <i>The $1s2s2p^23d^6L - 1s2p^33d^6D^o$ Transitions in O IV.</i>	86
5.3.2 <i>The $1s2s2p^23d^6L - 1s2p^33d^6D^o$ Transitions in F V.</i>	92
5.3.3 <i>The $1s2s2p^23d^6L - 1s2p^33d^6D^o$ Transitions in Ne VI.</i>	94
5.3.4 <i>Comparisons with Theory of the $1s2s2p^23d^6L - 1s2p^33d^6D$ Transitions along B I Isoelectronic Sequence</i>	96
5.3.5 <i>Summary</i>	104
5.4 The Sextet Transitions and Energy Levels in O IV, F V and Ne VI.....	105
5.5 Outlook - Preliminary Results of the $1s2s2p^3^6S - 1s2s2p^23l^6L$ Transitions from Beam-gas Experiment.....	130
CHAPTER 6: TRIPLET TRANSITIONS BETWEEN DISPLACED TERMS IN O V, F VI AND NE VII.....	135
<i>Configuration Interactions</i>	136
CHAPTER 7: CONCLUSIONS	141
REFERENCES	143

FIGURES

<p>FIG. 2.1 Term diagram of the doubly excited sextet states of O IV. The mean wavelength for $1s2s2p^2 3s \ ^6P-1s2p^3 3s \ ^6S^o$ transitions in O IV is shown. The energies are not to scale.</p>	6
<p>FIG. 2.2 The recent work of Lapierre and Knystautas [17] on possible sextet transitions in Ne VI highlights the significance and difficulties in this sequence. The tentative identifications of the two transitions at 12.0 and 10.6 nm show rather weak lines and overwhelming blending problem. [The Figure is taken from [17]].</p>	8
<p>FIG. 2.3 The work of Blanke et al. [15] in searching for the two possible sextet transitions in C III, O IV and F V indicates blending problems in all six cases. [The Figure is taken from [15]].</p>	9
<p>FIG. 3.1 The deviations between the theoretical and experimental transition energies for the $1s2s2p^2 \ ^5P_J-1s2p^3 \ ^5S^o_2$ transitions. Here the theoretical transition energy is the center of gravity of the $1s2s2p^2 \ ^5P_J-1s2p^3 \ ^5S^o_2$, $J=1,2,3$, transition energies (computed from the fine structure lines by this work) with the results of theoretical analysis, and the experimental transition energy is the center of gravity of the $1s2s2p^2 \ ^5P_J-1s2p^3 \ ^5S^o_2$, $J=1,2,3$, transition energies (computed from the observed lines [33-35]) with the results of theoretical analysis.</p>	24
<p>FIG. 3.2 The deviations between the theoretical (this work) and experimental [32-35,37] lifetimes for the $1s2p^3 \ ^5S^o_2$ state in the beryllium sequence.</p>	25
<p>FIG. 3.3 Theoretical fine and hyperfine structure of the $1s2s2p^2 \ ^5P_{J=1,2,3}$ and $1s2p^3 \ ^5S^o_2$ states in F VI by this work.</p>	28
<p>FIG. 4.1 The principle of fast beam-foil spectroscopy. Ions from the accelerator are mass-analyzed and sent through a thin foil. The radiation emitted on the downstream is studied by optical, x-ray, or electron spectroscopy. Lifetime and quantum beat experiments are also indicated. [The figure is taken from [19]].</p>	36

FIG. 4.2	Scaling of ion energy change with reduced ion velocity ($V/Z^{2/3}$). Bohr and others suggested that the charge states of an ion could be estimated by assuming that it would lose all of its electrons whose orbital velocity was less than the ion velocity. The orbital velocity of electron was considered to be proportional to $Z^{2/3}$, so this concept can be illustrated by plotting stopping vs an ion velocity reduced by this electron velocity factor. [The figure is taken from [57]].	41
FIG. 4.3	Scaling of ion energy change with reduced ion velocity [$V/(V_F Z^{2/3})$]. Bohr estimated the charge state of an ion by assuming that it was stripped of any of its electrons whose orbital velocity was less than its velocity see Fig. 4.2 This concept lasted for thirty years because of its elegant simplicity. Brandt extended this idea to relate the ion velocity and its electrons to the electron velocity of the foil, in which it is passing. Ziegler took this concept and made an explicit formulation that reduced the ion velocity could take the form of $V/(V_F Z^{2/3})$. [The figure is taken from [57]].	41
FIG. 4.4	The optimal ion beam energy of the ionization degrees with the effective charge $Z^* = (Z \cdot \bullet)$ and most abundant isotope of atomic weight A listed in Table 4.1	43
FIG. 4.5	Equilibrium charge distribution of: (a) carbon, (b) nitrogen, (c) oxygen, and (d) neon as a function of incident ion. From the above we can find the optimal energy to get the ions needed in the experiments. For oxygen we needed to study the higher lying sextet states in O IV (O^{3+}). In (c) we could see that 2-3 MeV is a possible optimal energy [82]. [The figure is taken from [82]].	44
FIG. 4.6	The 1-m Seya-Namioka Type Concave Grating Monochromator at Liège. The distances $l_a = l_b = 1$ meter and the angle $\bullet = 70.25$ degrees are fixed. In order to scan wavelengths, the grating is rotated about its vertical axis. For the maximum efficiency A and B are on the Rowland circle.	47
FIG. 4.7	The electronics and data-taking equipment in BFS experiments at the University of Liège	51
FIG. 4.8	The experimental setup of fast beam-foil spectroscopy on a 2 MeV Van de Graaff accelerator beam line at the University of Liège [14,106]. [The figure is taken from University of Liège].	52
FIG. 4.9	The beam-foil interaction in the vacuum chamber works as an efficient light source in the Beam-foil Spectroscopy at the University of Liège. [The figure is taken from University of Liège].	53

FIG.5.1.1 The transitions in the beam-foil spectra of oxygen, fluorine and neon, recorded at the different energies. The beam energies and spectrometer slit widths are shown.....	59
FIG. 5.1.2 Scaled isoelectronic plot of the center of gravity wavelength (a) and transition energy deviations from experiment (b) of the $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ multiplet transitions for the Boron sequence. The experimental and calculated points are denoted by the following: circles with error bars inside for experiments, + for SCHF, • for SCHFT, • for MCHF, * for MCHFT, • for SCDF, and diamonds for MCDF.....	65
FIG. 5.1.3 Comparison of the experimental (opened circles) and theoretical (others) fine structures for O IV, FV and Ne VI. The experimental and calculated points are denoted by the following: circles with error bars inside for experiments, • for SCHFT, • for MCHFT, diamonds for SCDF, and * for MCDF.....	66
FIG. 5.1.4 Isoelectronic comparison of the QED effects and higher-order corrections of the $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ transitions for the boron sequence. The calculated points are denoted by the following: diamonds for QED effects from SCHF, • for higher order corrections from SCHF, • for QED effects from MCHF, and • for higher order corrections from MCHF. The fitted QED effects from both SCHF and MCHF are denoted by solid line.....	67
FIG. 5.1.5 Isoelectronic comparison of the theoretical and experimental mean wavelengths of the $1s2s2p^23s\ ^6P-1s2p^33s\ ^6S^o$ transitions for the boron sequence.	68
FIG. 5.2.1 The $1s2s2p^23p\ ^6L$ and $1s2p^33p\ ^6P$ transitions in the beam-foil spectra of oxygen, fluorine and neon, recorded at the different energies. The beam energies and spectrometer slit widths are indicated.	72
FIG. 5.2.2 The identifications of the $1s2s2p^23p\ ^6D_J - 1s2p^33p\ ^6P_J$ and $1s2s2p^23p\ ^6S_{5/2} - 1s2p^33p\ ^6P_J$ transitions of O IV in the experimental spectrum of O at the energy of 2.5 MeV. In (a) * means all possible J's of the upper states allowed by E1 transition rules. In (b) * means 3/2, 5/2 and 7/2.....	75
FIG. 5.2.3 The deviations between the theoretical and experimental transition energies for the $1s2s2p^23p\ ^6D_J-1s2p^33p\ ^6P_J$ transitions. Here the theoretical transition energy is the center of gravity of the $1s2s2p^23p\ ^6D_J-1s2p^33p\ ^6P_J$ transition energies (computed from the fine structure lines by this work) with the results of theoretical analysis, and the experimental transition energy is the center of gravity of the $1s2s2p^23p\ ^6D_J-1s2p^33p\ ^6P_J$ transition energies (computed from the observed lines) with the results of theoretical analysis.	82

FIG. 5.2.4	The deviations between the theoretical and experimental transition energies for the $1s2s2p^23p\ ^6P_J-1s2p^33p\ ^6P_J$ transitions. Here the theoretical transition energy is the center of gravity of the $1s2s2p^23p\ ^6P_J-1s2p^33p\ ^6P_J$ transition energies (computed from the fine structure lines by this work) with the results of theoretical analysis, and the experimental transition energy is the center of gravity of the $1s2s2p^23p\ ^6P_J-1s2p^33p\ ^6P_J$ Transition energies (computed from the observed lines) with the results of theoretical analysis.	82
FIG. 5.2.5	The deviations between the theoretical and experimental transition energies for the $1s2s2p^23p\ ^6S_J-1s2p^33p\ ^6P_J$ transitions. Here the theoretical transition energy is the center of gravity of the $1s2s2p^23p\ ^6S_J-1s2p^33p\ ^6P_J$ transition energies (computed from the fine structure lines by this work) with the results of theoretical analysis, and the experimental transition energy is the center of gravity of the $1s2s2p^23p\ ^6S_J-1s2p^33p\ ^6P_J$ transition energies (computed from the observed lines) with the results of theoretical analysis.	83
FIG. 5.2.6	The QED and higher-order corrections for (a) the $1s2s2p^23p\ ^6D-1s2p^33p\ ^6P$, (b) $1s2s2p^23p\ ^6P-1s2p^33p\ ^6P$ and (c) $1s2s2p^23p\ ^6S-1s2p^33p\ ^6P$ transitions in the B I isoelectronic sequence.	84
FIG. 5.3.1	The $1s2s2p^23d\ ^6L_J - 1s2p^33d\ ^6D_J$, $L=P, D, F$ transitions in the beam-foil spectra of oxygen, fluorine and neon, recorded at different energies. The beam energies and spectrometer slit widths in are indicated.	88
FIG. 5.3.2	The identifications of the $1s2s2p^23d\ ^6F_J - 1s2p^33d\ ^6D_J$ transitions of O IV in the experimental spectrum of $^{20}\text{O}^+$ at the energy of 2.5 MeV.....	91
FIG. 5.3.3	The deviations between the theoretical and experimental transition energies for the $1s2s2p^23d\ ^6F-1s2p^33d\ ^6D$ transitions.....	97
FIG. 5.3.4	The deviations between the theoretical and experimental transition energies for the $1s2s2p^23d\ ^6D-1s2p^33d\ ^6D$ transitions.	98
FIG. 5.3.5	The deviations between the theoretical and experimental transition energies for the $1s2s2p^23d\ ^6P-1s2p^33d\ ^6D$ transitions.....	98
FIG. 5.3.6	The reduced experimental transition energies $\bullet E/(Z-1.86)$ for the $1s2s2p^23d\ ^6F_J-1s2p^33d\ ^6D_J$ transitions.	99
FIG. 5.3.7	The reduced experimental transition energies $\bullet E/(Z-1.86)$ for the $1s2s2p^23d\ ^6D_J-1s2p^33d\ ^6D_J$ transitions.	99
FIG. 5.3.8	The reduced experimental transition energies $\bullet E/(Z-1.86)$ for the $1s2s2p^23d\ ^6P_J-1s2p^33d\ ^6D_J$ transitions.	100
FIG. 5.3.9	The lifetimes (in ps) for the $1s2s2p^23d\ ^6P^e$ states in the boron sequence Experimental values are taken from Table 5.3.5.....	101
FIG. 5.3.10	The lifetimes (in ps) for the $1s2p^33d\ ^6D^o$ states in the boron sequence.....	102

FIG. 5.3.11 The QED and higher-order corrections for (a) the $1s2s2p^23d\ ^6F-1s2p^33d\ ^6D$, (b) $1s2s2p^23d\ ^6D-1s2p^33d\ ^6D$ and (c) $1s2s2p^23d\ ^6P-1s2p^33d\ ^6D$ transitions in the B I isoelectronic sequence.	103
FIG. 5.4.1 Doubly excited sextet levels $1s2p^2nln'l'\ ^6L$ of 5-electron O IV with aligned spins. The level energies are in cm^{-1}	123
FIG. 5.4.2 Grotrian Diagram of Doubly excited sextet levels $1s2p^2nln'l'\ ^6L$ of O IV with aligned spins. The wavelengths are in \AA	124
FIG. 5.4.3 Doubly excited sextet levels $1s2p^2nln'l'\ ^6L$ of 5-electron F V with aligned spins. The level energies are in cm^{-1}	125
FIG. 5.4.4 Grotrian Diagram of Doubly excited sextet levels $1s2p^2nln'l'\ ^6L$ of F V with aligned spins. The wavelengths are in \AA	126
FIG. 5.4.5 Doubly excited sextet levels $1s2p^2nln'l'\ ^6L$ of 5-electron Ne VI with aligned spins. The level energies are in cm^{-1}	127
FIG. 5.4.6 Grotrian Diagram of Doubly excited sextet levels $1s2p^2nln'l'\ ^6L$ of Ne VI with aligned spins. The wavelengths are in \AA	128
FIG. 5.5.1 Block diagram of the beam-gas experimental apparatus showing the procedures to be followed to prepare N^{5+} beam in low (3%) and high (48%) fractions of the metastable states.]. [The figure is taken from [128]].	129
FIG. 5.5.2 Spectra of F V in the 150-180 \AA region for direct and prepared beams. [The figure is taken from [129]].	130
FIG. 5.5.3 The Boron-like $2p-3d$ and $2p-3s$ transitions in normal and displaced terms for (a) O IV, (b) F V and (c) Ne VI. A, B and C: $2s2p^2\ ^4P-2s2p(^3P^o)3d\ ^4P^o$, $^4D^o$ and $^4F^o$. a, b and c: $2s2p^2\ ^2D-2s2p(^3P^o)3d\ ^2P^o$, $^2F^o$ and $^2D^o$. D and a': $2s2p^2\ ^4P-2s2p(^3P^o)3s\ ^4P^o$ and $2s2p^2\ ^2S-2s2p(^3P^o)3d\ ^2P^o$. • and • : $2s^22p\ ^2P^o-2s^23d\ ^2D$ and $2s^22p\ ^2P^o-2s^23d\ ^2S$. [The figure is taken from [129]].	132
FIG. 5.5.4 Reduced transition energy $\bullet E/(Z-5.77)$ of the $1s2s2p^3\ ^6S - 1s2s2p^23s\ ^6P$ transition of boron-like sequence. The squares are for the beam-gas experimental values by this work. The diamonds are for the beam-gas experimental values by Blanke and Lapierre taken from Table 5.5.1	134

TABLES

<p>TABLE 3.1 The energies E (in cm^{-1}) and wavelengths λ (in \AA) for the $1s2s2p^2$ 5P_J-$1s2p^3$ $^5S^{\circ}_2$ transitions in C III, N IV, O V, F VI, and Ne VII by this work. We also list the deviations between the theoretical and experimental transition energies for the $1s2s2p^2$ 5P_J-$1s2p^3$ $^5S^{\circ}_2$ transitions.</p>	23
<p>TABLE 3.2 Lifetimes (in ns) of the $1s2p^3$ $^5S^{\circ}_2$ states in O V, F VI and Ne VII.</p>	25
<p>TABLE 3.3 A short wavefunction expansion CI for the $1s2s2p^3$ $^6S^{\circ}$, $1s2s2p^23s$ 6P, $1s2p^33s$ $^6S^{\circ}$, $1s2s2p^23p$ $^6D^{\circ}$, $1s2p^33p$ $^6P^{\circ}$, $1s2s2p^23d$ 6F, $1s2p^33d$ $^6D^{\circ}$, $1s2s2p^24f$ $^6G^{\circ}$ and $1s2p^34f$ 6F states of boron-like O IV and F V. $\sum_{i=1}^4 c_i^2$ is the total contributions of the CSFs listed the above.</p>	31
<p>TABLE 3.4 Energies (in cm^{-1}) as well as the QED effects and higher order corrections of the $1s2s2p^3$ $^6S^{\circ}_{5/2}$ states in boron-like O IV, F V and Ne VI.</p>	32
<p>TABLE 4.1 The optimal ion beam energies (in MeV) of the ionization degrees with the effective ion charge $Z^* = (Z - \sigma)$, σ is electronic screening of the beam ion after collision with foil atom, and most abundant isotope of atomic weight A, by this work.</p>	42
<p>TABLE 4.2 Comparisons of the calculated optimal beam ion energies (in MeV) by this work with experimental data (in MeV) from [82].</p>	45
<p>TABLE 5.1.1 The energies (in cm^{-1}) and wavelengths (in \AA) for $1s2s2p^23s$ 6P_J-$1s2p^33s$ $^6S^{\circ}_{5/2}$ transitions in O IV, F V, and Ne VI by this work. Here we also list the QED effects and higher-order corrections for the $1s2s2p^23s$ 6P_J-$1s2p^33s$ $^6S^{\circ}_{5/2}$ transitions in O IV, F V, and Ne VI. The errors for calculations are root mean square differences between the calculated and experimental values.</p>	61
<p>TABLE 5.1.2 Fine structure (in cm^{-1}) of the $1s2s2p^23s$ 6P_J states in O IV, F V and Ne VI.</p>	64
<p>TABLE 5.1.3 Lifetime (in ps) of the $1s2s2p^23s$ 6P and $1s2p^33s$ $^6S^{\circ}_{5/2}$ states in O IV, F V and Ne VI.</p>	64

TABLE 5.1.4 Electron screening • for the 2s and 2p valence electron of the 1s2s2p ² 3s ⁶ P and 1s2p ³ 3s ⁶ S _{5/2} states in O IV, F V and Ne VI.	64
TABLE 5.2.1 The energies E (in cm ⁻¹) and wavelengths • (in Å) for the 1s2s2p ² 3p ⁶ L _J -1s2p ³ 3p ⁶ P _J transitions in O IV, F V and Ne VI by this work. We also list the deviations between the theoretical and experimental transition energies for the 1s2s2p ² 3p ⁶ L _J -1s2p ³ 3p ⁶ P _J transitions.	78
TABLE 5.2.1 Continued.	79
TABLE 5.2.1 Continued.	80
TABLE 5.3.1 The energies E (in cm ⁻¹) and wavelengths • (in Å) for the 1s2s2p ² 3d ⁶ L _J -1s2p ³ 3d ⁶ D _J transitions in O IV by this work. We also list the deviations between the theoretical and experimental transition energies for the 1s2s2p ² 3d ⁶ L _J -1s2p ³ 3d ⁶ D _J transitions.	89
TABLE 5.3.2 The wavelengths • (in Å) for the 1s ² 2p3p ¹ D ₂ -1s ² 2p4d ¹ F ₃ transitions in O V by this work.	91
TABLE 5.3.3 The energies E (in cm ⁻¹) and wavelengths • (in Å) for the 1s2s2p ² 3d ⁶ L _J -1s2p ³ 3d ⁶ D _J transitions in F V by this work. We also list the deviations between the theoretical and experimental transition energies for the 1s2s2p ² 3d ⁶ L _J -1s2p ³ 3d ⁶ D _J transitions.	93
TABLE 5.3.4 The energies E (in cm ⁻¹) and wavelengths • (in Å) for the 1s2s2p ² 3d ⁶ L _J -1s2p ³ 3d ⁶ D _J transitions in Ne VI by this work. We also list the deviations between the theoretical and experimental transition energies for the 1s2s2p ² 3d ⁶ L _J -1s2p ³ 3d ⁶ D _J transitions.	95
TABLE 5.3.5 Lifetimes (in ps) of the 1s2s2p ² 3d ⁶ P and 1s2p ³ 3d ⁶ D states in N III, O IV, F V, Ne VI and Na VII.	101
TABLE 5.4.1 The transition wavelengths (in Å) and energies (in cm ⁻¹) of the sextet states of O IV.	107
TABLE 5.4.2 The transition wavelengths (in Å) and energies (in cm ⁻¹) of the sextet states of F V.	112
TABLE 5.4.3 The transition wavelengths (in Å) and energies (in cm ⁻¹) of the sextet states of Ne VI.	117
TABLE 5.4.4 The relative energies (in cm ⁻¹) of the sextet states of O IV, F V and Ne VI.	122
TABLE 5.5.1 The sextet transition wavelengths (in Å) of O IV, F V and Ne VI.	133
TABLE 6.1 The transition wavelengths (in Å) and energies (in cm ⁻¹) of the triplet states of O V.	138

TABLE 6.2 The transition wavelengths (in Å) and energies (in cm ⁻¹) of the triplet states of F VI.	139
TABLE 6.3 The transition wavelengths (in Å) and energies (in cm ⁻¹) of the triplet states of Ne VII.	139

ACKNOWLEDGMENTS

My deepest appreciation is to Professor Henry Gordon Berry for his constant encouragement, stimulation and help in starting the program, performing the experiments and revising the manuscript.

I appreciate Professors Garnir, Bastin and the members of the beam-foil group at Liège for invaluable help during long data-collecting of this investigation.

I am very grateful to Professors A. Eugene Livingston and J. Désesquelles for allowing me to use your unpublished spectra. I am also greatly indebted to Drs. Tomohiro Shibata and Igor Savukov for much valuable help and many discussions.

I thank all members of the Atomic Physics Group at University of Notre Dame for your essential help and useful discussions. I also want to thank the faculties, staffs and many others in the Department of Physics for your strong supports and cordial amity, and the growing of my spirit during my stay as a graduate student in the big family of Notre Dame.

I acknowledge the financial support from the Department of Physics and Graduate School of the University of Notre Dame for my study, research and living at the University of Notre Dame.

Thank you all very very much.