

# Systematic measurement of keV-neutron capture cross sections and capture gamma-ray spectra of Sn isotopes

J. NISHIYAMA<sup>1,\*</sup>, M. IGASHIRA<sup>1</sup>, T. OHSAKI<sup>1</sup>, G. N. KIM<sup>2</sup>, W. C. CHUNG<sup>3</sup> and T. I. RO<sup>3</sup>

<sup>1</sup> *Research Laboratory for Nuclear Reactors, Tokyo Institute of Technology,  
2-12-1-N1-26 O-okayama, Meguro-ku, Tokyo 152-8550, Japan*

<sup>2</sup> *Institute of High Energy Physics, Kyungpook National University, Daegu 702-701, Korea*

<sup>3</sup> *Department of Physics, Dong-A University, Busan 604-714, Korea*

The neutron capture cross sections of long-lived fission products (LLFPs) are important physical quantities for the study on the transmutation of radioactive nuclear wastes. The nuclide <sup>126</sup>Sn is one of the LLFPs. However, there is no experimental data for <sup>126</sup>Sn, because the preparation of high-purity sample is difficult and, moreover, gamma-ray radiation from a sample causes a serious background.

On the other hand, keV-neutron capture cross sections and capture gamma-ray spectra of stable Sn isotopes contain important information which is useful for the evaluation of capture cross sections of <sup>126</sup>Sn. Thus, we have started a systematic measurement of keV-neutron capture cross sections and capture gamma-ray spectra of stable Sn isotopes. In the present contribution, the results for <sup>117,119</sup>Sn are shown.

The capture cross sections and capture gamma-ray spectra of <sup>117,119</sup>Sn were measured in the incident neutron energy region from 10 to 100 keV and at 550 keV, using the 3-MV Pelletron accelerator of the Research Laboratory for Nuclear Reactors at the Tokyo Institute of Technology. Pulsed keV neutrons were produced from the <sup>7</sup>Li(p,n)<sup>7</sup>Be reaction with a 1.5-ns bunched proton beam from the accelerator. The pulse-repetition rate was 4 MHz. The <sup>117,119</sup>Sn samples were highly enriched metal plates, and the net weight of each sample was about 1 g. Capture gamma rays were detected with a large anti-Compton NaI(Tl) spectrometer by means of a time-of-flight method. A pulse-height weighting technique was applied to the observed capture gamma-ray pulse-height spectra to obtain capture yields. Using the standard capture cross sections of <sup>197</sup>Au, the capture cross sections of <sup>117,119</sup>Sn were derived with the error of about 5%. Capture gamma-spectra were derived by unfolding the observed capture gamma-ray pulse-height spectra. The present cross section results were compared with other experimental data and the evaluated values in JENDL-3.3 and ENDF/B-VI.

The calculation of capture cross sections and capture gamma-ray spectra of <sup>117,119</sup>Sn were performed with the EMPIRE-II code in the incident neutron energy region from 10 to 1000 keV. The calculated results were compared with the present experimental results.

\* Tel. +81-3-5734-3378, Fax. +81-3-5734-2959, Email: 03m19193@nr.titech.ac.jp