

# Isospin symmetry of isobaric triplets along the $N \sim Z$ line\*

J. N. Orce<sup>1</sup>, V. Velázquez<sup>2</sup>, P. Petkov<sup>3,4</sup>, C. J. McKay<sup>1</sup>, S. R. Leshner<sup>1</sup>, S. W. Yates<sup>1,5</sup>,  
and M. T. McEllistrem<sup>1</sup>

<sup>1</sup> Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky  
40506-0055, USA

<sup>2</sup> Departamento de Física, Facultad de Ciencias, Universidad Nacional Autónoma de México,  
Apartado Postal 70-543, 04510 México, D.F., México

<sup>3</sup> Institute for Nuclear Research and Nuclear Energy, Sofia 11784, Bulgaria

<sup>4</sup> Institut für Kernphysik, Universität zu Köln, 50937 Köln, Germany

<sup>5</sup> Department of Chemistry, University of Kentucky, Lexington, Kentucky 40506-0055, USA

A careful measurement of the lifetime of the first  $2_{T=1}^+$  state in  $^{42}\text{Sc}$  has allowed an experimental test of isospin purity in the  $A=42$  isobaric triplet by using the isospin formalism [1]. In the case of testing charge independence of  $T = 1$  nuclear structures, the experimental test involves the comparison of  $E2$  strengths of  $2_{1(T=1)}^+ \rightarrow 0_{1(T=1)}^+$  transitions in isobaric triplets [2]. A lifetime of 69 (18) fs has been determined for the  $2_{1(T=1)}^+$  state in  $^{42}\text{Sc}$ , giving an isoscalar matrix element of  $6.9(9)$  (W.u.)<sup>1/2</sup> [3]. Previous measurements of the lifetimes in the mirror nuclei  $^{42}\text{Ca}$  and  $^{42}\text{Ti}$  provided an isoscalar matrix element of  $7.1(5)$  (W.u.)<sup>1/2</sup>, which is very close to the measured value for  $^{42}\text{Sc}$ . This agreement suggests that charge independence is conserved in the  $A = 42$  isobaric triplet. Globally, the systematics of isoscalar matrix elements,  $M_0$ , present a well-defined trend for isobaric triplets ranging from  $A = 18$  to  $A = 42$ . Shell model calculations have been carried out using the m-scheme numerical code ANTOINE [4] in order to understand this particular trend. The  $2_{1(T=1)}^+ \rightarrow 0_{1(T=1)}^+$  transition energies, reduced transition probabilities and isoscalar matrix elements are reproduced to a high degree of accuracy [5]. The equality of  $M_0$  values between mirror pairs and  $T_Z = 0$  nuclides supports our shell model calculations. The predicted results reproduce the isoscalar trend from  $A=18$  to  $A=42$ ; the variation of  $M_0$  along the  $sd$  shell is explained in terms of the dynamic shell structure, tying together the results from recent experiments. Discrepancies may arise at the extremes of the  $sd$  shell for the  $A = 18$  and  $A = 38$  isobaric triplets, and might be explained in terms of the occupancy of the orbits at the extremes of the shell.

\* This material is based upon work supported by the U.S. National Science Foundation under Grant No. PHY-0354656.

- [1] A. M. Bernstein, V. R. Brown and V. A. Madsen, Phys. Rev. Lett. **42**, 425 (1979).
- [2] P. D. Cottle *et al.*, Phys. Rev. Lett. **88**, 172502 (2002).
- [3] J. N. Orce *et al.*, Phys. Rev. C **70**, 014314 (2004).
- [4] E. Caurier and F. Nowacki, Acta Physica Polonica B **30**, 705 (1999).
- [5] J. N. Orce and V. Velázquez, submitted to Phys. Lett. B (2005).