

# Introduction to Probability and Statistics

## Review of the most common distributions

March 14, 2008

Here is a list of some of the most common probability distributions encountered in statistics, with their basic properties and uses. The first three in each category (discrete, continuous) are the most important.

### 1. Discrete

#### (a) Bernoulli

- i. **Parameter:**  $0 \leq p \leq 1$
- ii. **Mass function:**  $P(X = 1) = p, P(X = 0) = 1 - p$
- iii. **Mean and variance:**  $p$  and  $p(1 - p)$
- iv. **Use:** Models the outcome of a single trial, success probability  $p$

#### (b) Binomial

- i. **Parameters:** Positive integer  $n, 0 \leq p \leq 1$
- ii. **Mass function:**  $P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, k = 0, 1, \dots, n.$
- iii. **Mean and variance:**  $np$  and  $np(1 - p)$
- iv. **Use:** Counts the number of successes when  $n$  independent repetitions of a trial are performed, each with success probability  $p$

#### (c) Poisson

- i. **Parameter:**  $\lambda > 0$
- ii. **Mass function:**  $P(X = k) = \frac{\lambda^k}{k!} e^{-\lambda}, k = 0, 1, 2, \dots$
- iii. **Mean and variance:** Both  $\lambda$
- iv. **Uses:**
  - A. Approximates the binomial when number of trials  $n$  is large and probability of success  $p$  is small,  $\lambda = np$
  - B. Models the number of times a rare event occurs in a time period, when  $\lambda$  is the average number of occurrences

(d) **Hypergeometric**

- i. **Parameters:**  $N, M, n$
- ii. **Mass function:**  $P(X = k) = \frac{\binom{N}{k}\binom{M}{n-k}}{\binom{N+M}{n}}, k = 0, 1, \dots, \min\{N, n\}$
- iii. **Mean and variance:**  $\frac{nN}{N+M}$  and  $\frac{nNM}{(N+M)^2} \left(1 - \frac{n-1}{N+M-1}\right)$
- iv. **Use:** Models the number of desirable elements obtained, if  $n$  items are drawn from a pool containing  $N$  desirable and  $M$  undesirable elements

2. **Continuous**

(a) **Uniform**

- i. **Parameters:**  $\alpha < \beta$
- ii. **Density function:**  $f(x) = \frac{1}{\beta-\alpha}, \alpha \leq x \leq \beta$
- iii. **Mean and variance:**  $\frac{\alpha+\beta}{2}$  and  $\frac{(\beta-\alpha)^2}{12}$
- iv. **Use:** Models the selection of a random number between  $\alpha$  and  $\beta$

(b) **Exponential**

- i. **Parameter:**  $\lambda > 0$
- ii. **Density function:**  $f(x) = \lambda e^{-\lambda x}, x \geq 0$
- iii. **Mean and variance:**  $\frac{1}{\lambda}$  and  $\frac{1}{\lambda^2}$
- iv. **Use:** Models the time spent waiting until an event occurs,  $1/\lambda$  the average waiting time

(c) **Normal**

- i. **Parameters:**  $\mu, \sigma^2$
- ii. **Density function:**  $f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$
- iii. **Mean and variance:**  $\mu$  and  $\sigma^2$  (**Standard Normal** if  $\mu = 0, \sigma^2 = 1$ )
- iv. **Uses:**
  - A. Models the distribution of many observable physical quantities
  - B. Models the error made when measuring device makes measurement
  - C. Approximates the limit of the sum of independent random variables with the same mean and variance

(d) **Chi-squared**

- i. **Parameter:**  $n$
- ii. **Description:**  $X = Z_1^2 + Z_2^2 + \dots + Z_n^2$  where the  $Z_i$ 's are independent standard normals
- iii. **Uses:**
  - A. Models distance of a point in  $n$ -dimensional space from origin if coordinates are independent standard normals
  - B. Suitably scaled, is the sample variance of a sample drawn from a normal population