Math 30530 — Introduction to Probability

Quiz 5 – Friday November 30, 2012

Solutions

Fully grown Indiana grass snakes have a length that is normally distributed with mean 20cm and variance 16cm^2 (with the length of different grass snakes being independent of each other).

1. For what length ℓ is it correct to say that 85% of all fully grown Indiana grass snakes have length at least ℓ ?

Solution: Let X be he length; $X \sim \mathcal{N}(20, 16)$. We want x so that $\Pr(X > x) = .85$. This is the same as $\Pr(Z > (x - 20)/4) = .85$. From a standard normal table, $\Pr(Z < 1.04) = .85$, so by symmetry $\Pr(Z > -1.04) = .85$. So we want (x - 20)/4 = -1.04, or x = 15.84.

2. The Potawatami Zoo has 9 fully grown Indiana grass snakes. What is the probability that the average length of these 9 snakes is at least 18cm?

Solution: Let X_1 be the length of the *i*th snakes; each $X_i \sim \mathcal{N}(20, 16)$, and the X_i 's are independent. So $X_1 + \ldots + X_9 \sim \mathcal{N}(9 \times 20, 9 \times 16) = \mathcal{N}(180, 144)$ (note that this is an exact statement, not requiring Central Limit Theorem). We want to know about the average of the 9 lengths, which is $(X_1 + \ldots + X_9)/9 \sim \mathcal{N}(180/9, 144/81) = \mathcal{N}(20, 16/9)$. We want the probability that this is at least 18, which is $\Pr(Z > -1.5) = .9332$.

.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	,9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997

Entry is area A under the standard normal curve from $-\infty$ to z(A)

-

.9997

.9997

.9997

.9997

3.4

.9997

.9997

.9997

.9998

.9997

.9997