

The independent set sequence of regular bipartite graphs

David Galvin*

August 30, 2010

Abstract

Let $i_t(G)$ be the number of independent sets of size t in a graph G . Alavi, Erdős, Malde and Schwenk made the conjecture that if G is a tree then the independent set sequence $\{i_t(G)\}_{t \geq 0}$ of G is unimodal; Levit and Mandrescu further conjectured that this should hold for all bipartite G .

We consider the independent set sequence of finite regular bipartite graphs, and graphs obtained from these by percolation (independent deletion of edges). Using bounds on the independent set polynomial $P(G, \lambda) := \sum_{t \geq 0} i_t(G) \lambda^t$ for these graphs, we obtain partial unimodality results in these cases.

We then focus on the discrete hypercube Q_d , the graph on vertex set $\{0, 1\}^d$ with two strings adjacent if they differ on exactly one coordinate. We obtain asymptotically tight estimates for $i_{t(d)}(Q_d)$ in the case where $\lim t(d)/2^{d-1}$ is greater than $1 - 1/\sqrt{2}$, and nearly matching upper and lower bounds otherwise. We use these estimates to obtain a stronger partial unimodality result for the independent set sequence of Q_d .

1 Introduction and statement of results

For a (finite, simple, undirected, loopless) graph $G = (V, E)$ set

$$i_t(G) = \{I \in \mathcal{I}(G) : |I| = t\}$$

where $\mathcal{I}(G)$ is the collection of independent sets of G (sets of vertices spanning no edges). The *independent set sequence* of G is the sequence $\{i_t(G)\}_{t=0}^{\alpha(G)}$ where $\alpha(G)$

2010 Mathematics Subject Classification: 05C69 (Primary), 05C30 (Secondary)

Key words and phrases: independent set polynomial, stable set polynomial, regular graph, discrete hypercube, unimodal sequence

*Department of Mathematics, University of Notre Dame, 255 Hurley Hall, Notre Dame IN 46556; dgalvin1@nd.edu. Research supported in part by National Security Agency grant H98230-10-1-0364.

is the size of the largest independent set in G . The *independent set polynomial* or *stable set polynomial* of G , first introduced explicitly by Gutman and Harary [6], is the polynomial

$$P(G, \lambda) = \sum_{t=0}^{\alpha(G)} i_t(G) \lambda^t.$$

A sequence $\{a_i\}_{i=0}^n$ is said to be *unimodal* (with *mode* k) if

$$a_0 \leq a_1 \leq \dots \leq a_k \geq a_{k+1} \geq \dots \geq a_n,$$

and the polynomial $\sum_{i=0}^n a_i \lambda^i$ is said to be unimodal if its sequence of coefficients is. There has been some consideration in the literature of the question of the unimodality of the independent set polynomial of a graph. There are two major positive results. The first of these follows from the celebrated result of Heilmann and Lieb [9] to the effect that for any graph G the matching polynomial $\sum_{i \geq 0} m_i(G) \lambda^i$ of G (where $m_t(G)$ is the number of matchings of G of size t) has only real roots, implying by Newton's inequality (see for example [8]) that it is unimodal (in fact, log-concave). Since independent sets of a fixed size in the line graph of a graph are in bijection with matchings of that size in the original graph, Heilmann and Lieb's result shows that if G is the line graph of a graph, then $P(G, \lambda)$ is unimodal.

The second of these positive results is due Hamidoune [7], who showed that if G is claw-free (that is, does not contain a star on four vertices as an induced subgraph) then $P(G, \lambda)$ is unimodal. Later Chudnovsky and Seymour [3] showed that in this case $P(G, \lambda)$ also has all real roots. Since line graphs are claw-free, this result generalizes that of Heilmann and Lieb.

On the other hand, if G contains a claw then $P(G, \lambda)$ may not be unimodal. For example, the graph obtained from a claw by replacing each of the degree 1 vertices with a K_4 (the complete graph on 4 vertices), the degree 3 vertex with a K_{37} , and all edges with complete bipartite graphs, has independent set polynomial $1 + 49\lambda + 48\lambda^2 + 64\lambda^3$. In fact, Alvari, Erdős, Malde and Schwenk [1] showed that in general the independent set polynomial of a graph can have every possible pattern of increases and decreases. Specifically, they showed that for any integer $m \geq 1$ and any permutation π of $\{1, \dots, m\}$, there is a graph G with $\alpha(G) = m$ and with

$$i_{\pi(1)}(G) < i_{\pi(2)}(G) < \dots < i_{\pi(m)}(G). \quad (1)$$

In the language of [1], the independent set polynomial of a general graph is *unconstrained*. Note that there are at most $2^m = o(m!)$ permutations π of $\{1, \dots, m\}$ for which (1) holds for some graph G satisfying $\alpha(G) = m$ and $P(G, \lambda)$ unimodal.

Alvari, Erdős, Malde and Schwenk made the following positive conjecture.

Conjecture 1.1. *If T is a tree then $P(T, \lambda)$ is unimodal.*

In [11] Levit and Mandrescu make the stronger conjecture that $P(G, \lambda)$ is unimodal if G is a König-Egerváry graph (a graph in which the size of the largest independent set plus the size of the largest matching equals the number of vertices in the graph). In particular, since all bipartite graphs are König-Egerváry graphs, we have the following:

Conjecture 1.2. *If G is bipartite then $P(G, \lambda)$ is unimodal.*

Very little progress has been made towards this conjecture; indeed, even Conjecture 1.1 remains open. A partial result of Levit and Mandrescu [11] is that for G a bipartite graph, the final third of the coefficients of $P(G, \lambda)$ form a decreasing sequence, that is,

$$i_{\lceil (2\alpha(G)-1)/3 \rceil}(G) \geq i_{\lceil (2\alpha(G)-1)/3 \rceil + 1}(G) \geq \dots \geq i_{\alpha(G)}(G). \quad (2)$$

In this note we consider the independent set polynomial of a graph drawn from the family of regular and almost-regular bipartite graphs. One approach to showing unimodality of the independent set polynomial of a graph is to obtain upper and lower bounds on $i_t(G)$ for each $0 \leq t \leq \alpha(G)$, and this is the approach that we take here.

For d -regular G (not necessarily bipartite) we have the following upper bound on $i_t(G)$:

$$i_t(G) \leq \exp_2 \left\{ H \left(\frac{2t}{|V|} \right) \frac{|V|}{2} + \frac{|V|}{2d} \right\} \quad (3)$$

where $H(x) = -x \log x - (1-x) \log(1-x)$ is the binary entropy function. (Here and elsewhere $\log = \log_2$.) This is close to best possible; the graph consisting of disjoint copies of $K_{d,d}$, the complete bipartite graph with d vertices in each class, shows that $|V|/2d$ above cannot be replaced by $c|V|/d$ for any $c < 1/2$. In general this bound (whose derivation we postpone for a moment) may be far from the truth, but for d -regular bipartite G it has the same leading term in the exponent as the trivial lower bound

$$i_t(G) \geq \binom{|V|}{t} \geq \exp_2 \left\{ H \left(\frac{2t}{|V|} \right) \frac{|V|}{2} - \frac{1}{2} \log |V| \right\} \quad (4)$$

obtained by specifying a bipartition $V = \mathcal{E} \cup \mathcal{O}$ of G and considering only those independent sets which are subsets of \mathcal{E} . The second inequality in (4) follows for all $|V| \geq 1$ and $0 \leq t \leq |V|/2$ from Stirling's formula. (Note that for regular bipartite G , $\alpha(G) = |V(G)|/2$.)

The bound in (3) is based on the observation that for all $t > 0$ and $\lambda > 0$ we have $i_t(G)\lambda^t \leq P(G, \lambda)$ and so

$$i_t(G) \leq \min_{\lambda > 0} \left\{ \frac{P(G, \lambda)}{\lambda^t} \right\}.$$

From this we obtain (3) (at least for $t \neq 0, |V|/2$; but these two extreme cases are trivial) by taking $\lambda = \frac{2t}{|V(G)|-2t}$ in the following inequality, proved in [12]:

$$P(G, \lambda) \leq 2^{\frac{|V|}{2d}} (1 + \lambda)^{\frac{|V|}{2}}. \quad (5)$$

(The weaker bound with $|V|/d$ replacing $|V|/(2d)$ had earlier been obtained in [2]).

Inequalities (3) and (4) combine to show that for regular, bipartite G we have $i_t(G) \approx \binom{|V|/2}{t}$. This suggests that in contrast to the general situation, for regular bipartite G the independent set polynomial is quite constrained (in the language of [1]), in the sense that any sufficiently sparse subsequence of $\{i_t(G)\}_{t=0}^{|V(G)|/2}$ is unimodal.

To state a precise result in this direction, it will be helpful to set up some notation. Say that a sequence $\{a_i\}_{i \geq 0}$ is *s-step monotone increasing* on the interval $[a, b]$ (where a and b satisfy $0 \leq a \leq b$) if for every $a \leq i \leq j \leq b$ with $j - i \geq s$ we have $a_i \leq a_j$, and define *s-step monotone decreasing* analogously. (Note that $\{a_i\}_{i=0}^r$ being unimodal with mode k is equivalent to $\{a_i\}_{i=0}^r$ being 1-step monotone increasing on $[0, k]$ and 1-step monotone decreasing on $[k, r]$.) For a bipartite graph G on $2n$ vertices, say that G has *property* (β, γ, s) if the sequence $\{i_t(G)\}_{t \geq 0}$ is *s-step monotone increasing* on the interval $[\beta n, (1 - \gamma)n/2]$ and *s-step monotone decreasing* on the interval $[(1 + \gamma)n/2, (1 - \beta)n]$. Property $(0, 0, 1)$ is equivalent to the unimodality of $\{i_t(G)\}_{t=0}^n$ with mode $n/2$. We prove a number of results establishing property (β, γ, s) for β and γ arbitrarily small constants and $s = o(n)$.

Theorem 1.3. *Fix $\varepsilon > 0$. There is $C = C(\varepsilon) > 0$ such that if G is a $2n$ -vertex, d -regular bipartite graph, then G has property $(0, \varepsilon, s)$ with*

$$s = C \max \left\{ \log n, \frac{n}{d} \right\}.$$

Note that if $d = \omega(1)$ in the above theorem, then $s = o(n)$.

The condition that G be regular can be relaxed quite a bit; using a recent result in the spirit of (5) obtained in [4] we may extend Theorem 1.3 to all bipartite graphs which are suitably close to regular. Specifically, let G be a bipartite graph on $2n$ vertices with bipartition classes \mathcal{E} and \mathcal{O} (with $|\mathcal{O}| \geq |\mathcal{E}|$), let d be an arbitrary positive parameter and let

$$h(G, d) = \frac{1}{d} + \frac{|\{v \in \mathcal{E} : d(v) < d\}|}{n} + \frac{1}{dn} \sum_{v \in \mathcal{O}} (d(v) - d) \mathbf{1}_{\{d(v) \geq d\}} + \frac{|\mathcal{O}| - |\mathcal{E}|}{n}$$

(where $\mathbf{1}_E$ is the indicator function of the event E). The parameter $h(G, d)$ is intended to capture the degree to which G is “almost d -regular”: $h(G, d)$ being close to $1/d$ (the value it takes when G is d -regular) means that G has not too many low degree vertices, that the sum of the degrees of high degree vertices is not too large, and that the difference between the sizes of the partition classes is not too great.

Similar to Theorem 1.3, we have the following statement:

Theorem 1.4. *Fix $\varepsilon > 0$. There is $C = C(\varepsilon) > 0$ such that if G is a $2n$ -vertex bipartite graph and d is an arbitrary positive parameter, then G has property $(\varepsilon, \varepsilon, s)$ with*

$$s = C \max \{ \log n, nh(G, d) \}.$$

This theorem is only of interest when $s = o(n)$, that is, when there is a choice of d for which $h(G, d) = o(1)$. One quite general situation in which this occurs is in percolation. Given a graph G and a parameter $0 \leq p \leq 1$, let G^p be a random subgraph of G obtained by deleting each edge independently with probability $1 - p$ (so

$$\Pr(G^p = H) = p^{|E(H)|} (1 - p)^{|E(G)| - |E(H)|}.$$

In [4, Section 4] it is shown that if G is a d -regular bipartite graph on $2n$ vertices, and G^p is obtained from G by percolation with $p \geq f(d)/d$ for some function $f(d) = \omega(1)$, then there is a function $g(d) = o(1)$ such that with probability at least $1 - g(d)$, we have that $h(G^p, d') \leq g(d)$ (for $d' = dp - (2dp)^{1/2} f(d)^{1/4}$). We thus have the following corollary of Theorem 1.4.

Corollary 1.5. *Fix $\varepsilon > 0$. Let G be a d -regular bipartite graph on $2n$ vertices. Let $p = f(d)/d$ for some function $f(d) = \omega(1)$. There are functions $g(d) = o(1)$ and $s(n, d) = o(n)$ such that with probability at least $1 - g(d)$, G^p has property $(\varepsilon, \varepsilon, s(n, d))$.*

A particularly interesting application of Corollary 1.5 is to the random equi-bipartite graph $G(n, n, p)$. This is the graph on vertex set $\mathcal{E} \cup \mathcal{O}$ with $|\mathcal{E}| = |\mathcal{O}| = n$ in which the edge $\{u, v\}$ ($u \in \mathcal{E}, v \in \mathcal{O}$) is present with probability p , independently for all choices of u and v . (In other words, $G(n, n, p)$ is the result of percolation on $K_{n, n}$ with parameter p .)

Corollary 1.6. *Fix $\varepsilon > 0$. Let $p = \omega(1)/n$. There is a function $s(n) = o(n)$ so that almost surely (with probability tending to 1 as n goes to infinity) the random bipartite graph $G(n, n, p)$ has property $(\varepsilon, \varepsilon, s(n))$.*

In other words, almost all equi-bipartite graphs exhibit partial unimodality of the independent set sequence.

The bound in (5) is valid for *all* d -regular graphs, and we expect that it, as well as both the upper and lower bounds on $i_t(G)$ ((3) and (4)), can be significantly improved if further structural conditions are put on G . In [5] the programme of improving (5) is carried out in the case when G is the discrete hypercube Q_d . This is the graph on vertex set $V = \{0, 1\}^d$ with two strings adjacent if they differ on exactly one coordinate. It is a d -regular bipartite graph with bipartition classes \mathcal{E} and \mathcal{O} , where \mathcal{E} is the set of vertices with an even number of 1's. Note that $|\mathcal{E}| = |\mathcal{O}| = \alpha(Q_d) = 2^{d-1}$. The following bound is obtained in [5]. (All asymptotic statements in what follows are as $d \rightarrow \infty$.)

Theorem 1.7. *There is a $c > 0$ and a function $f(d) \rightarrow 0$ such that for $\lambda > \frac{c \log d}{d^{1/3}}$,*

$$P(Q_d, \lambda) = 2(1 + \lambda)^{2^{d-1}} \exp \left\{ \frac{\lambda}{2} \left(\frac{2}{1 + \lambda} \right)^d (1 + f(d)) \right\}.$$

This generalizes work of Korshunov and Sapozhenko [10], who had shown

$$P(Q_d, 1) = (2\sqrt{e} + o(1))2^{2^{d-1}}.$$

Using Theorem 1.7 (or rather, using two of the intermediate inequalities that ultimately lead to the theorem), we can significantly improve the bounds on $i_t(Q_d)$ given by (3) and (4), obtaining optimal (asymptotically matching) bounds for a large range of values of t .

Theorem 1.8. *There is a constant $c > 0$ and a function $f(d) \rightarrow 0$ such that if $t = t(d)$ satisfies*

$$2^{d-1} \left(\frac{c \log d}{d^{1/3}} \right) \leq t \leq 2^{d-1} \left(1 - \frac{1}{\sqrt{2}} + \frac{2 \log d}{d} \right) \quad (6)$$

for all sufficiently large d then

$$i_t(Q_d) = \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} (1 + f(d)) \right\}.$$

If t satisfies

$$2^{d-1} \left(1 - \frac{1}{\sqrt{2}} + \frac{2 \log d}{d} \right) \leq t \leq 2^{d-1} \quad (7)$$

for all sufficiently large d then

$$i_t(Q_d) \sim 2 \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right\}.$$

A corollary of Theorem 1.8 is that the quantity $i_t(Q_d)$ undergoes a transition around $t = 2^{d-2}$, in a window of width $O(1/d)$. This is analogous to [5, Corollary 1.2].

Corollary 1.9. *If $t = t(d) = 2^{d-1} \left(\frac{1}{2} + \frac{g(d)}{d} \right)$ then*

$$\frac{i_t(Q_d)}{2 \binom{2^{d-1}}{t}} \rightarrow \begin{cases} \infty & \text{if } g(d) \rightarrow -\infty \\ \exp \{ e^{-2k}/2 \} & \text{if } g(d) \rightarrow k, \text{ a constant} \\ 1 & \text{if } g(d) \rightarrow +\infty \end{cases}$$

as $d \rightarrow \infty$.

We now turn to considering the unimodality of $P(Q_d, \lambda)$. By direct counting this polynomial may be shown to be unimodal for all $d \leq 5$ (although for $3 \leq d \leq 5$ it has some complex roots). For larger d , a corollary of Theorem 1.3 is that for all $\varepsilon > 0$ there is a constant $C = C(\varepsilon) > 0$ such that Q_d has property $(0, \varepsilon, C2^d/d)$. By examining the error terms in the asymptotic estimates of $i_t(Q_d)$ provided by Theorem 1.8, we are able to obtain much stronger partial unimodality result for $P(Q_d, \lambda)$, and in particular extend Levit and Mandrescu's observation (2) (in the particular case of $G = Q_d$) to a wider range of coefficients.

Theorem 1.10. *For all sufficiently large d it holds that*

$$i_p(Q_d) < i_{p+1}(Q_d) < \dots < i_{2^{d-2}-15d^2}(Q_d)$$

where $p = \lceil (1 - 1/\sqrt{2} + 2 \log d/d)2^{d-1} \rceil$, and

$$i_{2^{d-2}+5d^4}(Q_d) > i_{2^{d-2}+5d^4+1}(Q_d) > \dots > i_{2^{d-1}}(Q_d).$$

We have not made an attempt to optimize the coefficients of d^2 and d^4 here. The proof strategy is to show that for $p \leq t \leq t+1 \leq 2^{d-2} - 15d^2$ our upper bound on $i_t(Q_d)$ from Theorem 1.8 is eventually smaller than our lower bound on $i_{t+1}(Q_d)$ (with a similar approach for $2^{d-2} + 5d^4 \leq t \leq t+1 \leq 2^{d-1}$). Below p the upper and lower bounds on $i_t(Q_d)$ provided by Theorem 1.8 are too far apart to be of any use, and in the neighbourhood of $t = 2^{d-2}$ the sequence $\{i_t(Q_d)\}$ seems to be too flat for the present approach to be helpful.

The remainder of this note is laid out as follows. In Section 2 we prove Theorems 1.3 and 1.4, while the proofs of Theorems 1.8 and 1.10 are given in Section 3.

2 Proofs of Theorems 1.3 and 1.4

We begin with the proof of Theorem 1.3. Let j and ℓ satisfy $0 \leq j \leq \ell \leq (1 - \varepsilon)n/2$. By (3) and (4), a sufficient condition for $i_\ell(G) > i_j(G)$ is

$$H\left(\frac{\ell}{n}\right) - H\left(\frac{j}{n}\right) > \frac{1}{d} + \frac{\log 2n}{2n}. \quad (8)$$

By the mean value theorem,

$$H\left(\frac{\ell}{n}\right) - H\left(\frac{j}{n}\right) = \left(\frac{\ell}{n} - \frac{j}{n}\right) H'(\xi)$$

for some $\xi \in (\ell/n, j/n)$. Since the minimum of $H'(x)$ on $[0, (1 - \varepsilon)/2]$ is achieved at $x = (1 - \varepsilon)/2$ and is a positive constant depending on ε , we get that a sufficient condition for $i_\ell(G) > i_j(G)$ is

$$\ell - j \geq C(\varepsilon) \left(\frac{n}{d} + \log 2n\right).$$

For j and ℓ satisfying $(1 - \varepsilon)n/2 \leq \ell \leq j \leq n/2$ the proof is almost identical, once we observe that $H(x)$ is symmetric around $x = 1/2$.

In order to prove Theorem 1.4, we need an analog of (3) for the family of G 's under consideration in that theorem. The following result is a special case of a general result on graph homomorphisms from [4, Section 3]. For $\lambda_1, \lambda_2 > 1$,

$$\sum_{t=0}^{2n} i_t(G) \lambda_1^t \lambda_2^{2n-t} \leq (\lambda_1 \lambda_2 + \lambda_2^2)^n C(\lambda_1, \lambda_2)^{nh(G,d)} \quad (9)$$

where $C(\lambda_1, \lambda_2)$ may be taken to be $\max\{256, \lambda_1 + \lambda_2\}$. It follows that for all fixed $\lambda > 0$,

$$P(G, \lambda) \leq (1 + \lambda)^n C(\lambda)^{nh(G,d)} \quad (10)$$

where $C(\lambda) > 0$ goes to infinity both as λ goes to 0 and as λ goes to infinity. Indeed, if $\lambda \geq 1$ then we may take $\lambda_1 = 2\lambda$ and $\lambda_2 = 2$ in (9) to get

$$P(G, \lambda) = \lambda_2^{-2n} \sum_{t=0}^{2n} i_t(G) (\lambda_1/\lambda_2)^t \leq (1 + \lambda)^n \max\{256, 2(1 + \lambda)\}^{nh(G,d)}$$

while if $\lambda < 1$ then we may take $\lambda_1 = 2$ and $\lambda_2 = 2/\lambda$ to get

$$P(G, \lambda) \leq (1 + \lambda)^n \max\{256, 2(1 + 1/\lambda)\}^{nh(G,d)},$$

showing that we may take $C(\lambda)$ to be $2(1 + \lambda)$ for $\lambda \geq 127$, to be 256 for $1/127 \leq \lambda \leq 127$, and to be $2(1 + 1/\lambda)$ for $\lambda \leq 1/127$.

Using (10) in place of (5), we may reproduce the derivation of (3) to obtain the following upper bound on $i_t(G)$ for G satisfying the conditions of Theorem 1.4:

$$i_t(G) \leq \exp_2 \left\{ H\left(\frac{t}{n}\right) n + C(t, n) nh(G, d) \right\}$$

where $C(t, n) > 0$ may be taken as follows:

$$C(t, n) = \begin{cases} \log\left(\frac{2n}{n-t}\right) & \text{if } \frac{127n}{128} \leq t \\ 8 & \text{if } \frac{n}{128} \leq t \leq \frac{127n}{128} \\ \log\left(\frac{2n}{t}\right) & \text{if } t \leq \frac{n}{128}. \end{cases}$$

It follows that for $\varepsilon > 0$ there is $C(\varepsilon) > 0$ such that in the range $t \in [\varepsilon n, (1 - \varepsilon)n]$ we have

$$i_t(G) \leq \exp_2 \left\{ H\left(\frac{t}{n}\right) n + C(\varepsilon) nh(G, d) \right\}. \quad (11)$$

Noting that (4) is still valid in the setting of Theorem 1.4, it follows from (11) (just as (8) followed from (3)) that for all $\varepsilon > 0$, if j and ℓ satisfy $\varepsilon n \leq j \leq \ell \leq (1 - \varepsilon)n/2$, then a sufficient condition for $i_\ell(G) > i_j(G)$ is

$$H\left(\frac{\ell}{n}\right) - H\left(\frac{j}{n}\right) > C(\varepsilon) h(G, d) + \frac{\log 2n}{2n}.$$

The proof of Theorem 1.4 now goes through exactly as the proof of Theorem 1.3.

3 Proofs of Theorems 1.8 and 1.10

3.1 Preliminaries

We begin with some notation. For $A \subseteq V (= \{0, 1\}^d)$ write $N(A)$ for the set of vertices outside A that are neighbours of a vertex in A and set

$$[A] = \{v \in V : N(\{v\}) \subseteq N(A)\}.$$

Note that if A is an independent set then $A \subseteq [A]$. Say that $A \subseteq \mathcal{E}$ (or \mathcal{O}) is *small* if $|[A]| \leq 2^{d-2}$ and *2-linked* if $A \cup N(A)$ induces a connected subgraph of Q_d . Any A can be decomposed into its maximal 2-linked subsets; we refer to these as the *2-components* of A , and write $k(A)$ for the number of 2-components of A and $\text{cl}(A)$ for the size of the largest 2-component of A . Finally, for all $\lambda > 0$ and $a, g \geq 0$ we define a function $F_\lambda(a, g)$ by

$$F_\lambda(a, g) = \lambda^a (1 + \lambda)^{-g}.$$

There are two bounds from [5] (intermediate steps in the derivation of Theorem 1.7) that will be of use here. Both are valid for all $\lambda > \frac{c \log d}{d^{1/3}}$ for some absolute constant $c > 0$. First, we have

$$\sum_{A \subseteq \mathcal{E} \text{ small}} F_\lambda(|A|, |N(A)|) \leq \exp \left\{ \frac{\lambda}{2} \left(\frac{2}{1 + \lambda} \right)^d + \frac{d^2 \lambda^2 (1 + \lambda)^{2d}}{(1 + \lambda)^{2d}} \right\} \quad (12)$$

(this is [5, Equation (23)]). Second, for fixed $k \geq 1$ we have

$$\sum_{A \subseteq \mathcal{E} \text{ small, 2-linked, } |A| \geq k} F_\lambda(a, g) \leq e^{k-1} d^{2k-2} 2^d F_\lambda(k, kd - 2k(k-1)) \quad (13)$$

(this is [5, Corollary 3.11]).

3.2 Proof of Theorem 1.8

We assume throughout that

$$t \geq \left(\frac{c \log d}{d^{1/3}} \right) 2^{d-1}$$

with c the same as the constant appearing in the range of validity of (12) and (13). We also, where necessary, assume that d is large enough to support our assertions. All asymptotic statements in what follows will be as $d \rightarrow \infty$.

For the lower bounds, we may assume $t \leq \frac{3}{4}2^{d-1}$ (any constant greater than $1/2$ would do in place of $3/4$ here) since for $t \geq \frac{3}{4}2^{d-1}$, $t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} = o(1)$ and so the bound

$$i_t(Q_d) \geq (2 - o(1)) \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} \right\}$$

is trivial. For the remaining range of t , let $f = f(t, d)$ be defined by

$$f = \max \left\{ d, 5^7 e t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} \right\}. \quad (14)$$

The lower bounds will be based on

$$\begin{aligned} i_t(Q_d) &\geq \sum_{A \subseteq \mathcal{E} \text{ or } A \subseteq \mathcal{O}, \text{cl}(A) \leq 1, |A| \leq f} \binom{2^{d-1} - |N(A)|}{t - |A|} \\ &= 2 \sum_{A \subseteq \mathcal{E}, \text{cl}(A) \leq 1, |A| \leq f} \binom{2^{d-1} - d|A|}{t - |A|}. \end{aligned} \quad (15)$$

This lower bound is obtained by first choosing a set of vertices of size no more than f , no two of which share a common neighbour, to be the intersection of the independent set with one of the two partition classes, and then extending it by choosing an arbitrary subset of the other class of appropriate size (note that for A with $\text{cl}(A) = 1$, $|N(A)| = d|A|$). The equality (15) is valid since $f < t/2$ and so there is no overlap between the independent sets of size t which intersect \mathcal{E} in no more than f vertices and those which intersect \mathcal{O} in no more than f vertices.

For all a and g with $g \geq a$, and $0 \leq t \leq 2^{d-1}$, we have the identity

$$\begin{aligned} \binom{2^{d-1} - g}{t - a} &= \binom{2^{d-1}}{t} \frac{t^a (2^{d-1} - t)^{g-a}}{(2^{d-1})^g} E(a, g) \\ &= F_{\lambda(t)}(a, g) \binom{2^{d-1}}{t} E(a, g) \end{aligned} \quad (16)$$

where $\lambda(t) = t/(2^{d-1} - t)$ and

$$E(a, g) = \frac{\prod_{i=0}^{a-1} \left(1 - \frac{i}{t}\right) \prod_{i=0}^{g-a-1} \left(1 - \frac{i}{2^{d-1}-t}\right)}{\prod_{i=0}^{g-1} \left(1 - \frac{i}{2^{d-1}}\right)}.$$

Note that

$$F_{\lambda(t)}(k, dk) = \left(\frac{t}{2^{d-1}} \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} \right)^k, \quad (17)$$

and that for fixed d and k , the quantity on the right-hand side in (17) is decreasing in t for the range of t that we are considering (indeed, it is decreasing for $t \geq (2^{d-1} - 1)/(d - 1)$), a fact that we will use repeatedly in the calculations that follow, usually without comment. For those A contributing to (15),

$$\begin{aligned} E(|A|, d|A|) &\geq \prod_{i=0}^{f-1} \left(1 - \frac{i}{t}\right) \prod_{i=0}^{(d-1)f-1} \left(1 - \frac{i}{2^{d-1} - t}\right) \\ &\geq \exp \left\{ -\frac{f^2}{t} - \frac{d^2 f^2}{2^{d-1} - t} \right\} \\ &\geq \exp \left\{ -\frac{2f^2}{t} \right\}. \end{aligned} \tag{18}$$

In the second inequality we use $1 - x \geq e^{-2x}$ for $0 \leq x \leq 1/2$. The use is valid since both $f \leq t/2$ and $f \leq (2^{d-1} - t)/(2d)$ hold.

The number of ways of choosing $A \subseteq \mathcal{E}$ with $\text{cl}(A) \leq 1$ and $|A| = k \leq f$ is at least

$$\begin{aligned} \frac{\prod_{i=0}^{k-1} (2^{d-1} - id^2)}{k!} &\geq \frac{(2^{d-1})^k}{k!} \exp \left\{ -\frac{f^2 d^2}{2^{d-1}} \right\} \\ &\geq \frac{(2^{d-1})^k}{k!} \exp \left\{ -\frac{f^2}{t} \right\} \end{aligned} \tag{19}$$

since each choice of vertex in A eliminates from consideration at most d^2 other vertices. By (16) each such A contributes

$$F_{\lambda(t)}(k, dk) \binom{2^{d-1}}{t} E(|A|, d|A|) \tag{20}$$

to the sum in (15). Combining (17), (18), (19) and (20) with (15) we get

$$i_t(Q_d) \geq 2 \binom{2^{d-1}}{t} \exp \left\{ -\frac{3f^2}{t} \right\} \sum_{k \leq f} \frac{1}{k!} \left(t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right)^k. \tag{21}$$

Using $k! \geq (k/e)^k$ and the lower bound on f from (14) we have

$$\sum_{k \leq f} \frac{1}{k!} \left(t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right)^k \geq E_0 \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right\}$$

where

$$E_0 = 1 - 2 \left(\frac{et \left(1 - \frac{t}{2^{d-1}} \right)^{d-1}}{f} \right)^f \exp \left\{ -t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right\}$$

and so (21) becomes

$$i_t(Q_d) \geq 2 \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right\} E_0 \quad (22)$$

where

$$E_1 = \exp \left\{ -\frac{3f^2}{t} \right\} E_0.$$

The dominating term here is $\exp\{-3f^2/t\}$. For t satisfying (6)

$$E_1 \geq \exp \left\{ -o \left(t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right) \right\}$$

and for t satisfying (7) $E_1 \geq 1 - o(1)$, completing the lower bounds. We also note the following more precise bounds:

$$E_1 \geq \begin{cases} 1 - \frac{1}{d^5} & \text{if } t \text{ satisfies (7)} \\ 1 - \frac{14d^2}{2^d} & \text{if } 2^{d-1} \left(\frac{1}{2} - \frac{1}{d} \right) \leq t \leq \frac{3}{4} 2^{d-1}. \end{cases} \quad (23)$$

We now turn to the upper bounds. For any $I \in \mathcal{I}(Q_d)$ we have $[I \cap \mathcal{E}] \cap [I \cap \mathcal{O}] = \emptyset$. Since Q_d has a perfect matching, it follows that at least one of $[I \cap \mathcal{E}]$, $[I \cap \mathcal{O}]$ is no larger than 2^{d-2} , that is, that at least one of $I \cap \mathcal{E}$, $I \cap \mathcal{O}$ is small. This together with \mathcal{E} - \mathcal{O} symmetry leads to the bound

$$i_t(Q_d) \leq 2 \sum_{A \subseteq \mathcal{E} \text{ small}} \binom{2^{d-1} - |N(A)|}{t - |A|}. \quad (24)$$

We will examine the sum in (24) in three parts. Say that small $A \subseteq \mathcal{E}$ is of *type I* if $|A| \leq f$, of *type II* if $|A| > f$ and $\text{cl}(A) \leq 5$, and of *type III* if $|A| > f$ and $\text{cl}(A) \geq 6$ (where f is as defined in (14)).

We first consider the contribution to the sum in (24) from A of type I. For these A we have

$$E(|A|, |N(A)|) \leq \exp \left\{ \frac{d^2 f^2}{2^{d-1}} \right\}$$

(this is similar to the derivation of (18), except that in this case we are lower bounding the numerator of $E(|A|, |N(A)|)$). Taking $\lambda = \lambda(t) = t/(2^{d-1} - t)$ in (12) and combining with (16) we find that the contribution to (24) from A of type I is at most

$$2 \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} + \frac{d^2 t^2 2^d}{(2^{d-1} - t)^2} \left(1 - \frac{t}{2^{d-1}} \right)^{2d-2} + \frac{d^2 f^2}{2^{d-1}} \right\}. \quad (25)$$

Next we consider the sum in (24) over A of type II. We have

$$\begin{aligned}
\binom{2^{d-1} - g}{t - a} &= \lambda(t)^{a-t} \binom{2^{d-1} - g}{t - a} \lambda(t)^{t-a} \\
&\leq \lambda(t)^{a-t} (1 + \lambda(t))^{2^{d-1}-g} \\
&= (1 + \lambda(t))^{2^{d-1}} \lambda(t)^{-t} F_{\lambda(t)}(a, g) \\
&= 2^{H(\frac{t}{2^{d-1}})} 2^{d-1} F_{\lambda(t)}(a, g).
\end{aligned}$$

By Stirling's formula, (more precisely, by the fact that for all $n \geq 1$,

$$2n^n e^{-n} \sqrt{n} \leq n! \leq 3n^n e^{-n} \sqrt{n},$$

this is at most $3 \cdot 2^{d/2} \binom{2^{d-1}}{t} F_{\lambda(t)}(a, g)$. It follows that the contribution to (24) from A of type II is at most

$$6 \binom{2^{d/2}}{t} \sum_{A \subseteq \mathcal{E} \text{ small, } \text{cl}(A) \leq 5, k(A) \geq f/5} F_{\lambda(t)}(|A|, |N(A)|). \quad (26)$$

To bound this sum, we make a number of observations.

- For each $k \geq f/5$ there are at most $2^{k(d-1)}/k!$ ways of choosing a fixed vertex in each of the k 2-components of A , and at most 5^k ways of assigning a size to each 2-component.
- For each $\ell = 1, \dots, 5$, the number of 2-linked subsets of \mathcal{E} of size ℓ that include a fixed vertex is at most $(\ell - 1)!(d^2)^{\ell-1}$. (Once j vertices have been chosen, there are at most $j d^2$ choices for the $(j + 1)$ st.)
- Each $A \subseteq \mathcal{E}$ with $|A| = \ell \leq 5$ satisfies $|N(A)| \geq d\ell - 2\ell(\ell - 1)$. (Each vertex of A has d neighbours, of which at least $d - 2(|A| - 1)$ must be unique to it, since a pair of vertices in Q_d can have at most two common neighbours.)
- The quantity $F_{\lambda}(a, g)$ is decreasing in g , and for each $\ell = 1, \dots, 5$, (and sufficiently large d)

$$(\ell - 1)!(d^2)^{\ell-1} F_{\lambda(t)}(\ell, d\ell - 2\ell(\ell - 1)) \leq F_{\lambda(t)}(1, d).$$

All this together serves to bound the expression in (26) by

$$6 \binom{2^{d/2}}{t} \sum_{k \geq f/5} \frac{1}{k!} \left(5t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right)^k \leq \frac{1}{3^f} \binom{2^{d-1}}{t}, \quad (27)$$

the inequality following from the choice of f and the bound $k! \geq (k/e)^k$.

Finally we consider the sum in (24) over A of type III. Beginning with the same steps as in the case of A of type II, this is at most

$$6 \binom{2^{d/2}}{t} \sum_{A \subseteq \mathcal{E} \text{ small, } \text{cl}(A) \geq 6} F_{\lambda(t)}(|A|, |N(A)|). \quad (28)$$

We now use a multiplicative property of F : for any λ , if A' is a 2-component of A then

$$F_{\lambda}(|A|, |N(A)|) = F_{\lambda}(|A'|, |N(A')|) F_{\lambda}(|A \setminus A'|, |N(A \setminus A')|)$$

and so the sum in (28) is at most

$$\left(\sum_{A \subseteq \mathcal{E} \text{ small, 2-linked, } |A| \geq 6} F_{\lambda(t)}(|A|, |N(A)|) \right) \left(\sum_{A \subseteq \mathcal{E} \text{ small}} F_{\lambda(t)}(|A|, |N(A)|) \right) = S_1 S_2.$$

By (12) we have

$$S_2 \leq \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} + \frac{d^2 t^2 2^d}{(2^{d-1} - t)^2} \left(1 - \frac{t}{2^{d-1}} \right)^{2d-2} \right\} \quad (29)$$

and by (13) we have

$$S_1 \leq e^5 d^{10} 2^d F_{\lambda(t)}(6, 6d - 60). \quad (30)$$

Combining (25), (27), (29) and (30) we find that

$$i_t(Q_d) \leq 2 \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right\} E_2 \quad (31)$$

where

$$E_2 = \exp \left\{ \frac{d^2 t^2 2^d}{(2^{d-1} - t)^2} \left(1 - \frac{t}{2^{d-1}} \right)^{2d-2} + \frac{d^2 f^2}{2^{d-1}} \right\} + \frac{1}{3f} + 3e^5 d^{10} 2^{3d/2} F_{\lambda(t)}(6, 6d - 60) \exp \left\{ \frac{d^2 t^2 2^d}{(2^{d-1} - t)^2} \left(1 - \frac{t}{2^{d-1}} \right)^{2d-2} \right\}.$$

For t satisfying (6)

$$E_2 \leq \exp \left\{ o \left(t \left(1 - \frac{t}{2^{d-1}} \right)^{d-1} \right) \right\}$$

and for t satisfying (7) $E_2 \leq 1 + o(1)$, completing the upper bounds. We also note the following more precise bounds:

$$E_2 \leq \begin{cases} 1 + \frac{1}{d^3} & \text{if } t \text{ satisfies (7)} \\ 1 + \frac{4d^4}{2^d} & \text{if } t \geq 2^{d-1} \left(\frac{1}{2} - \frac{1}{d} \right). \end{cases} \quad (32)$$

3.3 Proof of Theorem 1.10

Fix

$$2^{d-2} + 5d^4 \leq t < (3/4)2^{d-1}. \quad (33)$$

We will show that for large d (independent of t), $i_t(Q_d) > i_{t+1}(Q_d)$. (Note that by (2), for the purposes of proving Theorem 1.10 it is not necessary to consider larger t .) By (22), (23), (31) and (32) it is enough to show that

$$\frac{2 \left(1 - \frac{14d^2}{2^d}\right) \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} \right\}}{2 \left(1 + \frac{4d^4}{2^d}\right) \binom{2^{d-1}}{t+1} \exp \left\{ (t+1) \left(1 - \frac{t+1}{2^{d-1}}\right)^{d-1} \right\}} > 1.$$

Since $t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1}$ is decreasing in t , this inequality is implied by

$$\frac{\left(1 - \frac{14d^2}{2^d}\right) (t+1)}{\left(1 + \frac{4d^4}{2^d}\right) (2^{d-1} - t)} > 1$$

which holds for all t satisfying (33) (for large enough d , independent of t).

Next, fix

$$2^{d-1} \left(1 - \frac{1}{\sqrt{2}} + \frac{2 \log d}{d}\right) \leq t < 2^{d-1} \left(\frac{1}{2} - \frac{1}{d}\right). \quad (34)$$

In this range we wish to show that for large d (independent of t), $i_t(Q_d) < i_{t+1}(Q_d)$. Again by (22), (23), (31) and (32) it is enough to show that

$$\frac{2 \left(1 + \frac{1}{d^3}\right) \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} \right\}}{2 \left(1 - \frac{1}{d^5}\right) \binom{2^{d-1}}{t+1} \exp \left\{ (t+1) \left(1 - \frac{t+1}{2^{d-1}}\right)^{d-1} \right\}} < 1. \quad (35)$$

Writing $h(a, b)$ for $a(1 - b/2^{d-1})^{d-1}$ we have

$$\begin{aligned} h(t, t) - h(t+1, t+1) &\leq h(t, t) - h(t, t+1) \\ &= h(t, t) \left(1 - \left(1 - \frac{1}{2^{d-1} - t}\right)^{d-1}\right) \\ &\leq h(t, t) \left(\frac{2(d-1)}{2^{d-1} - t}\right) \end{aligned} \quad (36)$$

with (36) valid for sufficiently large d as long as $2^{d-1} - t \geq 2$. For t satisfying (34) we therefore have

$$\begin{aligned} \exp \{h(t, t) - h(t+1, t+1)\} &\leq \exp \left\{ h(t, t) \left(\frac{2(d-1)}{2^{d-1} - t}\right) \right\} \\ &\leq \exp \left\{ h(2^{d-3}, 2^{d-3}) \left(\frac{2(d-1)}{2^{d-1} - 2^{d-2}}\right) \right\} \\ &\leq 1 + .76^d \end{aligned}$$

(for large enough d) and so (35) in this range is implied by

$$\frac{\left(1 + \frac{2}{d^3}\right)(t+1)}{\left(1 - \frac{1}{d^5}\right)(2^{d-1} - t)} < 1$$

which holds for all t satisfying (34) (for large d , independent of t).

Finally, in the range

$$2^{d-1} \left(\frac{1}{2} - \frac{1}{d}\right) \leq t < 2^{d-2} - 15d^2 \quad (37)$$

we again wish to show that for large d (independent of t), $i_t(Q_d) < i_{t+1}(Q_d)$, and again by (22), (23), (31) and (32) it is enough to show that

$$\frac{2 \left(1 + \frac{4d^4}{2^d}\right) \binom{2^{d-1}}{t} \exp \left\{ t \left(1 - \frac{t}{2^{d-1}}\right)^{d-1} \right\}}{2 \left(1 - \frac{14d^2}{2^d}\right) \binom{2^{d-1}}{t+1} \exp \left\{ (t+1) \left(1 - \frac{t+1}{2^{d-1}}\right)^{d-1} \right\}} < 1. \quad (38)$$

For t in this range we use (36) to obtain

$$\exp \{h(t, t) - h(t+1, t+1)\} \leq 1 + \frac{d^2}{2^d}$$

and so (38) is implied by

$$\frac{\left(1 + \frac{5d^4}{2^d}\right)(t+1)}{\left(1 - \frac{14d^2}{2^d}\right)(2^{d-1} - t)} < 1$$

which holds for all t satisfying (37) (for large d , independent of t).

Acknowledgement: Part of this work was carried while the author was a participant in the programme on Combinatorics and Statistical Mechanics at the Isaac Newton Institute for Mathematical Sciences, University of Cambridge, in spring 2008. The author thanks the institute and the programme organizers for the support provided.

References

- [1] Y. Alavi, P. Erdős, P. Malde and A. Schwenk, The vertex independence sequence of a graph is not constrained, *Congressus Numerantium* **58** (1987), 15–23.
- [2] T. Carroll, D. Galvin and P. Tetali, Matchings and Independent Sets of a Fixed Size in Regular Graphs, *J. Combin. Theory Ser. A* **116** (2009), 1219–1227.

- [3] M. Chudnovsky and P. Seymour, The Roots of The Stable Set Polynomial of a Claw-free Graph *J. Combin. Theory. Ser. B* **97** (2007), 350–357.
- [4] J. Engbers and D. Galvin, H -colouring bipartite graphs, submitted. (Preprint available at <http://www.nd.edu/~dgalvin1/pdf/colourings.pdf>.)
- [5] D. Galvin, A threshold phenomenon for random independent sets in the discrete hypercube, *Combin. Probat. Comput.*, in press. (Available at <http://www.nd.edu/~dgalvin1/pdf/Qdthreshold.pdf>.)
- [6] I. Gutman and F. Harary, Generalizations of the matching polynomial, *Utilitas Mathematica* **24** (1983), 97–106.
- [7] Y. Hamidoune, On the number of independent k -sets in a claw-free graph, *J. Combin. Theory B* **50** (1990), 241–244.
- [8] G. Hardy, J. Littlewood, and G. Pólya, *Inequalities* (2nd ed.), Cambridge University Press, Cambridge, 1952.
- [9] O. Heilmann and E. Lieb, Theory of monomer-dimer systems, *Comm. Math. Phys.* **25** (1972), 190–232.
- [10] A. Korshunov and A. Sapozhenko, The number of binary codes with distance 2, *Problemy Kibernet.* **40** (1983), 111–130. (Russian)
- [11] V. Levit and E. Mandrescu, Partial unimodality for independence polynomials of König-Egerváry graphs, *Congr. Numer.* **179** (2006), 109–119.
- [12] Y. Zhao, The Number of Independent Sets in a Regular Graph, *Combin. Probab. and Comput.* **19** (2010), 315–320.