

Toric Poisson Structures

AMS Eastern Sectional Meeting

Pennsylvania State University

Oct. 24, 2009

Arlo Caine

arlo.caine@nd.edu

arXiv:0910.0229

Goal of the construction

Desire: Π_Σ should be not-symplectic, yet non-degenerate on an open dense set, and have symmetry.

(M, π)	$\pi = 0$	$(\mathfrak{g}^*, \pi_{\mathfrak{g}^*})$	$(K/T, \pi_{K/T})$	π non-deg.
$S(M)$	points	Co-Ad orbits	Bruhat Cells	one leaf = M
	reg.	not reg.	not reg.	reg.

- Flag manifold K/T

KKS symplectic structure $\leftarrow K/T \rightarrow$ Bruhat Poisson structure


- Smooth toric variety $X(\Sigma)$

Delzant symplectic structures $\leftarrow X(\Sigma) \rightarrow$ Toric Poisson structure

Symplectic vs. Algebraic Geometry


Symplectic Geometry:

- $\Delta \subset \mathfrak{t}^*$ Delzant polytope with d facets.
 $(X_\Delta, \omega_\Delta)$ = symplectic reduction of $(\mathbb{C}^d, \omega_{\text{std}})$
with respect to action of $N \triangleleft \mathbb{T}^d$.
- Torus N defined by “shape” of Δ .
- Level determined by “size” of Δ .



Algebraic Geometry:

- Σ = the dual fan of Δ (“shape”)
- Smooth \mathbb{C} -variety $X(\Sigma) = \mathbb{C}^d // N_{\mathbb{C}}$ with action
of $T_{\mathbb{C}} \simeq \mathbb{T}_{\mathbb{C}}^d / N_{\mathbb{C}}$ having an open dense orbit.
- Fact: compact X_Δ is homeomorphic to $X(\Sigma)$.



Construction of Π_Σ

Poisson Geometry:


- $\pi = \sum_{\ell=1}^d i \partial_{\bar{z}_\ell} \wedge \partial_{z_\ell}$ is translation invariant under action of \mathbb{C}^d .
- $\Pi := \exp_*(\pi) = \sum_{\ell=1}^d i |z_\ell|^2 \partial_{\bar{z}_\ell} \wedge \partial_{z_\ell}$ extends smoothly to all of \mathbb{C}^d .
- Π is invariant under action of $\mathbb{T}_{\mathbb{C}}^d$.
- $[\Pi, \Pi] = [\exp_*(\pi), \exp_*(\pi)] = \exp_*([\pi, \pi]) = 0$.
- $\mathbb{C}^d // N_{\mathbb{C}}$ means $\mathcal{U}_\Sigma / N_{\mathbb{C}}$ where $\mathcal{U}_\Sigma \subset \mathbb{C}^d$ is the free locus of $N_{\mathbb{C}}$ action.
- Π_Σ is the Poisson structure on $X(\Sigma)$ co-induced from Π by the quotient map $\mathcal{U}_\Sigma \rightarrow \mathcal{U}_\Sigma / N_{\mathbb{C}}$.

Symplectic Leaves

Theorem 1 *The symplectic leaves of $(X(\Sigma), \Pi_\Sigma)$ are the orbits of $T_{\mathbb{C}} \simeq \mathbb{T}^d / N_{\mathbb{C}}$. In particular, Π_Σ is non-degenerate on an open dense set.*

Local picture when $\dim_{\mathbb{C}} T_{\mathbb{C}} = 1$:

- $\Pi_\Sigma = 2i|w|^2 \partial_{\bar{w}} \wedge \partial_w = (x^2 + y^2) \partial_x \wedge \partial_y$.
- T action Hamiltonian with primitive $\approx \log |w|$.
- $T_{\mathbb{C}}/T$ action not Hamiltonian due to topology of open leaf, local primitive $\approx \text{Arg}(w)$.



Some generators of $H^1(X(\Sigma), \Pi_\Sigma)$

Theorem 2 *The action of $T_{\mathbb{C}}$ on $(X(\Sigma), \Pi_\Sigma)$ is Poisson but not Hamiltonian, however, each symplectic leaf admits a Hamiltonian action by a sub-torus of T .*

- In hol. coordinates (w_1, \dots, w_n) assoc. to an open cone $V \in \Sigma$,

$$\Pi_\Sigma = \sum_{p,q=1}^n i B_V^{pq} \bar{w}_p w_q \partial_{\bar{w}_p} \wedge \partial_{w_q}$$

where B_V is a symm. pos. def. integral matrix determined by V .

Corollary 1 $\mathfrak{t} + i\mathfrak{t} \hookrightarrow H^1(X(\Sigma), \Pi_\Sigma)$

The Modular Class

- (M, π) orientable, μ a volume form on M .
- θ_μ (modular vector field), char. by $\mathcal{L}_{\pi^\#(df)}\mu = \theta_\mu(f)\mu$.
- π non-deg. and $\mu = \pi^{-n} \Rightarrow \theta_\mu = 0$.
- $\mathcal{L}_{\theta_\mu}\pi = 0 \Rightarrow [\theta_\mu] \in H^1(M, \pi)$ (modular class).
- $\theta_{a\mu} = \theta_\mu - \pi^\#(d \log |a|)$ (class independence of μ .)

$[\theta_\mu]$ as an invariant:

(M, π)	$\pi = 0$	$(\mathfrak{g}^*, \pi_{\mathfrak{g}^*})$	$(K/T, \pi_{K/T})$	π non-deg.
$[\theta_\mu]$	$\theta_\mu = 0$	$[\theta_\mu] = 0$ iff \mathfrak{g} unimodular	$[\theta_\mu] \neq 0$	$\theta_\mu = 0$

Key Point: $[\theta_\mu]$ is an interesting invariant away from the extremes.

Estimate for $H^1(X(\Sigma), \Pi_\Sigma)$

Theorem 3 $\dim_{\mathbb{R}} H^1(X(\Sigma), \Pi_\Sigma) \geq 2 \dim_{\mathbb{R}} T + 1$

Proof:

- In an affine chart \mathbb{C}^n associated an open cone $V \in \Sigma$, $\mu = a\lambda$ where $\lambda = i^n dw_1 \wedge d\bar{w}_1 \wedge \cdots \wedge dw_n \wedge d\bar{w}_n$ and $a \in C^\infty(\mathbb{C}^n)$ is non-vanishing.
- Local representation of Π_Σ shows that θ_λ belongs to the range of $\mathfrak{t} + i\mathfrak{t} \hookrightarrow \mathcal{V}^1(\mathbb{C}^n)$.
- Hence $\theta_\mu|_{\mathbb{C}^n} = \theta_\lambda - \Pi_\Sigma^\#(d \log |a|)$ and thus by continuity $[\theta_\mu]$ belongs to the image of $\mathfrak{t} + i\mathfrak{t}$ in $H^1(X(\Sigma), \Pi_\Sigma)$ if and only if $\log |a|$ extends continuously off of the affine chart.
- But $X(\Sigma)$ is compact, so we must have $\lim_{\|w\| \rightarrow \infty} |a(w)| = 0$. Thus $[\theta_\mu]$ is independent of the image of $\mathfrak{t} + i\mathfrak{t}$ in $H^1(X(\Sigma), \Pi_\Sigma)$. \square

Bound is tight for $\mathbb{C}P^1$

Theorem 4 (Nakanishi) For $\Pi = 2i|z|^2\partial_{\bar{z}} \wedge \partial_z$ on \mathbb{C} ,

$$H^0(\mathbb{C}, \Pi) = \mathbb{R},$$

$$H^1(\mathbb{C}, \Pi) = \mathbb{R}\langle R, D \rangle, \text{ and}$$

$$H^2(\mathbb{C}, \Pi) = \mathbb{R}\langle \pi, \Pi \rangle.$$

- Mayer-Vietoris argument yields:




Theorem 5 For Π_Σ on $\mathbb{C}P^1$,

$$H^0(\mathbb{C}P^1, \Pi_\Sigma) = \mathbb{R},$$

$$H^1(\mathbb{C}P^1, \Pi_\Sigma) = \mathbb{R}\langle R, D, \theta_\mu \rangle, \text{ and}$$

$$H^2(\mathbb{C}P^1, \Pi_\Sigma) \simeq \mathbb{R}^4.$$

Special Representatives for $H^1(\mathbb{C}P^1, \Pi_\Sigma)$

$R = iw\partial_w + c.c.$	$D = w\partial_w + c.c.$	θ_μ
		

θ_μ **for** $(\mathbb{C}P^n, \Pi_\Sigma)$ **for** $\mu = \frac{1}{n!} \omega_\Delta^n$

- The standard n -simplex Δ' is a Delzant polytope in \mathfrak{t}^* .
- Set $\Delta = \Delta' - \nu$ where ν is the center of mass of Δ' and let Σ be the dual fan of Δ .
- Apply Delzant construction to get $(\mathbb{C}P^n, \omega_\Delta, \Phi_\Delta)$ and $\mu = \frac{1}{n!} \omega_\Delta^n$.

Theorem 6 For each open cone V in Σ

$$\theta_\mu = \frac{(n+1)}{2\pi} \sum_{\ell=1}^n \langle \Phi_\Delta, (uB_V)_\ell \rangle R_\ell$$

where R_1, \dots, R_n are the vector fields corresponding to the basis $u = (u_1, \dots, u_n)$ of $\Lambda \subset \mathfrak{t}$ generating the cone V .

Zeroes of θ_μ on $\mathbb{C}P^n$

Theorem 7 *If $Z = \{\theta_\mu = 0\} \subset \mathbb{C}P^n$, then $\Phi_\Delta(Z)$ is the set of centroids of the faces of Δ .*

