Department of Mathematics University of Notre Dame Math 10120 – Finite Math Spring 2012

Name:_____

Instructors: Garbett & Migliore

Practice Exam II Answers

March 5, 2015

This exam is in two parts on 11 pages and contains 14 problems worth a total of 100 points. You have 1 hour and 15 minutes to work on it. You may use a calculator, but no books, notes, or other aid is allowed. Be sure to write your name on this title page and put your initials at the top of every page in case pages become detached.

You must record on this page your answers to the multiple choice problems.

The partial credit problems should be answered on the page where the problem is given. The spaces on the bottom right part of this page are for me to record your grades, **not** for you to write your answers.

Place an \times through your answer to each problem.

1.	(a)	(b)	(c)	(d)	(e)
2.	(a)	(b)	(c)	(d)	(e)
3.	(a)	(b)	(c)	(d)	(e)
4.	(a)	(b)	(c)	(d)	(e)
5.	(a)	(b)	(c)	(d)	(e)
6.	(a)	(b)	(c)	(d)	(e)
7.	(a)	(b)	(c)	(d)	(e)
8.	(a)	(b)	(c)	(d)	(e)
9.	(a)	(b)	(c)	(d)	(e)
10.	(a)	(b)	(c)	(d)	(e)

MC. ______ 11. _____ 12. _____ 13. _____ 14. _____ Tot. _____

Multiple Choice

	Outcome	Probability		Outcome	Probability
	s_1	0.50			
	s_1	0.25		s_1	0.50
(a)	s_1	0.10	(b)	s_2	0.25
(a)		0.15	(5)	s_3	0.10
	s_1	1		s_4	0.10
	Note	all s_1 's			
	Outcome	Probability		Outcome	Probability
	s_1	0.50		s_1	0.50
(a)	s_2	0.25	(d)	s_2	0.25
(c)	s_3	0.10	(u)	s_3	0.25
	s_4	0.15		s_4	0.10
	Outcome	Probability			
	s_1	0.50			
(a)	s_2	0.25			
(e)	s_3	0.35			
	s_4	-0.10			

1. (5 pts.) Which of the following could be a set of probability assignments?

Correct answer: (c). Answer (a) gives different probabilities to the same outcome. In answer (b) the probabilities add to less than 100%. In answer (d) the answers add to more than 100%. In answer (e) there is a negative probability. So (c) is the only one that could work.

2. (5 pts.) Suppose I roll a die twice and record the sum of the numbers that come up. What is the probability that this sum is 3?

$\langle \rangle$	1	(1) 1	1	(1) 1	, 1
(a)	$\overline{9}$	(b) $\frac{1}{18}$	$(c) \overline{36}$	$(d) \overline{6}$	(e) $\frac{12}{12}$

Correct answer: (b). The sample space consists of all 36 ordered pairs (a, b) where a and b are integers between 1 and 6. Of these, only (1, 2) and (2, 1) add up to 3. So the probability is $\frac{2}{36} = \frac{1}{18}$.

3. (5 pts.) An urn contains some balls. Each of the balls has a number on it, and each is colored. Consider the events $E_{i} = - \text{"The ball has a 2 on it"}$

$$E =$$
 "The ball has a 3 on it
 $F =$ "The ball is green"

Someone asks you "What is the probability that the ball is green and has a 3 on it?" Which of the following probabilities are they asking for?

(a) Pr(E|F) (b) Pr(F|E) (c) $Pr(E \cup F)$ (d) Pr(E) + Pr(F) (e) $Pr(E \cap F)$

Correct answer: (e). We want E to be true and F to be true, which is $E \cap F$.

4. (5 pts.) It is determined that the probability that Stewball does NOT win any specific race is $\frac{3}{5}$. Assuming that the race results are independent, what is the probability that Stewball DOES win the next TWO races?

- (a) 16% (b) 36% (c) 40%
- (d) 12% (e) 60%

Correct answer: (a). The probability that Stewball wins a race is $\frac{2}{5}$ and the events are independent, so the probability of winning the next two races is $\frac{2}{5} \cdot \frac{2}{5} = \frac{4}{25} = 16\%$.

Math 10120 Spring 2012, Practice Exam II Answers

5. (5 pts.) Three students (strangers) meet in line at the dining hall and start chatting. What is the probability that **at least** two were born on the same day of the week?

(a)
$$\frac{1}{7}$$
 (b) $\frac{1}{49}$ (c) $\frac{133}{343}$
(d) $\frac{152}{343}$ (e) $\frac{241}{343}$

Correct answer: (c). We use the complement rule. First we compute the probability that they were all born on different days: $\frac{7 \cdot 6 \cdot 5}{7^3} = \frac{30}{49}.$

So the desired probability is

$$1 - \frac{30}{49} = \frac{19}{49} \left(= \frac{133}{343} \right).$$

6. (5 pts.) An urn contains balls with the following distribution of colors and numbers:

9 red balls: 3 labelled "A," 3 labelled "B" and 3 labelled "C" 10 blue balls: 2 labelled "A," 4 labelled "B" and 4 labelled "C" 12 green balls: 4 labelled "A," 5 labelled "B" and 3 labelled "C"

A ball is chosen at random and found to be labeled "B." What is the probability that it is blue?

(a)	$\frac{2}{5}$	(b)	$\frac{10}{31}$	(c)	$\frac{12}{31}$
(d)	$\frac{1}{3}$	(e)	$\frac{5}{6}$		

Correct answer: (d). There are 12 balls labeled "B," of which 4 are blue. So given that the ball is labeled "B," the probability is $\frac{4}{12} = \frac{1}{3}$ that it is blue.

Initials:_____

Math 10120 Spring 2012, Practice Exam II Answers

7. (5 pts.) In a football game, a coin is tossed before the game starts to determine who receives the ball first. Suppose that the local team plays 13 games this season. What is the probability that they will win exactly seven of the coin tosses? (Careful: a lot of these answers look similar. Make sure which one you're picking.)

(a)
$$\frac{\binom{13}{7}}{2^7}$$
 (b) $\frac{\binom{13}{7}}{2^{13}}$ (c) $\frac{P(13,7)}{2^{13}}$
(d) $\frac{7}{13}$ (e) $\frac{\binom{13}{7}}{P(13,7)}$

Correct answer: (b). There are 2^{13} sequences of W's and L's (wins and losses) for the flips. Of these, $\binom{13}{7}$ have exactly 7 wins. (Choose the 7 wins out of 13 tosses.) So the probability is given by (b).

8. (5 pts.) A (not very high quality) company manufactures stereo equipment. Experience shows that defects in manufacturing are independent of one another. Quality control studies reveal that

- 10% of their CD players are defective;
- 20% of their amplifiers are defective;
- 50% of their speakers are defective.

A system consists of a CD players, an amplifier, and **TWO** speakers. What is the probability that the entire system is NOT defective?

- (a) 0.5% (b) 18% (c) 82%
- (d) 97% (e) 99.5%

Correct answer: (b). The probability of the CD player not being defective is 1 - 0.10 = 0.90. The probability of the amp not being defective is 1 - 0.20 = 0.80. The probability of one speaker not being defective is 1 - 0.50 = 0.50. So since there are two speakers, the probability of all four components simultaneously not being defective is

$$(.9)(.8)(.5)(.5) = 0.18.$$

9. (5 pts.) The scores of top 10 finalists for the Uneven Parallel Bars at the USA Gymnastics Collegiate Nationals Event Finals 2014 are as follow:

9.85, 9.80, 9.82, 9.75, 9.70, 9.70, 9.85, 9.85, 9.80, 9.82.

Compute the mean score (to three decimal places).

- (a) 9.800 (b) 9.850 (c) 9.784
- (d) 9.794 (e) 9.810

Correct answer: (c). Add the scores and divide by 10.

10. (5 pts.) Let *E* and *F* be **MUTUALLY EXCLUSIVE** events. Assume that Pr(E) = 0.20 and Pr(F) = 0.30. Find $Pr(E \cup F)$.

(a) 0.20 (b) 0.30 (c) 0.44

(d) 0.50 (e) There is not enough information.

Correct answer: (d). We have the formula $Pr(E \cup F) = Pr(E) + Pr(F) - Pr(E \cap F)$. Since E and F are mutually exclusive, $E \cap F = \emptyset$, so $Pr(E \cap F) = 0$ and we get $Pr(E \cup F) = Pr(E) + Pr(F) = 0.20 + 0.30 = 0.50$.

Partial Credit

You must show all of your work on the partial credit problems to receive credit! Make sure that your answer is clearly indicated. You're more likely to get partial credit for a wrong answer if you explain your reasoning.

11. (10 pts.) The Census Bureau has determined that among all families with four children (no multiple births), all 16 possible orders of boys and girls (e.g. BGBB) are equally likely to occur.

(a) If a family is chosen at random from the Bureau's list of families with four children, what is the probability that exactly two are boys?

Answer: There are four "slots" for the births: first, second, third and fourth. There are $\binom{4}{2} = 6$ ways to choose which two slots are the boys. As noted in the problem, there are $16 (= 2^4)$ possible orders all together. Hence the probability of exactly two boys is $\frac{6}{16} = \frac{3}{8} = 0.375 = 37.5\%$.


(b) Suppose that a family is chosen at random, and it is noticed that they have at least two boys. What is the probability that they have exactly two boys?

Answer: Besides the 6 ways of having exactly two boys, there are 4 ways to have 3 boys and one girl, and there is 1 way to have all boys. So there are 6 + 4 + 1 = 11 ways to have at least two boys, of which 6 consist of exactly two boys. So the probability is $\frac{6}{11} \sim 0.5455 = 54.55\%$.

Math 10120 Spring 2012, Practice Exam II Answers

Initials:____


12. (10 pts.) For this problem, you may leave your answer in terms of mixtures of combinations and permutations (i.e. C(n,r) and P(n,r) for appropriate n and r) if you like.



(a) Refer to the above city map. Sam starts at school (marked S) and needs to go home (marked H). He chooses a random path traveling only east and south. A bully is lurking in the spot marked B. What is the probability that Sam goes by the bully? (Hint: How many paths all together go from S to H? How many paths go through B?)

Answer: There are 12 blocks from S to H, of which 4 are south. So there are a total of $\binom{12}{4}$ routes, as we discussed in chapter 5. That will be the denominator. For the numerator, there are 5 blocks from S to B, of which 2 are south. So there are $\binom{5}{2}$ routes from S to B. For each of these, there are $\binom{7}{2}$ routes from B to H. So the probability is

$$\frac{\binom{5}{2}\cdot\binom{7}{2}}{\binom{12}{4}} = \frac{210}{495} = \frac{14}{33} \sim 42.42\%.$$




(b) Sam starts at school (marked S) and needs to go home (marked H). He randomly chooses a route traveling only east and south. A leprechaun has left two pots of gold at the points marked P_1 and P_2 . What is the probability that he passes by **both** of the pots?

Answer: In the same way, the answer is

$$\frac{\binom{4}{1} \cdot \binom{4}{2} \cdot \binom{4}{1}}{\binom{12}{4}} = \frac{96}{495} = \frac{32}{165} \sim 19.39\%.$$

Math 10120 Spring 2012, Practice Exam II Answers

13. (10 pts.) Consider the following Venn diagram:



Note that there are 180 outcomes all together. If an outcome is selected at random from U, find the following probabilities:

(a)
$$Pr(E) = \frac{n(E)}{n(U)} = \frac{10+5+20+15}{180} = \frac{50}{180} = \frac{5}{18} \sim 27.78\%.$$

(b)
$$Pr(E|F) = \frac{Pr(E \cap F)}{Pr(F)} = \frac{5+15}{5+15+25+30} = \frac{20}{75} = \frac{4}{15} \sim 26.67\%.$$

(c)
$$Pr(E \cap G') = \frac{10+5}{180} = \frac{15}{180} = \frac{1}{12} \sim 8.33\%.$$

(d)
$$Pr(F \cup G) = \frac{5 + 25 + 15 + 30 + 20 + 35}{180} = \frac{130}{180} = \frac{13}{18} \sim 72.22\%$$

14. (10 pts.) Jack plays the following game. First he rolls a die and observes the number that comes up; then he draws a card from a standard deck and observes the suit. (The number on the card is irrelevant; only the suit matters. Remember that there are four suits, all equally likely to be drawn.) Here's how the game works:


If the die shows a 1 or a 2, then he needs a "club" to win.

If the die shows a 3,4,5 or 6, then he needs a "diamond" or a "heart" to win.

(a) Draw a tree diagram for this game, with all the probabilities labelled and indicating which endpoints correspond to winning the game.

Let

- E = "die shows 1 or 2"
- F = "die shows 3, 4, 5 or 6"
- C = "card is a club"
- D = "card is a diamond"
- H = "card is a diamon" H = "card is a heart"
- S = "card is a spade"



(b) What is the probability of winning the game?

The probability is

$$\left(\frac{1}{3}\right)\left(\frac{1}{4}\right) + \left(\frac{2}{3}\right)\left(\frac{1}{4}\right) + \left(\frac{2}{3}\right)\left(\frac{1}{4}\right) = \frac{5}{12} \sim 0.4167 = 41.67\%.$$

(c) If Jack won the game, what is the probability that he rolled a 1 or a 2 in the first part?

We'll use the answer to part (b) as part of this answer (the denominator), instead of re-calculating it.

$$Pr(E|\text{win}) = \frac{Pr(E \cap \text{win})}{Pr(\text{win})} = \frac{\left(\frac{1}{3}\right)\left(\frac{1}{4}\right)}{\frac{5}{12}} = \frac{1}{5} = 20\%.$$

Math 10120 Spring 2012, Practice Exam II Answers

15. (10 pts.)


Zoologists studying wolfpacks close to the arctic circle recorded the sizes of the packs that they found, as follows:

6, 10, 7, 5, 7, 7, 2, 4, 3, 2, 2, 3, 9, 4, 4, 2, 8, 7, 8, 4.

(a) Organize the data in a relative frequency table.

Outcome	Frequency	
2	4	4/20 = 1/5 = 0.2
3	2	2/20 = 1/10 = 0.1
4	4	4/20 = 4/5 = 0.2
5	1	1/20 = 0.05
6	1	1/20 = 0.05
7	4	4/20 = 1/5 = 0.2
8	2	2/20 = 1/10 = 0.1
9	1	1/20 = 0.05
10	1	1/20 = 0.05
Totals	20	1.00

(b) Draw a histogram for the data on the axes provided below:



Initials:_____

Department of Mathematics University of Notre Dame Math 10120 – Finite Math Spring 2012

Name:_____

Instructors: Garbett & Migliore

Practice Exam II Answers

March 5, 2015

This exam is in two parts on 11 pages and contains 14 problems worth a total of 100 points. You have 1 hour and 15 minutes to work on it. You may use a calculator, but no books, notes, or other aid is allowed. Be sure to write your name on this title page and put your initials at the top of every page in case pages become detached.

You must record on this page your answers to the multiple choice problems.

The partial credit problems should be answered on the page where the problem is given. The spaces on the bottom right part of this page are for me to record your grades, **not** for you to write your answers.

Place an \times through your answer to each problem.

1.	(a)	(b)	(\bullet)	(d)	(e)
2.	(a)	(\mathbf{b})	(c)	(d)	(e)
3.	(a)	(b)	(c)	(d)	(ullet)
4.	(\mathbf{a})	(b)	(c)	(d)	(e)
5.	(a)	(b)	(ullet)	(d)	(e)
6.	(a)	(b)	(c)	(\mathbf{q})	(e)
7.	(a)	(\mathbf{b})	(c)	(d)	(e)
8.	(a)	(\mathbf{b})	(c)	(d)	(e)
9.	(a)	(b)	(c)	(\mathbf{q})	(e)
10.	(a)	(b)	(c)	(\mathbf{q})	(e)

MC. ______ 11. _____ 12. _____ 13. _____ 14. _____ Tot. _____