

DIRAC OPERATORS ON COBORDISMS: DEGENERATIONS AND SURGERY

DANIEL F. CIBOTARU AND LIVIU I. NICOLAESCU

ABSTRACT. We investigate the Dolbeault operator on a pair of pants, i.e., an elementary cobordism between a circle and the disjoint union of two circles. This operator induces a canonical selfadjoint Dirac operator D_t on each regular level set C_t of a fixed Morse function defining this cobordism. We show that as we approach the critical level set C_0 from above and from below these operators converge in the gap topology to (different) selfadjoint operators D_{\pm} that we describe explicitly. We also relate the Atiyah-Patodi-Singer index of the Dolbeault operator on the cobordism to the spectral flows of the operators D_t on the complement of C_0 and the Kashiwara-Wall index of a triplet of finite dimensional lagrangian spaces canonically determined by C_0 .

CONTENTS

Introduction	1
1. A model degeneration	4
2. The Atiyah-Patodi-Singer theorem	12
3. Dolbeault operators on two-dimensional cobordisms	18
4. The Kashiwara-Wall index	26
References	31

Introduction

Suppose (M, g) is compact oriented odd dimensional Riemann manifold. We let \widehat{M} denote the cylinder $[0, 1] \times M$ and \widehat{g} denote the cylindrical metric $dt^2 + g$.

Let \widehat{D} be a first order elliptic operator on \widehat{M} that has the form

$$\widehat{D} = \sigma(dt)(\nabla_t - D(t)), \quad (\dagger)$$

where σ denotes the principal symbol of \widehat{D} , and for every $t \in [0, 1]$ the operator $D(t)$ on $\{t\} \times M$ is elliptic and symmetric. For simplicity we assume that both $A(0)$ and $A(1)$ are invertible.

A classical result of Atiyah, Patodi and Singer [2, §7] (see also [12, §17.1]) relates the index $i_{APS}(\widehat{A})$ of the Atiyah-Patodi-Singer problem associated to \widehat{D} to the spectral flow $SF(D(t))$ of the family of Fredholm selfadjoint operators $D(t)$. More precisely, they show that

$$i_{APS}(\widehat{D}) + SF(D(t), 0 \leq t \leq 1) = 0. \quad (\text{A})$$

We can regard the cylinder \widehat{M} as a trivial cobordism between $\{0\} \times M$ and $\{1\} \times M$, and the coordinate t as a Morse function on \widehat{M} with no critical points.

Date: Started July 14, 2009. Completed on August 14, 2009. This is the *September 14, 2009* version.

2000 Mathematics Subject Classification. Primary 58J20, 58J28, 58J30, 58J32, 53B20, 35B25.

Key words and phrases. Atiyah-Patodi-Singer index theorem, spectral flows, elliptic boundary value problems, lagrangian spaces, Kashiwara index.

In this paper we initiate an investigation of the case when \widehat{M} is no longer a trivial cobordism. We outline below the main themes of this investigation.

First, we will concentrate only on elementary cobordisms, the ones that trace a single surgery. We regard such a cobordism as a pair (\widehat{M}, f) , where \widehat{M} is an even dimensional, compact oriented manifold with boundary, and f is a Morse function on \widehat{M} with a single critical point p_0 such that

$$f(\widehat{M}) = [-1, 1], \quad f(\partial M) = \{-1, 1\}, \quad f(p_0) = 0.$$

We set $M_{\pm} := f^{-1}(\pm 1)$ so that we have a diffeomorphism of oriented manifolds $\partial M = M_+ \cup -M_-$. By removing the critical level set $M_0 = f^{-1}(0)$ we obtain two cylinders

$$\widehat{M}_- = \{f < 0\} \cong [-1, 0) \times M_-, \quad \widehat{M}_+ = \{f > 0\} \cong (0, 1] \times M_+.$$

Suppose \hat{g} is a Riemann metric on \widehat{M} and $\widehat{D} : C^\infty(E_+) \rightarrow C^\infty(E_-)$ is a Dirac type operator on \widehat{M} , where $E_+ \oplus E_-$ is a $\mathbb{Z}/2$ -graded bundle of Clifford modules.

Using the bundle isomorphism $\sigma(dt)$ we can regard \widehat{D} as an operator $C^\infty(E_+) \rightarrow C^\infty(E_+)$. As explained in [8] (see also Section 2 of this paper), for every $t \neq 0$, there is a canonically induced symmetric Dirac operator $D(t)$ on the slice $M_t = f^{-1}(t)$. We regard $D(t)$ as a linear operator $D(t) : C^\infty(E_+|_{M_t}) \rightarrow C^\infty(E_+|_{M_t})$, so that if \hat{g} were a cylindrical metric then formula (†) would hold.

The Riemann metric \hat{g} defines finite measures dV_t on all the slices M_t , including the singular slice M_0 . In particular we obtain a one parameter family of Hilbert spaces

$$\mathbf{H}_t := L^2(M_t, dV_t; E_+).$$

We can now regard $D(t)$ as a closed, densely defined linear operator on \mathbf{H}_t .

Problem 1. Organize the family $(\mathbf{H}_t)_{t \in [-1, 1]}$ as a trivial Hilbert bundle over the interval $[-1, 1]$

$$\mathcal{H} = \mathbf{H} \times [-1, 1] \rightarrow [-1, 1].$$

Under reasonable assumptions on f and \hat{g} we can use the gradient flow of f to address this issue. Once this problem is solved we can regard the operators $D(t)$, $t \neq 0$ as closed densely defined operators on the same Hilbert space \mathbf{H} . We can then formulate our next problem.

Problem 2. Investigate whether the limits

$$SF_- := \lim_{\varepsilon \searrow 0} SF(D(t), -1 \leq t \leq -\varepsilon), \quad SF_+ := \lim_{\varepsilon \searrow 0} SF(D(t), \varepsilon \leq t \leq 1).$$

exist and are finite.

If Problem 2 has a positive answer we are interested in a version of (A) relating these limits to the Atiyah-Patodi-Singer index of \widehat{D} in the noncylindrical formulation of [8, 9].

Problem 3. Express the quantity

$$\delta := i_{APS}(\widehat{D}) + SF_- + SF_+ \tag{B}$$

in terms of invariants of the singular level set M_0 .

The existence of the limits in Problem 2 is a consequence of a much more refined analytic behavior of the family of operators $D(t)$ that we now proceed to explain. We set

$$\widehat{\mathbf{H}} := \mathbf{H} \oplus \mathbf{H}, \quad \mathbf{H}_+ = \mathbf{H} \oplus 0, \quad \mathbf{H}_- = 0 \oplus \mathbf{H},$$

and denote by Lag the Grassmannian of hermitian lagrangian subspaces $\widehat{\mathbf{H}}$. These are complex subspaces $L \subset \widehat{\mathbf{H}}$ satisfying $L^\perp = JL$, where $J : \mathbf{H} \oplus \mathbf{H} \rightarrow \mathbf{H} \oplus \mathbf{H}$ is the operator with block decomposition

$$J = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Following [5] we denote by Lag^- the open subset of Lag consisting of lagrangians L such that the pair of subspaces $(L, \widehat{\mathbf{H}}_-)$ is a Fredholm pair, i.e.,

$$L + \mathbf{H}_- \text{ is closed and } \dim L \cap \mathbf{H}_- < \infty$$

As explained in [5], the space Lag^- equipped with the gap topology of [10, §IV.2] is a classifying spaces for the complex K -theoretic functor K^1 .

To a closed densely defined operator $T : \text{Dom}(T) \subset \mathbf{H} \rightarrow \mathbf{H}$ we associate its switched graph

$$\widetilde{\Gamma}_T := \left\{ (Th, h) \in \widehat{\mathbf{H}}; h \in \text{Dom}(T) \right\}.$$

Then T is selfadjoint if and only if $\widetilde{\Gamma}_T \in \text{Lag}$. It is also Fredholm if and only if $\widetilde{\Gamma}_T \in \text{Lag}^-$. We can now formulate a refinement of Problem 2.

Problem 2*. Investigate whether the limits $\widetilde{\Gamma}_\pm = \lim_{t \searrow 0} \widetilde{\Gamma}_{D(\pm t)}$ exist in the gap topology and, if so, do they belong to Lag^- .

The gap convergence of the switched graphs of operators is equivalent to the convergence in norm as $t \rightarrow 0^\pm$ of the resolvents $R_t = (i + D(t))^{-1}$. To show that $\widetilde{\Gamma}_\pm \in \text{Lag}^-$ it suffices to show that the limits $R_\pm = \lim_{t \rightarrow 0^\pm} R_t$ are compact operators. If in addition¹ $\widetilde{\Gamma}_\pm \cap \widehat{\mathbf{H}}_- = 0$ then the limits in Problem 2 exist and are finite.

An even analog of Problem 2* was investigated in [16]. The role of the smooth slices M_t was played there by a 1-parameter family of Riemann surfaces degenerating to a Riemann surface with single singularity of the simplest type, a node. The authors show that the gap limit of the graphs of Dolbeault operators on M_t exists and then described it explicitly.

In this paper we solve Problems 1, 2* and 3 in the symplectic case, when \widehat{M} is an elementary 2-dimensional cobordism, i.e., a pair of pants (see Figure 1) and \widehat{D} is the Dolbeault operator on the Riemann surface \widehat{M} .

We solved Problem 1 by an ad-hoc intuitive method. The limits $\widetilde{\Gamma}_\pm$ in Problem 2* turned out to be switched graphs of certain Fredholm-selfadjoint operators D_\pm , $\widetilde{\Gamma}_\pm = \widetilde{\Gamma}_{D_\pm}$.

We describe these operators as realizations of two different boundary value problems associated to the same symmetric Dirac operator D_0 defined on the disjoint union of four intervals. These intervals are obtained by removing the singular point of the critical level set M_0 and then cutting in two each of the resulting two components. The boundary conditions defining D_\pm are described by some (4-dimensional) lagrangians Λ_\pm determined by the geometry of the singular slice M_0 . The operators D_\pm have well defined eta invariants η_\pm . If $\ker D_\pm = 0$ then we can express the defect δ in (B) as

$$\delta = \frac{1}{2}(\eta_- - \eta_+). \quad (\text{C})$$

The above difference of eta invariants admits a purely symplectic interpretation very similar to the signature additivity defect of Wall [19]. More precisely, we show that

$$\delta = -\omega(\Lambda_0^\perp, \Lambda_+, \Lambda_-), \quad (\text{D})$$

¹The condition $\widetilde{\Gamma}_\pm \cap \widehat{\mathbf{H}}_- = 0$ is not really needed, but it makes our presentation more transparent. In any case, it is generically satisfied.

where Λ_0 is the Cauchy data space of the operator D_0 and $\omega(L_0, L_1, L_2)$ denotes the Kashiwara-Wall index of a triplet of lagrangians canonically determined by M_0 ; see [4, 11, 19] or Section 4.

Here is briefly how we structured the paper. In Section 1 we investigate in great detail the type of degenerations that occur in the family $D(t)$ as $t \rightarrow 0^\pm$. It boils down to understanding the behavior of families of operators of the unit circle S^1 of the type

$$L_\varepsilon = -i \frac{d}{d\theta} + a_\varepsilon(\theta),$$

where $\{a_\varepsilon\}_{\varepsilon>0}$ is a family of smooth functions on the unit circle that converges in a rather weak sense way as $\varepsilon \rightarrow 0$ to a Dirac measure supported at a point θ_0 . For example if we think of a_ε as densities defining measures converging weakly to the Dirac measure, then the corresponding family of operators has a well defined gap limit; see Corollary 1.5.

In Theorem 1.8 we give an explicit description of this limiting operator as an operator realizing a natural boundary value problem on the *disjoint* union of the two intervals, $[0, \theta_0]$ and $[\theta_0, 2\pi]$. This section also contains a detailed discussion of the eta invariants of operators of the type $-i \frac{d}{d\theta} + a(\theta)$, where a is allowed to be the “density” of any finite Radon measure.

In Section 2 we survey mostly known facts concerning the Atiyah-Patodi-Singer problem when the metric near the boundary is not cylindrical. Because the various orientation conventions vary wildly in the existing literature, we decided to go careful through the computational details. We discuss two topics. First, we explain what is the restriction of a Dirac operator to a cooriented hypersurface and relate this construction to another conceivable notion of restriction. In the second part of this section we discuss the noncylindrical version of the Atiyah-Patodi-Singer index theorem. Here we follow closely the presentation in [8, 9].

In Section 3 we formulate and prove the main result of this paper, Theorem 3.2. The solution to Problem 2* is obtained by reducing the study of the degenerations to the model degenerations investigated in Section 1. The equality (C) follows immediately from the noncylindrical version of the Atiyah-Patodi-Singer index theorem discussed in Section 2 and the eta invariant computations in Section 1. In the last section we present a few facts about the Kashiwara-Wall index and then use them to prove (D).

Finally a few words about conventions and notation. We consistently orient the boundaries using the outer-normal-first convention. We let i stand for $\sqrt{-1}$ and we let $L^{k,p}$ denote Sobolev spaces of functions that have weak derivatives up to order k that belong to L^p .

1. A model degeneration

Let $L > 0$ be a positive number. Denote by \mathbf{H} the Hilbert space $L^2([0, L], \mathbb{C})$. To any smooth function $a : \mathbb{R} \rightarrow \mathbb{R}$ which is L -periodic we associate the selfadjoint operator

$$D_a : \text{Dom}(D_a) \subset \mathbf{H} \rightarrow \mathbf{H},$$

where

$$\text{Dom}(D_a) = \{ u \in L^{1,2}([0, L], \mathbb{C}); u(0) = u(L) \}, \quad D_a u = -i \frac{du}{dt} + au. \quad (1.1)$$

In this section we would like to understand the dependence of D_a on the potential a , and in particular, we would like to allow for more singular potentials such as a Dirac distribution concentrated at an interior point of the interval. We will reach this goal via a limiting procedure that we implement in several steps.

We observe first that D_a can be expressed in terms of the resolvent $R_a := (i + D_a)^{-1}$ as $D_a = R_a^{-1} - i$. The advantage of this point of view is that we can express R_a in terms of the more regular

function

$$A(t) := \int_0^t a(s) ds. \quad (*)$$

which continues to make sense even when there is no integrable function a such that $(*)$ holds. For example, we can allow $A(t)$ to be any function with bounded variation so that, formally, a ought to be the density of any Radon measure on $[0, L]$.

This will allow us to conclude that when we have a family of smooth potentials a_n that converge in a suitable sense to something singular such as a Dirac function, then the operators D_{a_n} have a limit in the gap topology to a Fredholm selfadjoint operator with compact resolvent. We show that in many cases this limit operator can be expressed as the Fredholm operator defined by a boundary value problem.

We begin by expressing R_a as an integral operator. We set

$$A(t) := \int_0^t a(s) ds, \quad \Phi_A(t) := iA(t) - t.$$

For $f \in H$ the function $u = R_a f$ is the solution of the boundary value problem

$$\left(i - i \frac{d}{dt} \right) u + au = f, \quad u(0) = u(L).$$

We rewrite the above equation as

$$\frac{du}{dt} + (ia - 1)u = if$$

from which we deduce

$$\frac{d}{dt} \left(e^{\Phi_A(t)} u(t) \right) = i e^{\Phi_A(t)} f(t).$$

This implies that

$$e^{\Phi_A(t)} u(t) - u(0) = i \int_0^t e^{\Phi_A(s)} f(s) ds, \quad \forall t \in [0, L].$$

If in the above equality we let $t = L$ and use condition $u(0) = u(L)$ we deduce

$$u(0) = \frac{i}{e^{\Phi_A(L)} - 1} \int_0^L e^{\Phi_A(s)} f(s) ds.$$

Finally we deduce

$$u(t) = R_a f = \frac{i e^{-\Phi_A(t)}}{e^{\Phi_A(L)} - 1} \int_0^L e^{\Phi_A(s)} f(s) ds + i \int_0^t e^{-(\Phi_A(t) - \Phi_A(s))} f(s) ds. \quad (1.2)$$

The key point of the above formula is that R_a can be expressed in terms of the antiderivative $A(t)$ which typically has milder singularities than a . To analyze the dependence of R_a on A we introduce a class of admissible functions.

Definition 1.1. (a) We say that $A : [0, L] \rightarrow \mathbb{R}$ is *admissible* if A has bounded variation, it is right continuous, and $A(0) = 0$. We denote by \mathcal{A} or \mathcal{A}_L the class of admissible functions.

(b) We say that a sequence $\{A_n\}_{n \geq 0} \subset \mathcal{A}$ converges *very weakly* to $A \in \mathcal{A}$ if there exists a negligible subset $\Delta \subset (0, L)$ such that

$$\lim_{n \rightarrow \infty} A_n(t) = A(t), \quad \forall t \in [0, L] \setminus \Delta. \quad \square$$

Remark 1.2. (a) Note that if A_n converges very weakly to A then $A_n(L)$ converges to $A(L)$.
 (b) Let us explain the motivation behind the “very weak” terminology. An admissible function A defines a finite Lebesgue-Stieltjes measure μ_A on $[0, L]$, and the resulting map $A \mapsto \mu_A$ is a linear isomorphism between \mathcal{A} and the space of finite Borel measures on $[0, L]$, [7, Thm. 3.29]. Thus, we can identify \mathcal{A} with the space of finite Borel measures on $[0, L]$. As such it is equipped with a weak topology.

According to [6, §4.22], a sequence of Borel measures μ_{A_n} is weakly convergent to μ_A if and only if $\mu_{A_n}(\mathcal{O}) \rightarrow \mu_A(\mathcal{O})$, for any (relatively) open subset \mathcal{O} of $[0, L]$. This clearly implies the very weak convergence introduced in Definition 1.1. \square

Inspired by (1.2) we define for every $A \in \mathcal{A}$ the function $\Phi_A(t) = iA(t) - t$ and the integral kernels

$$\mathfrak{S}_A : [0, L] \times [0, L] \rightarrow \mathbb{C}, \quad \mathfrak{S}_A(t, s) = \frac{i}{e^{\Phi_A(L)} - 1} e^{-(\Phi_A(t) - \Phi_A(s))}, \quad \forall t, s \in [0, L],$$

$$\mathfrak{K}_A : [0, L] \times [0, L] \rightarrow \mathbb{C}, \quad \mathfrak{K}_A(t, s) = \begin{cases} 0 & t < s \\ ie^{-(\Phi_A(t) - \Phi_A(s))} & t \geq s. \end{cases}$$

Observe that there exists a constant $C > 0$ such that

$$\|\mathfrak{S}_A\|_{L^\infty([0, L] \times [0, L])} + \|\mathfrak{K}_A\|_{L^\infty([0, L] \times [0, L])} \leq C, \quad \forall A \in \mathcal{A}. \quad (1.3)$$

Thus, these kernels define bounded compact operators $S_A, K_A : \mathbf{H} \rightarrow \mathbf{H}$; see [18, §X.2]. Moreover, if we denote by $\|\bullet\|_{\text{op}}$ the operator norm on the space $\mathcal{B}(\mathbf{H})$ of bounded linear operators $\mathbf{H} \rightarrow \mathbf{H}$ then we have the estimates that

$$\|S_A\|_{\text{op}} \leq \|\mathfrak{S}_A\|_{L^2([0, L] \times [0, L])}, \quad \|K_A\|_{\text{op}} \leq \|\mathfrak{K}_A\|_{L^2([0, L] \times [0, L])}. \quad (1.4)$$

We can now rewrite (1.2) as

$$R_a = R_A := S_A + K_A. \quad (1.5)$$

Proposition 1.3. *If A_n converges very weakly to A then S_{A_n} and K_{A_n} converge in the operator norm topology to S_A and respectively K_A .*

Proof. The very weak convergence implies that

$$S_{A_n}(t, s) \xrightarrow{k \rightarrow \infty} \mathfrak{S}_A(t, s), \quad \mathfrak{K}_{A_n}(t, s) \xrightarrow{k \rightarrow \infty} \mathfrak{K}_A(t, s) \quad \text{a.e. on } [0, L] \times [0, L].$$

Using (1.3), the above pointwise convergence and the dominated convergence theorem we deduce

$$\lim_{n \rightarrow \infty} \left(\|S_{A_n} - S_A\|_{L^2([0, L] \times [0, L])} + \|\mathfrak{K}_{A_n} - \mathfrak{K}_A\|_{L^2([0, L] \times [0, L])} \right) = 0.$$

Using (1.4) we deduce that

$$\lim_{n \rightarrow \infty} \left(\|S_{A_n} - S_A\|_{\text{op}} + \|\mathfrak{K}_{A_n} - \mathfrak{K}_A\|_{\text{op}} \right) = 0. \quad \square$$

We want to describe the spectral decompositions of the operators $R_A, A \in \mathcal{A}$. To do this we rely on the fact that for certain A 's the operator R_A is the resolvent of an elliptic selfadjoint operator on S^1 . We use this to produce an intelligent guess for the spectrum of R_A in general.

Let a be a smooth, real valued, L -period function on \mathbb{R} and form again the operator D_a defined in (1.1). We set as usual

$$A(t) = \int_0^t a(s) ds.$$

The operator D_a has discrete real spectrum. If $u(t)$ is an eigenfunction corresponding to an eigenvalue λ then

$$-i \frac{du}{dt} + au = \lambda u \Rightarrow \frac{du}{dt} + i(a - \lambda)u = 0$$

so that $u(t) = u(0)e^{-iA(t)+i\lambda t}$. The periodicity assumption implies $\lambda L - A(L) \in 2\pi\mathbb{Z}$ so the spectrum of D_a is

$$\text{spec}(D_a) = \left\{ \lambda_{A,n} := \frac{2\pi}{L}(\omega_A + n); n \in \mathbb{Z} \right\}, \text{ where } \omega_A := \frac{A(L)}{2\pi}. \quad (1.6)$$

The eigenvalue $\lambda_{A,n}$ is simple and the eigenspace corresponding to $\lambda_{A,n}$ is spanned by

$$\psi_{A,n}(t) := e^{\frac{2\pi nit}{L}} e^{-i(A(t) - \frac{A(L)t}{L})}.$$

The numbers $\lambda_{A,n}$ and the functions $\psi_{A,n}$ are well defined for any $A \in \mathcal{A}$.

Lemma 1.4. *Let $A \in \mathcal{A}$. Then the collection $\{\psi_{A,n}(t); n \in \mathbb{Z}\}$ defines a Hilbert basis of \mathbf{H} .*

Proof. Observe first that the collection

$$e_n(t) = \psi_{A=0,n}(t) = e^{\frac{2\pi nit}{L}}, n \in \mathbb{Z}$$

is the canonical Hilbert basis of \mathbf{H} that leads to the classical Fourier decomposition. The map

$$U_A : \mathbf{H} \rightarrow \mathbf{H}, \mathbf{H} \ni f(t) \mapsto e^{-i(A(t) - \frac{A(L)t}{L})} f(t)$$

is unitary. It maps e_n to $\psi_{A,n}$ which proves our claim. \square

A direct computation shows that

$$R_A \psi_{A,n} = \frac{1}{i + \lambda_{A,n}} \psi_{A,n}, \quad \forall A \in \mathcal{A}, A \in \mathcal{A}.$$

This proves that for any $A \in \mathcal{A}$ the collection $\{\psi_{A,n}\}_{n \in \mathbb{Z}}$ is a Hilbert basis that diagonalizes the operator R_A . Observe that R_A is injective and compact. We define

$$T_A := R_A^{-1} - i.$$

The operator T_A is unbounded, closed and densely defined with domain $\text{Dom}(T_A) = \text{Range}(R_A)$. We will present later a more explicit description of $\text{Dom}(T_A)$ for a large class of A 's.

Note that when

$$A = \int_0^t a(s) ds, \quad a \text{ smooth and } L\text{-periodic},$$

the operator T_A coincides with the operator D_a defined in (1.1). Proposition 1.3 can be rephrased as follows.

Corollary 1.5. *If the sequence $(A_n)_{n \geq 1} \subset \mathcal{A}$ converges very weakly to $A \in \mathcal{A}$ then the sequence of unbounded operators $(T_{A_n})_{n \geq 1}$ converges in the gap topology to the unbounded operator T_A . \square*

The spectrum of T_A consists only of the simple eigenvalues $\lambda_{A,n}$, $n \in \mathbb{Z}$. The function ψ_{A_n} is an eigenfunction of T_A corresponding to the eigenvalue $\lambda_{A,n}$. The eta invariant of T_A is now easy to compute. For $s \in \mathbb{C}$ we have

$$\eta_A(s) := \sum_{\lambda > 0} \frac{1}{\lambda^s} \left(\dim \ker(\lambda - T_A) - \dim \ker(\lambda + T_A) \right)$$

$$= \sum_{n \in \mathbb{Z} \setminus \{-\omega_A\}} \frac{\text{sign } \lambda_{A,n}}{|\lambda_{A,n}|^s} = \frac{L^s}{2\pi^s} \sum_{n \in \mathbb{Z} \setminus \{-\omega_A\}} \frac{\text{sign}(n + \omega_A)}{|n + \omega_A|^s}.$$

Let

$$\rho_A := \omega_A - \lfloor \omega_A \rfloor = \frac{A(L)}{2\pi} - \left\lfloor \frac{A(L)}{2\pi} \right\rfloor \in [0, 1). \quad (1.7)$$

If $\rho_A = 0$ then $\eta_A(s) = 0$ because in this case the spectrum of T_A is symmetric about the origin. If $\rho_A \neq 0$ then we have

$$\eta_A(s) = \frac{L^s}{2\pi^s} \left(\sum_{n \geq 0} \frac{1}{(n + \rho_A)^s} - \sum_{n \geq 0} \frac{1}{(n + 1 - \rho_A)^s} \right) = \frac{L^s}{2\pi^s} \left(\zeta(s, \rho_A) - \zeta(s, 1 - \rho_A) \right),$$

where for every $a \in (0, 1]$ we denoted by $\zeta(s, a)$ the Riemann-Hurwitz zeta function

$$\zeta(s, a) = \sum_{n \geq 0} \frac{1}{(n + a)^s}.$$

The above series is convergent for any $s \in \mathbb{C}$, $\text{Res} > 1$ and admits an analytic continuation to the punctured plane $\mathbb{C} \setminus \{s = 1\}$. Its value at the origin $s = 0$ is given by Hermite's formula [17, 13.21]

$$\zeta(0, a) = \frac{1}{2} - a. \quad (1.8)$$

We deduce that $\eta_A(s)$ has an analytic continuation at $s = 0$ and we have

$$\eta_A(0) = \begin{cases} 0 & \text{if } \rho_A = 0, \\ 1 - 2\rho_A & \text{if } \rho_A \in (0, 1). \end{cases} \quad (1.9)$$

If we introduce the function

$$\xi_A = \frac{1}{2} (\dim \ker T_A + \eta_A(0)),$$

then we can rewrite the above equality in a more compact way

$$\xi_A = \frac{1}{2} (1 - 2\rho_A) = \frac{1}{2} - \rho_A. \quad (1.10)$$

Suppose we have $A_0, A_1 \in \mathcal{A}$. We set $A_s = A_0 + s(A_1 - A_0) \in \mathcal{A}$. The map $[0, 1] \ni s \mapsto A_s \in \mathcal{A}$ is continuous in the weak topology on \mathcal{A} and thus the family of operators T_{A_s} is continuous with respect to the gap topology. The eigenvalues of the family T_{A_s} can be organized in smooth families

$$\lambda_{s,n} = \frac{2\pi}{L} (\omega_s + n) = \frac{2\pi}{L} (\omega_{A_0} + s(\omega_1 - \omega_0) + n), \quad \omega_s := \omega_{A_s}, \quad \forall s \in [0, 1].$$

Assume for simplicity that $\omega_0, \omega_1 \notin \mathbb{Z}$, i.e., the operators T_{A_0} and T_{A_1} are invertible. Denote by $SF(A_1, A_0)$ the spectral flow of the affine family² T_{A_s} . Then

$$\begin{aligned} SF(A_1, A_0) &= \#\{n \in \mathbb{Z}; \omega_0 + n < 0, \omega_1 + n > 0\} - \#\{n \in \mathbb{Z}; \omega_0 + n > 0, \omega_1 + n < 0\} \\ &= \#(\mathbb{Z} \cap (\omega_0, \omega_1)) - \#(\mathbb{Z} \cap (\omega_1, \omega_0)). \end{aligned}$$

We conclude

$$SF(A_1, A_0) = (\lfloor \omega_1 \rfloor - \lfloor \omega_0 \rfloor), \quad \omega_i = \frac{A_i(L)}{2\pi}. \quad (1.11)$$

Using (1.10) we deduce

$$SF(A_1, A_0) = \lfloor \omega_{A_1} \rfloor - \lfloor \omega_{A_0} \rfloor = \omega_{A_1} - \omega_{A_0} + (\xi_{A_1} - \xi_{A_0}). \quad (1.12)$$

²The quantity $SF(A_1, A_0)$ is independent of the weakly continuous path A_s connecting A_0 to A_1 since the space \mathcal{A} equipped with the weak topology is contractible. It is thus an invariant of the pair (A_1, A_0) .

Remark 1.6 (Rescaling trick). Note that the rescaling

$$[0, L_1] \ni \tau \mapsto t = \frac{\tau}{c} \in [0, L_0], \quad c = \frac{L_1}{L_0}.$$

induces an isometry $\mathcal{J}_{L_1, L_0} : \mathbf{H}_{L_0} = L^2(0, L_0; \mathbb{C}) \rightarrow \mathbf{H}_{L_1} = L^2(0, L_1; \mathbb{C})$,

$$\mathbf{H}_{L_0} \ni f(t) \mapsto \mathcal{J}_{L_1, L_0} f(\tau) := c^{1/2} f\left(\frac{\tau}{c}\right) \in \mathbf{H}_{L_1}.$$

The unbounded operator $\frac{d}{dt}$ on \mathbf{H}_{L_0} is the conjugate to the operator $c\frac{d}{d\tau}$ on \mathbf{H}_{L_1} .

If $\alpha(t)$ is a real bounded measurable function on $[0, L_0]$, then the bounded operator on \mathbf{H}_{L_0} defined by pointwise multiplication by $\alpha(t)$ is conjugate to the bounded operator on \mathbf{H}_{L_1} defined by the multiplication by $a(\tau) = \alpha(\tau/c)$. Hence the unbounded operator D_b on \mathbf{H}_{L_0} is conjugate to the unbounded operator $cD_{c^{-1}a}$ on \mathbf{H}_{L_1} ,

$$cD_{c^{-1}a} = \mathcal{J}_{L_1, L_0} D_\alpha \mathcal{J}_{L_1, L_0}^{-1}. \quad (1.13)$$

Its resolvent is obtained by solving the periodic boundary value problem

$$\mathbf{i}u + c \left(-\mathbf{i} \frac{d}{d\tau} + c^{-1}a(\tau) \right) u(\tau) = f(\tau), \quad u(0) = u(L_1),$$

or equivalently

$$\frac{d}{d\tau} u + c^{-1}(a(\tau) - \mathbf{i})u = c^{-1}\mathbf{i}f, \quad u(0) = u(L_1).$$

If we set

$$A(\tau) = \int_0^\tau a(\sigma) d\sigma \quad \text{and} \quad \Phi_{A,c}(t) = c^{-1}\Phi_A(\tau) = c^{-1}(\mathbf{i}A(\tau) - \tau),$$

then we see that R_α is conjugate to the integral operator $R_{A,c}$

$$R_{A,c}f(\tau) = \frac{c^{-1}\mathbf{i}e^{-\Phi_{A,c}(\tau)}}{e^{\Phi_{A,c}(L_1)} - 1} \int_0^{L_1} e^{\Phi_{A,c}(\sigma)} f(s) ds + c^{-1}\mathbf{i} \int_0^\tau e^{-(\Phi_{A,c}(\tau) - \Phi_A(\sigma))} f(\sigma) d\sigma.$$

Arguing exactly as in the proof of Proposition 1.3 we deduce that if A_n covers very weakly to $A \in \mathcal{A}_{L_1}$ and the sequence of positive numbers c_n converges to the positive number c then R_{A_n, c_n} converges in the operator norm to $R_{A,c}$.

For any $c > 0$ and $A \in \mathcal{A}$ we define the operator

$$T_{A,c} = R_{A,c}^{-1} - \mathbf{i}, \quad c > 0.$$

Note that $T_{A,c} = cT_{c^{-1}A}$. Then for every $c > 0$ the spectrum of $T_{A,c}$ is

$$\text{spec}(T_{A,c}) = c \text{spec}(T_{c^{-1}A}). \quad \square$$

We want to give a more intuitive description of the operators R_A , and T_A for a large class of A 's. We begin by introducing a nice subclass \mathcal{A}_* of \mathcal{A} . Let $H(t)$ denote the Heaviside function

$$H(t) = \begin{cases} 1, & t \geq 0 \\ 0, & t < 0. \end{cases}$$

Definition 1.7. We say that $A \in \mathcal{A}$ is *nice* if there exists $a \in L^\infty(0, L)$, a finite subset $\mathcal{P} \subset (0, L)$, and a function $c : \mathcal{P} \rightarrow \mathbb{R}$ such that if we define

$$A_*(t) := \int_0^t a(s) ds$$

then

$$A(t) = A_*(t) + \sum_{p \in \Delta} c(p)H(t-p), \quad \forall t \in [0, L].$$

We denote by \mathcal{A}_* the subcollection of nice functions. □

Let us first point out that \mathcal{A}_* is a vector subspace of \mathcal{A} . Next, observe that $A \in \mathcal{A}_*$ if and only if there exists a finite subset $\mathcal{P}_A \subset (0, L)$ such that the restriction of A to $[0, L] \setminus \mathcal{P}$ is Lipschitz continuous. In this case A admits left and right limits at any point $t \in [0, L]$ and we define

$$c : \mathcal{P}_A \rightarrow \mathbb{R}, \quad c(p) = \lim_{t \searrow p} A(t) - \lim_{t \nearrow p} A(t).$$

Then

$$A_*(t) = A(t) - \sum_{p \in \mathcal{P}} c(p)H(t-p)$$

is Lipschitz continuous, it is differentiable a.e. on $[0, L]$ and we define a to be the derivative of A_* .

Let us next observe that if $A \in \mathcal{A}_*$ then the operator T_A can be informally described as

$$T_A = -i \frac{d}{dt} + a(t) + \sum_{p \in \mathcal{P}_A} c(p)\delta_p.$$

In other words, T_A would like to be a Dirac type operator whose coefficients are measures. In the above informal discussion we left out a description of the domain of T_A . Below we would like to give a precise description of T_A as a closed unbounded selfadjoint operator defined by an elliptic boundary value problem.

For any partition of $[0, L]$, $\mathcal{P} = \{0 < t_1 < \dots < t_{n-1} < L\}$, we set

$$t_0 := 0, \quad t_n := L, \quad I_k := [t_{k-1}, t_k], \quad k = 1, \dots, n.$$

We define the Hilbert space

$$\mathbf{H}_{\mathcal{P}} = \bigoplus_{k=1}^n L^2(I_k, \mathbb{C}),$$

and the Hilbert space isomorphism

$$\mathcal{J}_{\mathcal{P}} : \mathbf{H} \rightarrow \mathbf{H}_{\mathcal{P}}, \quad \mathbf{H} \ni f \mapsto (f|_{I_1}, \dots, f|_{I_n}) \in \mathbf{H}_{\mathcal{P}}.$$

Let $A \in \mathcal{A}_*$ and \mathcal{P} be a partition

$$\mathcal{P} = \{0 < t_1 < \dots < t_{n-1} < L\}$$

that contains the set of discontinuities of A , $\mathcal{P} \supset \mathcal{P}_A$. We set

$$a = \frac{dA_*}{dt}, \quad ; a_k = a|_{I_k}, \quad k = 1, \dots, n.$$

For $j = 1, \dots, n-1$ we denote by $c_j = c_j(A)$ the jump of A at t_j ,

$$c_j = A(t_j^+) - A(t_j^-).$$

Finally we define the closed unbounded linear operator

$$L_{\mathcal{A}, \mathcal{P}} : \text{Dom}(L_{\mathcal{A}, \mathcal{P}}) \subset \mathbf{H}_{\mathcal{P}} \rightarrow \mathbf{H}_{\mathcal{P}},$$

where $\text{Dom}(L_{\mathcal{A}, \mathcal{P}})$ consists of n -uples $(u_k)_{1 \leq k \leq n} \in \mathbf{H}_{\mathcal{P}}$ such that

$$u_k \in L^{1,2}(I_k), \quad k = 1, \dots, n, \tag{1.14a}$$

$$u_{j+1}(t_j) = e^{-ic_j} u_j(t_j), \quad j = 1, \dots, n-1, \tag{1.14b}$$

$$u_n(L) = u_1(0). \tag{1.14c}$$

and

$$L_{A,\mathcal{P}}(u_1, \dots, u_n) = \left(-i \frac{du_1}{dt} + a_1 u_1, \dots, -i \frac{du_n}{dt} + a_n u_n \right). \quad (1.15)$$

A standard argument shows that $L_{A,\mathcal{P}}$ is closed, densely defined and selfadjoint. In particular, the operator $(L_{A,\mathcal{P}} + i)$ is invertible, with bounded inverse.

Theorem 1.8. *For any $A \in \mathcal{A}_*$ and any partition*

$$\mathcal{P} = \{0 < t_1 < \dots < t_{n-1} < L\}$$

that contains the set of discontinuities of A we have the equality

$$L_{A,\mathcal{P}} = \mathcal{J}_{\mathcal{P}} T_A \mathcal{J}_{\mathcal{P}}^{-1}.$$

Proof. For simplicity we write L_A instead of $L_{A,\mathcal{P}}$. We will prove the equivalent statement

$$(i + L_A)^{-1} = \mathcal{J}_{\mathcal{P}} (T_A + i)^{-1} \mathcal{J}_{\mathcal{P}}^{-1} = \mathcal{J}_{\mathcal{P}} R_A \mathcal{J}_{\mathcal{P}}^{-1}.$$

In other words we have to prove that for any $u, f \in \mathbf{H}$ if $u = R_A f$, then $u \in \text{Dom}(L_A)$ and $(L_A + i)\mathcal{J}_{\mathcal{P}} u = \mathcal{J}_{\mathcal{P}} f$. More precisely, we have to show that the collection $\mathcal{J}_{A} u = (u_k)_{1 \leq k \leq n}$ satisfies (1.14a–1.14c) and (1.15). Using (1.2) we deduce

$$u(t) = \frac{i e^{-\Phi_A(t)}}{e^{\Phi_A(L)} - 1} \int_0^L e^{\Phi_A(s)} f(s) ds + i e^{-\Phi_A(t)} \int_0^t e^{\Phi_A(s)} f(s) ds. \quad (1.16)$$

This implies the condition (1.14a). The condition (1.15) follows by direct computation using (1.16).

Next, we observe that

$$\begin{aligned} u(t_j^+) &= \frac{i e^{-\Phi_A(t_j^+)}}{e^{\Phi_A(L)} - 1} \int_0^L e^{\Phi_A(s)} f(s) ds + i e^{-\Phi_A(t_j^+)} \int_0^{t_j} e^{\Phi_A(s)} f(s) ds, \\ u(t_j^-) &= \frac{i e^{-\Phi_A(t_j^-)}}{e^{\Phi_A(L)} - 1} \int_0^L e^{\Phi_A(s)} f(s) ds + i e^{-\Phi_A(t_j^-)} \int_0^{t_j} e^{\Phi_A(s)} f(s) ds, \end{aligned}$$

from which we conclude that

$$u(t_j^+) = e^{-i(\Phi_A(t_j^+) - \Phi_A(t_j^-))} u(t_j^-), \quad \forall j = 1, \dots, n-1.$$

This proves (1.14b). The equality (1.14c) follows directly from (1.5). \square

Remark 1.9. We would like to place the above operator L_A in a broader perspective that we will use extensively in Section 4. Consider a compact, oriented 1-dimensional manifold with boundary I . In other words I is a disjoint union of finitely many compact intervals

$$\mathbf{I} = \sqcup_{k=1}^n I_k.$$

If $I_k := [a_k, b_k]$, $a_k < b_k$, then we set

$$\partial_+ I_k := \{b_k\}, \quad \partial_- I_k := \{a_k\}, \quad \partial_+ \mathbf{I} := \{b_1, \dots, b_n\}, \quad \partial_- \mathbf{I} := \{a_1, \dots, a_n\}.$$

In particular, we have a direct sum decomposition of (finite dimensional) Hilbert spaces

$$\mathbf{E} := L^2(\partial \mathbf{I}, \mathbb{C}) = L^2(\partial_+ \mathbf{I}) \oplus L^2(\partial_- \mathbf{I}) =: \mathbf{E}_+ \oplus \mathbf{E}_-.$$

On the space $C^\infty(\mathbf{I}, \mathbb{C})$ of smooth complex valued functions on \mathbf{I} we have a canonical, symmetric Dirac \mathcal{D} operator described on each I_k by $-i \frac{d}{dt}$. Let σ denote the principal symbol of this operator. If ν_* denotes the *outer* conormal to the boundary. We then get an operator

$$J = \sigma(\nu_*) : L^2(\partial \mathbf{I}, \mathbb{C}) \rightarrow L^2(\partial \mathbf{I}, \mathbb{C}).$$

It is a unitary operator satisfying $J^2 = -1$, $\ker(\mathbf{i} + J) = \mathbf{E}_+$, and $\ker(\mathbf{i} - J) = \mathbf{E}_-$. It thus defines a Hermitian symplectic structure in the sense of [1, 5, 14]. A (hermitian) lagrangian subspace of \mathbf{E} is then a complex subspace L such that $L^\perp = JL$. We denote by $\text{Lag}(\mathbf{E}, J)$ the Grassmannian of hermitian lagrangian spaces. We denote by $\text{Iso}(\mathbf{E}_+, \mathbf{E}_-)$ the space of linear isometries $\mathbf{E}_+ \rightarrow \mathbf{E}_-$. As explained in [1] there exists a natural bijection³

$$\text{Iso}(\mathbf{E}_+, \mathbf{E}_-) \rightarrow \text{Lag}(\mathbf{E}), \quad \text{Iso}(\mathbf{E}_+, \mathbf{E}_-) \ni T \longmapsto \Gamma_T$$

where Γ_T is the graph of T viewed as a subspace of \mathbf{E} . Our spaces \mathbf{E}_\pm are equipped with natural bases and through these bases we can identify $\text{Iso}(\mathbf{E}_+, \mathbf{E}_-)$ with the unitary group $U(n)$. We denote by Δ the Lagrangian subspace corresponding to the identity operator.

Any subspace $V \subset \mathbf{E}$ defines a Fredholm operator

$$\mathcal{D}_V : \text{Dom}(\mathcal{D}_V) \subset L^2(\mathbf{I}, \mathbb{C}) \rightarrow L^2(\mathbf{I}, \mathbb{C}),$$

where

$$\text{Dom}(\mathcal{D}_V) = \{ u \in L^{1,2}(\mathbf{I}, \mathbb{C}); \quad u|_{\partial \mathbf{I}} \in V \}, \quad \mathcal{D}_V u = \mathcal{D}u.$$

The index of this operator is

$$\begin{aligned} i_V &= \dim(V \cap \Delta) - \dim(\Delta \cap JV^\perp) = \dim(V \cap \Delta) - \dim(J\Delta \cap V^\perp) \\ &= \dim(V \cap \Delta) - \dim(\Delta^\perp \cap V^\perp) = \dim V - \text{codim } \Delta = \dim V - n. \end{aligned}$$

A simple argument shows that \mathcal{D}_V is selfadjoint if and only if $V \in \text{Lag}(\mathbf{E})$. As we explained above, in this case V can be identified with the graph of an isometry $T : \mathbf{E}_+ \rightarrow \mathbf{E}_-$. We say that T is the *transmission operator* associated to the selfadjoint boundary value problem.

For example, if in Theorem 1.8 we let $A(t) = \sum_{j=1}^{n-1} c_j H(t - t_j)$, then we see that the operator L_A can be identified with the operator \mathcal{D}_{Γ_T} , where the transmission operator $T \in \text{Iso}(\mathbf{E}_+, \mathbf{E}_-)$ is given by the unitary $n \times n$ matrix

$$T = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & 1 \\ e^{-ic_1} & 0 & 0 & \cdots & 0 & 0 \\ 0 & e^{-ic_2} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdots & e^{-ic_{n-1}} & 0 \end{bmatrix}.$$

□

2. The Atiyah-Patodi-Singer theorem

We review here the Atiyah-Patodi-Singer index theorem for Dirac operators on manifold with boundary, when the metric is not assumed to be cylindrical near the boundary. Our presentation follows closely, [8, 9], but we present a few more details since the various orientation conventions and the terminology in [8, 9] are different from those in [3, 13] that we use throughout this paper.

Suppose $(\widehat{M}, \widehat{g})$ is a compact, oriented Riemann, and $M \subset \widehat{M}$ be a hypersurface in \widehat{M} co-oriented by a unit normal vector field ν along M . Let $n := \dim M$ so that $\dim \widehat{M} = n + 1$. We denote by g the induced metric on M . We first want to define a canonical restriction to M of a Dirac operator on \widehat{M} .

³There are various conventions in the definition of this bijection. We follow the conventions in [5].

Let $\exp^{\hat{g}} : T\widehat{M} \rightarrow \widehat{M}$ denote the exponential map determined by the metric \hat{H} . For sufficiently small $\varepsilon > 0$ the map

$$(-\varepsilon, \varepsilon) \times M \ni (t, p) \mapsto \exp_p^{\hat{g}}(t\nu(p)) \in \widehat{M}$$

is a diffeomorphism onto a small open tubular neighborhood \mathcal{O}_ε of M . The metric g determines a cylindrical metric $dt^2 + g$ on $(-\varepsilon, \varepsilon) \times M$. Via the above diffeomorphism we get a metric \hat{g}_0 on \mathcal{O}_ε . We say that \hat{g}_0 is the *cylindrical approximation* of \hat{g} near M .

We denote by $\widehat{\nabla}$ the Levi-Civita connection of the metric \hat{g} and by $\widehat{\nabla}^0$ the Levi-Civita connection of the metric \hat{g}_0 . We set

$$\Xi := \widehat{\nabla} - \widehat{\nabla}^0 \in \Omega^1(\mathcal{O}_\varepsilon, \text{End}(T\widehat{M})).$$

To get a more explicit description of Ξ we fix a local oriented, g -orthonormal frame (e_1, \dots, e_n) on M . Together with the unit normal vector field ν we obtain a local oriented orthonormal frame (ν, e_1, \dots, e_n) of $T\widehat{M}|_M$. We extend it by parallel transport along the geodesics orthogonal to M to a local, oriented orthonormal frame $(\hat{\nu}, \hat{e}_1, \dots, \hat{e}_n)$ of $T\widehat{M}$.

Denote by $\hat{\omega}$ the connection form associated to $\widehat{\nabla}$ by this frame, and by $\hat{\theta}$ the connection form associated to $\widehat{\nabla}^0$ by this frame. We can represent both $\hat{\omega}$ and $\hat{\theta}$ as skew-symmetric $(n+1) \times (n+1)$ matrices

$$\hat{\omega} = (\hat{\omega}_j^i)_{0 \leq i, j \leq n}, \quad \hat{\theta} = (\hat{\theta}_j^i)_{0 \leq i, j \leq n},$$

where the entries are 1-forms. Then $\Xi = \hat{\omega} - \hat{\theta}$.

We set $\hat{e}_0 := \hat{\nu}$, and we denote by $(\hat{e}^k)_{0 \leq k \leq n}$ the dual orthonormal frame of $T^*\widehat{M}$. Then we have

$$\hat{\omega}_j^i = \hat{\omega}_{kj}^i \hat{e}^k, \quad \hat{\theta}_j^i = \hat{\theta}_{kj}^i \hat{e}^k, \quad \widehat{\nabla}_k \hat{e}_j = \hat{\omega}_{kj}^i \hat{e}_i, \quad \widehat{\nabla}_k^0 \hat{e}_j = \hat{\theta}_{kj}^i \hat{e}_i, \quad \forall 0 \leq j, k \leq n,$$

where we have used Einstein's summation convention.

Observe that $\widehat{\nabla}^0 \hat{e}_0 = 0$ so that $\hat{\theta}_0^i = \hat{\theta}_i^0 = 0$. Also,

$$\hat{\omega}_{jk}^i = \hat{\theta}_{jk}^i, \quad \forall 1 \leq i, j, k \leq n.$$

If we write

$$\Xi = (\Xi_j^i)_{0 \leq i, j \leq n}, \quad \Xi_j^i = \Xi_{jk}^i \hat{e}^k,$$

and we let $o(1)$ denote any quantity that vanishes along M . then we have

$$\Xi_j^i = -\Xi_i^j, \quad \forall 0 \leq i, j \leq n, \tag{2.1}$$

$$\Xi_{kj}^i = o(1), \quad \forall 1 \leq i, j \leq n, \quad 0 \leq k \leq n. \tag{2.2}$$

We set

$$\Xi_{kij} := \Xi_{kj}^i = \hat{g}(\widehat{\nabla}_k \hat{e}_j, \hat{e}_i), \quad \hat{\omega}_{ij} = \hat{\omega}_j^i, \quad \hat{\theta}_{ij} = \hat{\theta}_j^i, \quad \omega_{kij} = \omega_{kj}^i, \quad \theta_{kij} = \theta_{kj}^i.$$

We denote by Q the second fundamental form⁴ of the embedding $M \hookrightarrow \widehat{M}$,

$$Q(e_i, e_j) = g(\widehat{\nabla}_{e_i} \nu, e_j).$$

Along the boundary we have the equalities

$$\Xi_{kj0} = \Xi_{jk0} = -\Xi_{k0j} = Q(e_j, e_k) \quad \forall 1 \leq i, j \leq n, \tag{2.3a}$$

$$\Xi_{ij0} = 0, \quad \forall 0 \leq i, j \leq n. \tag{2.3b}$$

To understand the nature of the restriction to a hypersurface of a Dirac operator we begin with a special case. Namely, we assume that \widehat{M} is equipped with a *spin* structure. We denote by $\hat{\mathbb{S}}$ the

⁴Our definition of the second fundamental form differs by a sign from the usual definition. With our definition the round sphere cooriented by the outer normal has positive mean curvature.

associated complex spinor bundle so that $\widehat{\mathbb{S}}$ is $\mathbb{Z}/2$ -graded is $\dim \widehat{M}$ is even, and ungraded otherwise. We have a Clifford multiplication

$$\hat{c} : T^*M \rightarrow \text{End}(\widehat{\mathbb{S}}).$$

The metrics \hat{g} and \hat{g}_0 define connections $\widehat{\nabla}^{spin}$ and $\widehat{\nabla}^{spin,0}$ on $\widehat{\mathbb{S}}|_{\mathcal{O}_\varepsilon}$. Using the local frame $(\hat{e}_i)_{0 \leq i, j \leq n}$ we can write

$$\widehat{\nabla}_k^{spin} = \partial_k - \frac{1}{4} \hat{\omega}_{kij} \hat{c}(\hat{e}^i) \hat{c}(\hat{e}^j), \quad \widehat{\nabla}_k^{spin,0} = \partial_k - \frac{1}{4} \hat{\theta}_{kij} \hat{c}(\hat{e}^i) \hat{c}(\hat{e}^j),$$

where we again use Einstein's summation convention.

Using the connections $\widehat{\nabla}^{spin}$ and $\widehat{\nabla}^{spin,0}$ we obtain two Dirac operators \hat{D} and respectively \hat{D}_0 on $\widehat{\mathbb{S}}|_{\mathcal{O}_\varepsilon}$

$$\hat{D} = \sum_{i=0}^n \hat{c}(\hat{e}^i) \widehat{\nabla}_i^{spin}, \quad \hat{D}_0 = \sum_{i=0}^n \hat{c}(\hat{e}^i) \widehat{\nabla}_i^{spin,0}.$$

Identifying \mathcal{O}_ε with $(-\varepsilon, \varepsilon) \times M$ we obtain a projection

$$\pi : \mathcal{O}_\varepsilon \rightarrow M.$$

We set $\mathbb{S} := \widehat{\mathbb{S}}|_M$. The parallel transport given by $\widehat{\nabla}^{spin}$ yields a bundle isomorphism $\widehat{\mathbb{S}}|_{\mathcal{O}_\varepsilon} \cong \pi^*\mathbb{S}$. Using these identifications we can rewrite the operators \hat{D} and \hat{D}_0 as

$$\begin{aligned} \hat{D} &= \hat{c}(\hat{e}_0) (\widehat{\nabla}_0^{spin} - D(t)) : C^\infty(\pi^*\mathbb{S}) \rightarrow C^\infty(\pi^*\mathbb{S}), \\ \hat{D}_0 &= \hat{c}(\hat{e}^0) (\partial_0 - D_0(t)) : C^\infty(\pi^*\mathbb{S}) \rightarrow C^\infty(\pi^*\mathbb{S}). \end{aligned}$$

The operators $D(t)$ and $D_0(t)$ are first order differential operators $C^\infty(\widehat{\mathbb{S}}|_{\{t\} \times M}) \rightarrow C^\infty(\widehat{\mathbb{S}}|_{\{t\} \times M})$ and thus can be viewed as t -dependent operators on \mathbb{S} .

The operator $D_0(t)$ is in fact independent of t and thus we can identify it with a Dirac operator on $C^\infty(\mathbb{S}) \rightarrow C^\infty(M)$. It is called the *canonical restriction* of \hat{D} to M , and we will denote it by $\mathcal{R}_M(\hat{D})$. This operator is intrinsic to M . More precisely when $\dim \widehat{M}$ is even then \mathbb{S} is the direct sum of two copies of the spinor bundle on M and the operator D_0 is the direct sum of two copies of the *spin*-Dirac operator determined by the Riemann metric on M .

When $\dim \widehat{M}$ is odd then \mathbb{S} is the spinor bundle on M and D_0 is the *spin*-Dirac operator determined by the metric on the boundary and the induced *spin* structure. We would like to express $\mathcal{R}_M(\hat{D})$ in terms of $D(t)|_{t=0}$.

Let $\nu_* := \hat{e}^0 \in C^\infty(T^*\widehat{M}|_{\partial\widehat{M}})$, set $J := \hat{c}(\nu_*)$ and define $c : T^*M \rightarrow \text{End}(\mathbb{S})$ by setting

$$c(\alpha) = \hat{c}(\nu_*(p)) \hat{c}(\alpha) = J \hat{c}(\alpha), \quad \forall p \in M, \quad \alpha \in T^*M \subset (T^*\widehat{M})|_M.$$

Observe first that

$$\mathcal{R}_M(\hat{D}) = D_0(t) = \partial_0 + J \hat{D}_0.$$

Next we observe that

$$\widehat{\nabla}^{spin} - \widehat{\nabla}^{spin,0} = -\frac{1}{4} \sum_{i,j} \Xi_{ij} \hat{c}(\hat{e}^i) \hat{c}(\hat{e}^j).$$

so that

$$\begin{aligned} \widehat{\nabla}_0^{spin} - \widehat{\nabla}_0^{spin,0} &= \widehat{\nabla}_0^{spin} - \partial_0 = -\frac{1}{4} \Xi_{0ij} J \hat{c}(\hat{e}^i) \hat{c}(\hat{e}^j) = o(1), \\ \hat{D} - \hat{D}_0 &= -\frac{1}{4} \sum_{i,j,k} \Xi_{kij} \hat{c}(\hat{e}^k) \hat{c}(\hat{e}^i) \hat{c}(\hat{e}^j) =: \mathcal{L}. \end{aligned}$$

We denote by $\mathcal{L}(t)$ the restriction of \mathcal{L} to the slice $\{t\} \times M$ so that $\mathcal{L}(t)$ is an endomorphism of $\widehat{\mathbb{S}}|_{\{t\} \times M}$.

Hence

$$\hat{D} = J\partial_0 - JD(t), \quad D(t) = J\hat{D} + \partial_0 = J\hat{D}_0 + \partial_0 + J\mathcal{L} = D_0(t) + J\mathcal{L},$$

so that

$$D(0) = \mathcal{R}_M(\hat{D}) + J\mathcal{L}(t)|_{t=0}.$$

Thus, we need to compute the endomorphism $J\mathcal{L}(t)|_{t=0}$. We have

$$J\mathcal{L} = -\frac{1}{4} \sum_{i,j,k} J\Xi_{kij} \hat{c}(e^k) \hat{c}(e^i) \hat{c}(e^j).$$

There are many cancellations in the above sum. Using (2.2) we deduce that the terms corresponding to $k = 0$ vanish. Using (2.1) we deduce that the terms corresponding to $i, j > 0$ or $i = j$ also vanish along the boundary. Thus

$$\begin{aligned} J\mathcal{L} &= -\frac{1}{4} \sum_{i \neq j, k \neq 0} \Xi_{kij} J\hat{c}(e^k) \hat{c}(e^i) \hat{c}(e^j) + o(1) = -\frac{1}{2} \sum_{i > j, k > 0} \Xi_{kij} J\hat{c}(e^k) \hat{c}(e^i) \hat{c}(e^j) + o(1) \\ &= -\frac{1}{2} \sum_{i > 0, k > 0} \Xi_{ki0} J\hat{c}(e^k) \hat{c}(e^i) \hat{c}(e^0) + o(1). \end{aligned}$$

Using the equalities $J = \hat{c}(e^0)$, $J\hat{c}(e^\ell) = -\hat{c}(e^\ell)J$ for $\ell > 0$ we deduce

$$J\mathcal{L} = \frac{1}{2} \sum_{i,k > 0} \Xi_{ki0} \hat{c}(e^k) \hat{c}(e^i) = -\frac{1}{2} \sum_{j > 0} \Xi_{ii0} + o(1) = -\frac{1}{2} \text{tr } Q.$$

The scalar $\text{tr } Q$ is the mean curvature of $M \hookrightarrow \widehat{M}$ and we denote it by h_M . Hence

$$D(t)|_{t=0} = \mathcal{R}_M(\hat{D}) - \frac{1}{2}h_M. \quad (2.4)$$

A similar equality was proved in [12, Lemma 4.5.1], although in [12] they use a different definition for the induced Clifford multiplication on the boundary that leads to some sign differences.

If now $\widehat{E} \rightarrow \widehat{M}$ is a hermitian vector bundle over \widehat{M} and $\widehat{\nabla}^E$ is a Hermitian connection on \widehat{E} then we obtain in standard fashion a twisted Dirac operator $\widehat{D}_E : C^\infty(\widehat{\mathbb{S}} \otimes \widehat{E}) \rightarrow C^\infty(\widehat{\mathbb{S}} \otimes \widehat{E})$. Using the parallel transport given by $\widehat{\nabla}^E$ we obtain an isomorphism

$$\widehat{E}|_{\mathcal{O}_\varepsilon} \cong \pi^*E, \quad \text{where } E := \widehat{E}|_M.$$

Along \mathcal{O}_ε the operator \widehat{D}_E has the form

$$\widehat{D}_E = J(\partial_t - D_E(t)).$$

If on \mathcal{O}_ε we replace the metric \hat{g} with its cylindrical approximation \hat{g}_0 we obtain a new Dirac operator

$$\widehat{D}_{E,0} : C^\infty(\pi^*(\mathbb{S} \otimes E)) \rightarrow C^\infty(\pi^*(\mathbb{S} \otimes E))$$

which along the boundary has the form $J(\partial_t - D_{E,0})$, where $D_{E,0} : C^\infty(\mathbb{S} \otimes E) \rightarrow C^\infty(\mathbb{S} \otimes E)$. We set $\mathcal{R}_M(\widehat{D}_E) := D_{E,0}$ and as before we obtain the identity

$$D_E(t)|_{t=0} = \mathcal{R}_M(\widehat{D}_E) - \frac{1}{2}h_M. \quad (2.5)$$

This is a purely local result so that a similar formula holds for the geometric Dirac operators determined by a $spin^c$ structure.

We want to apply the above discussion to a very special case. Consider a compact oriented surface Σ with possibly disconnected boundary $\partial\Sigma$. We think of $\partial\Sigma$ as a hypersurface in Σ cooriented by the outer normal.

Fix a Riemann metric \hat{g} on Σ , smooth up to the boundary. Denote by s the arclength coordinate on a component $\partial_0\Sigma$ of the boundary. As before we can identify an open neighborhood \mathcal{O} of this component of the boundary with a cylinder $(-\varepsilon, 0] \times S^1$. In this neighborhood the metric \hat{g} has the form

$$\hat{g} = dt^2 + w^2 ds^2$$

where $w : (-\varepsilon, 0] \times S^1 \rightarrow (0, \infty)$ is a smooth positive function in the variables t, s such that $w(0, s) = 1, \forall s$.

The metric and the orientation on Σ defines an integrable almost complex structure $J : T\Sigma \rightarrow T\Sigma$. More precisely, J is given by the counterclockwise rotation by $\pi/2$. We denote by K_Σ the canonical complex line bundle determined by J . We get a Dolbeault operator

$$(\bar{\partial} + \bar{\partial}^*) : C^\infty(\underline{\mathbb{C}}_\Sigma \oplus K_\Sigma^{-1}) \rightarrow C^\infty(\underline{\mathbb{C}}_\Sigma \oplus K_\Sigma^{-1}).$$

We regard this as the Dirac operator defined by the metric \hat{g} , a $spin^c$ structure. The twisting line bundle is $K_\Sigma^{-1/2}$, where the connection on K_Σ is the connection induced by the Levi-Civita connection of the metric \hat{g} . We analyze the form of $\bar{\partial} : C^\infty(\underline{\mathbb{C}}_\Sigma) \rightarrow C^\infty(K_\Sigma^{-1})$ on the cylindrical neighborhood \mathcal{O} . We set

$$e^0 = dt, \quad e^1 = w ds.$$

Then $\{e^0, e^1\}$ is an oriented, orthonormal frame of $T^*\Sigma|_{\mathcal{O}}$. We denote by $\{e_0, e_1\}$ its dual frame of $T\Sigma$. We let $c : T^*\Sigma \rightarrow \text{End}(\underline{\mathbb{C}}_\Sigma \oplus K_\Sigma)$ be the Clifford multiplication normalized by the condition that the operator $dV := c(e^0)c(e^1)$ on $\underline{\mathbb{C}}_\Sigma \oplus K_\Sigma^{-1}$ has the block decomposition [3, §3.2],

$$c(e^0)c(e^1) = \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}. \quad (2.6)$$

The Levi-Civita induces a natural connection on K_Σ^{-1} and if we use the trivial connection on $\underline{\mathbb{C}}_\Sigma$ we get a connection ∇ on $\underline{\mathbb{C}}_\Sigma \oplus K_\Sigma^{-1}$. The associated Dirac operator is $D_\Sigma = \sqrt{2}(\bar{\partial} + \bar{\partial}^*)$. The even part of this operator is

$$D^+ = \sqrt{2}\bar{\partial} : C^\infty(\underline{\mathbb{C}}_\Sigma) \rightarrow C^\infty(K_\Sigma^{-1}).$$

We want to compute its canonical restriction to the boundary.

The Levi-Civita connection $\widehat{\nabla}$ determined by \hat{g} is described on \mathcal{O} by a 1-form ω uniquely determined by Cartan's structural equations

$$\widehat{\nabla} = d + \begin{bmatrix} 0 & -\omega \\ \omega & 0 \end{bmatrix}, \quad d \begin{bmatrix} e^0 \\ e^1 \end{bmatrix} = \begin{bmatrix} 0 & \omega \\ -\omega & 0 \end{bmatrix} \wedge d \begin{bmatrix} e^0 \\ e^1 \end{bmatrix}.$$

We deduce $\omega = ae^1$, $a \in C^\infty(\mathcal{O})$ and from the equality

$$\frac{w'_t}{w} e^0 \wedge e^1 = w'_t dt \wedge ds = de^1 = ae^0 \wedge e_1$$

we conclude $a = \partial_t \log w$ so that

$$\omega = \partial_t(\log w)e^1 = w'_t ds.$$

The mean curvature h of the boundary component $\partial_0\Sigma$ is the restriction to $t = 0$ of the function w'_t . The Riemann curvature is described by the matrix

$$\begin{bmatrix} 0 & -d\omega \\ d\omega & 0 \end{bmatrix} = \begin{bmatrix} 0 & -w'_t dt \wedge ds \\ w'_t dt \wedge ds & 0 \end{bmatrix} = \begin{bmatrix} 0 & -\frac{w''_t}{w} \\ \frac{w''_t}{w} & 0 \end{bmatrix} e^0 \wedge e^1.$$

If we denote by ∂ the trivial connection on $\underline{\mathbb{C}}_\Sigma$ then we deduce

$$D_\Sigma^+ := c(e^0)\partial_{e_0} + c(e^1)\partial_{e_1} = c(e_0)(\partial_t - c(e^0)c(e^1)\partial_{e_1})$$

so that

$$D_{\Sigma}^{\pm}(t) = \mathbf{c}(e^0)D_{\Sigma}^{\pm} + \partial_t = \mathbf{c}(e^0)\mathbf{c}(e^1)\partial_{e_1} \stackrel{(2.6)}{=} -i\partial_{e_1}.$$

Above, the operator $D_{\Sigma}^{\pm}(t)$ is, *canonically*, a differential operator

$$D_{\Sigma}^{\pm}(t) : C^{\infty}(\underline{\mathbb{C}}_{\partial\Sigma}) \rightarrow C^{\infty}(\underline{\mathbb{C}}_{\partial\Sigma}),$$

where $\underline{\mathbb{C}}_{\partial\Sigma}$ denotes the trivial complex line bundle over $\partial\Sigma$. The boundary restriction is then according to (2.5)

$$\mathcal{R}_{\partial\Sigma}(\bar{\partial}) = D_{\Sigma}^{\pm}(t) + \frac{1}{2}h = -i\partial_{e_1} + \frac{1}{2}h. \quad (2.7)$$

Let us observe that along the boundary we have $\partial_{e_1} = \partial_s$.

Consider the Atiyah-Patodi-Singer operator

$$\bar{\partial}_{APS} : \text{Dom}(\bar{\partial}_{APS}) \subset L^2(\Sigma) \rightarrow L^2(\Sigma), \quad \bar{\partial}_{APS}u = \bar{\partial}u, \forall u \in \text{Dom}(\bar{\partial}_{APS}),$$

where

$$\text{Dom}(\bar{\partial}_{APS}) = \{u \in L^{1,2}(\Sigma, \mathbb{C}); u|_{\partial\Sigma} \in \Lambda_{\bar{\partial}}^{-}\},$$

and $\Lambda_{\bar{\partial}}^{-}$ is the closed subspace of $L^2(\partial\Sigma)$ generated by the eigenvectors of the operator $B := \mathcal{R}_{\partial\Sigma}(\bar{\partial})$ corresponding to strictly negative eigenvalues.

The index theorem of [8, 9] implies $\bar{\partial}_{APS}$ is Fredholm and

$$i_{APS}(\Sigma, g) := \text{index}(\bar{\partial}_{APS}) = \frac{1}{2} \int_{\Sigma} c_1(\Sigma, g) - \xi_B, \quad \xi_B := \frac{1}{2}(\dim B + \eta_B(0)).$$

Above, $c_1(\Sigma, g) \in \Omega^2(\Sigma)$ is the 2-form $\frac{1}{2\pi}K_g dV_g$, where K_g denotes the sectional curvature of g and dV_g denotes the metric volume form on Σ . From the Gauss-Bonnet theorem for manifolds with boundary [15, §6.6] we deduce

$$\int_{\Sigma} c_1(\Sigma, g) + \frac{1}{2\pi} \int_{\partial\Sigma} h ds = \chi(\Sigma),$$

where $h : \partial\Sigma \rightarrow \mathbb{R}$ is the mean curvature function defined as above. We deduce

$$i_{APS}(\Sigma, g) = \frac{1}{2}\chi(\Sigma) - \frac{1}{4\pi} \int_{\Sigma} h ds - \xi_B. \quad (2.8)$$

If $\partial\Sigma$ has several components $\partial\Sigma = \partial_1\Sigma \sqcup \dots \sqcup \partial_n\Sigma$, then we have n scalars

$$H_i = \frac{1}{4\pi} \int_{\partial_i\Sigma} h ds,$$

and a direct sum decomposition $B = \oplus_{i=1}^n B_i$, where each of the operators B_i is described by (2.7). We set

$$\rho_i = H_i - [H_i], \quad i = 1, \dots, n.$$

Then using (2.7) and (1.10) we deduce

$$\xi_{B_i} = \frac{1}{2}(\dim \ker B_i + \eta_{B_i}(0)) = \frac{1}{2}(1 - 2\rho_i).$$

We can rewrite (2.8) as

$$i_{APS}(\Sigma, g) = \frac{1}{2}\chi(\Sigma) - \sum_{i=1}^n H_i - \frac{1}{2} \sum_{i=1}^n (1 - 2\rho_i) = \frac{1}{2}(\chi(\Sigma) - n) - \sum_{i=1}^n [H_i]. \quad (2.9)$$

3. Dolbeault operators on two-dimensional cobordisms

When thinking of cobordisms we adopt the Morse theoretic point of view. For us an elementary (nontrivial) 2-dimensional cobordism will be a pair (Σ, f) where Σ is a compact, *connected*, oriented surface with boundary, $f : \Sigma \rightarrow \mathbb{R}$ is a Morse function with a unique critical point p_0 located in the interior of Σ such that

$$f(\Sigma) = [-1, 1], \quad f(\partial\Sigma) = \{-1, 1\}, \quad f(p_0) = 0.$$

In more intuitive terms, an elementary cobordism looks like one of the two pair of pants in Figure 1, where the Morse function is understood to be the altitude.




FIGURE 1. Elementary 2-dimensional cobordisms.

We set

$$\partial_{\pm}\Sigma := f^{-1}(\pm 1).$$

In the sequel, for simplicity, we will assume that $\partial_+\Sigma$ is connected, i.e., the pair (Σ, f) looks like the left-hand-side of Figure 1.

We fix a Riemann metric g on Σ . For simplicity⁵ we assume that in an open neighborhood \mathcal{O} near p_0 there exist local coordinates such that, in these coordinates we have

$$g = dx^2 + dy^2, \quad f(x, y) = -\alpha x^2 + \beta y^2, \quad (3.1)$$

where α, β are positive constants. We let ∇f denote the gradient of f with respect to this metric and we set

$$C_t := f^{-1}(t), \quad t \neq 0.$$

For $t \neq 0$ we regard C_t cooriented by the gradient ∇f . Observe that C_t has two connected components when $t < 0$. We let $h_t : C_t \rightarrow \mathbb{R}$ be the mean curvature of this cooriented surface. For $t \neq 0$ we set

$$L_t = \int_{C_t} ds = \text{length}(C_t), \quad \omega_t := \frac{1}{4\pi} \int_{C_t} h_t ds.$$

Observe that even the singular level set C_0 is equipped with a natural measure defined by the arclength measure on $C_0 \setminus \{0\}$. The length of C_0 is finite since in a neighborhood of the singular point p_0 the level set isometric to a pair of intersecting line segments in an Euclidean space.

Denote by W^{\pm} the stable/unstable manifolds of p_0 with respect to the flow Φ^t generated by $-\nabla f$. The unstable manifold intersects the region $\{-1 \leq f < 0\}$ in two smooth paths (see Figure 2)

$$[-1, 0) \ni t \mapsto a_t, \quad b_t \in C_t, \quad \forall t \in [-1, 0),$$

while the stable manifold intersects the region $\{0 < f \leq 1\}$ in two smooth paths (the top red arcs in Figure 2)

$$(0, 1] \ni t \mapsto a_t, \quad b_t \in C_t, \quad \forall t \in (0, 1].$$

⁵The results to follow do not require the simplifying assumption (3.1) but the computations would be less transparent.

Observe that $\lim_{t \rightarrow 0} a_t = \lim_{t \rightarrow 0} b_t = p_0$. For this reason we set $a_0 = b_0 = p_0$.




FIGURE 2. Cutting an elementary 2-dimensional cobordism.

As we have mentioned before, for $t < 0$ the level set C_t consists of two curves. We denote by C_t^a the component containing the point a_t and by C_t^b the component containing b_t . For $t < 0$ we set

$$L_t^a := \int_{C_t^a} ds, \quad L_t^b := \int_{C_t^b} ds, \quad \omega_t^a := \frac{1}{4\pi} \int_{C_t^a} h_t ds, \quad \omega_t^b := \frac{1}{4\pi} \int_{C_t^b} h_t ds$$

so that

$$L_t = L_t^a + L_t^b, \quad \omega_t = \omega_t^a + \omega_t^b, \quad \forall t < 0.$$

Fix a point $\bar{a}_{-1} \in C_{-1}^a \setminus \{a_{-1}\}$ and a point $\bar{b}_{-1} \in C_{-1}^b \setminus \{b_{-1}\}$. For $t \in [-1, 1]$ we denote by \bar{a}_t (respectively \bar{b}_t) the intersection of C_t with the negative gradient flow line through \bar{a}_{-1} (respectively \bar{b}_{-1}). We obtain in this fashion two smooth maps (see Figure 2)

$$\bar{a}, \bar{b} : [-1, 1] \rightarrow \Sigma.$$

For $t > 0$ we denote by I_t^a the component of $C_t \setminus \{a_t, b_t\}$ that contains the point \bar{a}_t and by I_t^b the component of $C_t \setminus \{a_t, b_t\}$ that contains the point \bar{b}_t .

The regular part $C_0^* = C_0 \setminus \{p_0\}$ consists of two components C_0^a and C_0^b . We set

$$\frac{1}{4\pi} \omega_0^a := \frac{1}{4\pi} \int_{C_0^a} h_0 ds, \quad \omega_0^b := \frac{1}{4\pi} \int_{C_0^b} h_0 ds, \quad \omega_0 := \frac{1}{4\pi} \int_{C_0^*} h_0 ds = \omega_0^a + \omega_0^b. \quad (3.2)$$

Note that the limits $\lim_{t \rightarrow 0} L_t^a, \lim_{t \rightarrow 0} L_t^b$ exist and are finite. We denote them by L_0^a and respectively L_0^b . We have

$$L_0^a + L_0^b = L_0 := \text{length}(C)_0.$$

Let D_t denote the restriction of $\bar{\partial}$ to the cooriented curve C_t , $t \neq 0$. As explained in the previous section we have

$$D_t = \begin{cases} -i \frac{d}{ds} + \frac{1}{2} h_t, & t > 0, \\ (-i \frac{d}{ds} + \frac{1}{2} h_t)|_{C_t^a} \oplus (-i \frac{d}{ds} + \frac{1}{2} h_t)|_{C_t^b}, & t < 0. \end{cases}$$

If we set

$$\rho_t = \omega_t - [\omega_t], \quad \rho_t^a = \omega_t - [\omega_t^a], \quad \rho_t^b = \omega_t - [\omega_t^b],$$

then the computations in Section 1 imply

$$\xi(t) := \xi_{D_t} = \frac{1}{2} \begin{cases} 1 - 2\rho_t, & t > 0 \\ (1 - 2\rho_t^a) + (1 - 2\rho_t^b), & t < 0. \end{cases} \quad (3.3)$$

Throughout this and the next section we assume that both $D_{\pm 1}$ and are invertible.

We organize the family of complex Hilbert spaces $L^2(C_t, ds; \mathbb{C})$, $t \in [-1, 1]$ as a trivial bundle of Hilbert spaces as follows.

First observe that $C_0 \setminus \{\bar{a}_0, \bar{b}_0, p_0\}$ is a disjoint union of four open arcs I_1, \dots, I_4 labeled as in Figure 2. Denote by ℓ_j the length of I_j so that

$$L_0 = \ell_1 + \dots + \ell_4, \quad L_0^a = \ell_1 + \ell_4, \quad L_0^b = \ell_2 + \ell_3.$$

For $t > 0$ we can isometrically identify the oriented open arc $C_t \setminus \bar{a}_t$ with the open interval $(0, L_t)$. We obtain in this fashion a canonical isomorphism

$$\mathcal{J}_t^+ := L^2(C_t, ds; \mathbb{C}) \rightarrow L^2(0, L_t; \mathbb{C}).$$

The rescaling

$$(0, L_0) \rightarrow (0, L_t), \quad (0, L_0) \ni t \mapsto \frac{t}{\lambda_t}, \quad \lambda_t = \frac{L_0}{L_t}$$

induces a Hilbert space isomorphism

$$\mathcal{R}_t^+ : L^2(0, L_t; \mathbb{C}) \rightarrow L^2(0, L_0; \mathbb{C}) =: \mathbf{H}_0.$$

Note that we have a partition \mathcal{P}_+ of $[0, L_0]$

$$0 = t_0 < t_1 < t_2 < t_3 < t_4 = L_0, \quad t_j - t_{j-1} = \ell_j, \quad \forall j = 1, \dots, 4. \quad (3.4)$$

This defines a Hilbert space isomorphism

$$\mathcal{U}_+ : L^2(0, L_0; \mathbb{C}) \rightarrow \bigoplus_{j=1}^4 L^2(t_{j-1}, t_j; \mathbb{C}) = \bigoplus_{j=1}^4 L^2(I_j, ds; \mathbb{C}) =: \mathbf{H}_0.$$

For $t < 0$ we have

$$L^2(C_t, ds; \mathbb{C}) = L^2(C_t^a, ds; \mathbb{C}) \oplus L^2(C_t^b, ds; \mathbb{C}).$$

By removing the points \bar{a}_t and \bar{b}_t we obtain Hilbert space isomorphisms

$$L^2(C_t^a, ds; \mathbb{C}) \rightarrow L^2(0, L_t^a; \mathbb{C}), \quad L^2(C_t^b, ds; \mathbb{C}) \rightarrow L^2(0, L_t^b; \mathbb{C})$$

that add up to a Hilbert space isomorphism

$$\mathcal{J}_t^- : L^2(C_t, ds; \mathbb{C}) \rightarrow L^2(0, L_t^a; \mathbb{C}) \oplus L^2(0, L_t^b; \mathbb{C}).$$

By rescaling we obtain a Hilbert space isomorphism

$$\mathcal{R}_t^- : L^2(0, L_t^a; \mathbb{C}) \oplus L^2(0, L_t^b; \mathbb{C}) \rightarrow L^2(0, L_0^a; \mathbb{C}) \oplus L^2(0, L_0^b; \mathbb{C}) \cong L^2(0, L_0; \mathbb{C}).$$

Next observe that we have isomorphisms

$$\begin{aligned} \mathcal{U}_-^a : L^2(0, L_0^a; \mathbb{C}) &\rightarrow L^2(I_1, ds; \mathbb{C}) \oplus L^2(I_4, ds; \mathbb{C}), \\ \mathcal{U}_-^b : L^2(0, L_0^b; \mathbb{C}) &\cong L^2(I_3, ds; \mathbb{C}) \oplus L^2(I_2, ds; \mathbb{C}), \end{aligned}$$

that add up to an isomorphisms

$$\mathcal{U}_- : L^2(0, L_0; \mathbb{C}) \rightarrow \bigoplus_{j=1}^4 L^2(I_j, ds; \mathbb{C}).$$

For $t = 0$ we let \mathcal{J}_0 be the natural isomorphism

$$\mathcal{J}_0 : L^2(C_0, ds; \mathbb{C}) \rightarrow \bigoplus_{j=1}^4 L^2(I_j, ds; \mathbb{C}) \cong \mathbf{H}_0.$$

Now define

$$\mathcal{J}_t := \begin{cases} \mathcal{U}_+ \mathcal{R}_t^+ \mathcal{J}_t^+, & t > 0, \\ \mathcal{U}_- \mathcal{R}_t^- \mathcal{J}_t^-, & t < 0, \\ \mathcal{J}_0, & t = 0. \end{cases}$$

We use the collection of isomorphisms \mathcal{J}_t organizes the collection $L^2(C_t, ds; \mathbb{C})$ as a trivial Hilbert \mathcal{H} bundle over $[-1, 1]$.

Remark 3.1. Let us observe that any continuous function $f : \Sigma \rightarrow \mathbb{C}$ induces elements $f|_{C_t} \in L^2(C_t, ds; \mathbb{C})$, $\forall t \in [-1, 1]$ which in turn define a continuous section of the trivial Hilbert bundle \mathcal{H} . \square

Theorem 3.2. (a) The operators $\mathcal{D}_t := \mathcal{J}_t D_t \mathcal{J}_t^{-1}$ converge in the gap topology as $t \rightarrow 0^\pm$ to Fredholm, selfadjoint operators \mathcal{D}_0^\pm .

(b) The eta invariants of \mathcal{D}_0^\pm exist, and we set

$$\xi_\pm := \frac{1}{2} \left(\dim \ker \mathcal{D}_0^\pm + \eta_{\mathcal{D}_0^\pm}(0) \right),$$

If $\ker \mathcal{D}_0^\pm = 0$ then we have⁶

$$i_{APS}(\bar{\partial}) + \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t; \varepsilon < t \leq 1) + \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t, -1 \leq t < -\varepsilon) = -(\xi_+ - \xi_-). \quad (3.5)$$

Proof. We set

$$\mathcal{S}_t := \begin{cases} \mathcal{U}_+^{-1} \mathcal{D}_t \mathcal{U}_+, & t > 0 \\ \mathcal{U}_-^{-1} \mathcal{D}_t \mathcal{U}_-, & t < 0. \end{cases}$$

To establish the convergence statements we show that the limits $\lim_{t \rightarrow 0^\pm} \mathcal{S}_t$ exist in the gap topology of the space of unbounded selfadjoint operators on $L^2(0, L_0; \mathbb{C})$. We discuss separately the cases $\pm t > 0$, corresponding to restrictions to level sets above/below the critical level set $\{f = 0\}$.

A. $t > 0$. We observe that

$$\text{Dom}(\mathcal{S}_t) = \left\{ u \in L^{1,2}(0, L_0; \mathbb{C}); u(L_0) = u(0) \right\}, \quad \mathcal{S}_t(u) = -i\lambda_t \frac{d}{ds} + \frac{1}{2} h_t(s/\lambda_t),$$

where we recall that the constant λ_t is the rescaling factor L_0/L_t . We set

$$A_t(s) := \frac{1}{\lambda_t} \int_0^s h_t(\sigma/\lambda_t) d\sigma$$

Using the fact that $\lambda_t \rightarrow 1$ and Proposition 1.3 we see that it suffices to show that A_t is very weakly convergent in \mathcal{A}_{L_0} ; see Definition 1.1. Thus it suffices to prove two things.

$$\text{The limit } \lim_{t \rightarrow 0^+} A_t(L_0) \text{ exists.} \quad (\mathbf{A}_1)$$

$$\text{The limits } \lim_{t \rightarrow 0^+} A_t(s) \text{ exists for almost any } s \in (0, L_0). \quad (\mathbf{A}_2)$$

⁶The condition $\ker \mathcal{D}_0^\pm = 0$ is satisfied for an open and dense set of metrics g satisfying (3.1). When this condition is violated the identity (3.5) needs to be slightly modified to take into account these kernels.

Proof of (A₁). Observe that

$$A_t(L_0) = \int_{C_t} h_t ds = \int_{C_t \setminus \mathcal{O}} h_t ds + \int_{\mathcal{O} \cap C_t} h_t ds,$$

where \mathcal{O} is the neighborhood where (3.1) holds. The intersection of C_t with \mathcal{O} is depicted in Figure 3.




FIGURE 3. The behavior of C_t near the critical point.

The integral $\int_{C_t \setminus \mathcal{O}} h_t ds$ converges as $t \rightarrow 0^+$ to $\int_{C_0 \setminus \mathcal{O}} h_0 ds$. Next observe that the intersection $C_t \cap \mathcal{O}$ consists of two oriented arcs (see Figure 3) and the integral $\int_{\mathcal{O} \cap C_t} h_t$ computes the total angular variation of the oriented unit tangent vector field along these oriented arcs. Using the notations in Figure 3 we see that this total variation approaches $-2\theta_+$ as $t \rightarrow 0^+$. Hence

$$\lim_{t \rightarrow 0^+} A_t(L_0) = \int_{C_0 \setminus \mathcal{O}} h_0 ds - 2\theta_+,$$

so that

$$\omega_0^+ = \lim_{t \rightarrow 0^+} \omega_t = \frac{1}{4\pi} \lim_{t \rightarrow 0^+} \int_{C_t} h_t ds = \omega_0 - \frac{\theta_+}{2\pi}. \quad (3.6)$$

Proof of (A₂). Let $C_t^* := C_t \setminus \{\bar{a}_t\}$ and define $s = s(q) : C_t^* \rightarrow (0, \infty)$ to be the coordinate function on C_t^* such that the resulting map

$$C_t^* \rightarrow \mathbb{R}, \quad q \mapsto \sigma(q) = s(q)/\lambda_t$$

is an orientation preserving isometry onto $(0, L_t)$. In other words σ is the oriented arclength function measured starting at \bar{a}_t , and s defines a diffeomorphism $C_t^* \rightarrow (0, L_0)$. Let $q_t : (0, L_0) \rightarrow C_t^*$ be the inverse of this diffeomorphism.

Consider the partition (3.4). Observe that there exists positive constants c and ε such that whenever

$$\forall t \in (0, \varepsilon), \quad \forall s \in [t_1 - c, t_1 + c] \cup [t_3 - c, t_3 + c] : \quad q_t(s) \in \mathcal{O}$$

the numbers t_j are defined by (3.4). Intuitively the intervals $[t_1 - c, t_1 + c] \cup [t_3 - c, t_3 + c]$ collect the parts of C_t that are close to the critical point p_0 . The length of each of the two components of C_t that are close to p_0 is bounded from below by $2c/\lambda_t$.

To prove part (b) it suffices to understand the behavior of $A_t(s)$ for $s \in [t_1 - c, t_1 + c] \cup [t_3 - c, t_3 + c]$. We do this for one of the components since the behavior for the other component is entirely similar. We look at the component of $C_t \cap \mathcal{O}$ that lies in the lower half-plane in Figure 3).

Here is a geometric approach. As explained before the difference $A_t(s) - A_t(t_1 - c)$ computes the angular variation of the unit tangent over the interval $[t_1 - c, s]$. A close look at Figure 3 shows that the absolute value of this is bounded above by θ_+ . This proves the boundedness part of the bounded convergence. The almost everywhere convergence is also obvious in view of the above geometric interpretation. The limit function is a bounded function $A_0 : [0, L_0] \rightarrow \mathbb{R}$ that has jumps $-\theta_+$ at t_1 and t_3

$$A_0(t_1^+) - A_0(t_1^-) = A_0(t_3^+) - A_0(t_3^-) = -\theta_+,$$

while the continuous function

$$A_0(t) + \theta_+ H(t - t_1) + \theta_+ H(t - t_3)$$

is differentiable everywhere on $[0, L_0] \setminus \{t_1, t_3\}$ and the derivative is the mean curvature function h_0 of $C_0 \setminus \{p_0\}$.

We can now invoke Theorem 1.8 to conclude that the operators \mathcal{D}_t converge as $t \rightarrow 0^+$ to the operator

$$\mathcal{D}_0^+ : \text{Dom}(\mathcal{D}_0^+) \subset L^2(0, L_0; \mathbb{C}) \rightarrow L^2(0, L_0; \mathbb{C}),$$

where $\text{Dom}(\mathcal{D}_0^+)$ consists of functions $u \in L^2(0, L_0; \mathbb{C})$ such that

$$\begin{aligned} u|_{(t_{j-1}, t_j)} &\in L^{1,2}(t_{j-1}, t_j), \quad \forall j = 1, \dots, 4, \\ u(t_i^+) &= e^{i\theta_+/2} u(t_i^-), \quad i = 1, 3, \\ u(t_2^-) &= u(t_2^+), \quad u(t_4^-) = u(t_4^+), \end{aligned}$$

while for $u \in \text{Dom}(\mathcal{D}_0^+)$ we have

$$\left(\mathcal{D}_0^+ u \right)|_{(t_{j-1}, t_j)} = \left(-i \frac{d}{ds} + \frac{1}{2} h_0(s) \right) u|_{(t_{j-1}, t_j)}, \quad \forall j = 1, \dots, 4.$$

Using the point of view elaborated in Remark 1.9 we let I denote the disjoint union of the intervals I_j , $j = 1, \dots, 4$. We regard \mathcal{D}_0^+ as a closed densely defined operator on the Hilbert space $L^2(I, \mathbb{C})$ with domain consisting of quadruples $\mathbf{u} = (u_1, \dots, u_4) \in L^{1,2}(I)$ satisfying the boundary condition

$$\partial_- \mathbf{u} = \mathbf{T}_+ \partial_+ \mathbf{u},$$

where ∂_{\pm} denotes the restriction to the outgoing/incoming boundary component of I , while

$$\mathbf{T}_+ : \mathbb{C}^4 \cong L^2(\partial_+ I) \rightarrow L^2(\partial_+ I) \cong \mathbb{C}^4,$$

is the transmission operator given by the unitary 4×4 matrix

$$\mathbf{T}_+ = \begin{bmatrix} 0 & 0 & 0 & 1 \\ e^{i\theta_+/2} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & e^{i\theta_+/2} & 0 \end{bmatrix}, \quad \text{and} \quad \mathcal{D}_0^+ \begin{bmatrix} u_1 \\ \vdots \\ u_4 \end{bmatrix} = \left(-i \frac{d}{ds} + \frac{1}{2} h_0 \right) \begin{bmatrix} u_1 \\ \vdots \\ u_4 \end{bmatrix}.$$

Using (1.10) we deduce that

$$\xi^+ = \xi_{\mathcal{D}_0^+} = \frac{1}{2}(1 - 2\rho_+), \quad \rho_+ = \omega_0^+ - [\omega_0^+] = \omega_0 - \frac{\theta_+}{2\pi} - \left[\omega_0 - \frac{\theta_+}{2\pi} \right]. \quad (3.7)$$

B. $t < 0$. We observe that $\mathcal{S}_t = \mathcal{S}_t^a \oplus \mathcal{S}_t^b$, where for $\bullet = a, b$ we have

$$\mathcal{S}_t^\bullet : \text{Dom}(\mathcal{S}_t^\bullet) \subset L^2(0, L_0^\bullet; \mathbb{C}) \rightarrow L^2(0, L_0^\bullet; \mathbb{C}),$$

$$\text{Dom}(\mathcal{S}_t^\bullet) = \{u \in L^{1,2}(0, L_0^\bullet; \mathbb{C}); u(L_0^\bullet) = u(0)\}, \quad \mathcal{S}_t^\bullet u = -i\lambda_t^\bullet \frac{d}{ds} + \frac{1}{2}h_t(s/\lambda_t^\bullet),$$

and λ_t^\bullet is the rescaling factor $\frac{L_0^\bullet}{L_t^\bullet}$. It is convenient to regard \mathcal{S}_t^\bullet as defined on the component C_0^\bullet of C_0^* .

Observe that $C_0^a \setminus \{\bar{a}_0\} = I_1 \cup I_4$ and $C_0^b \setminus \{\bar{b}_0\} = I_2 \cup I_3$. Arguing as in the case $t > 0$ we conclude that

$$\lim_{t \nearrow 0} \omega_t^a = \omega_0^a + \frac{\theta_-}{4\pi}, \quad \lim_{t \nearrow 0} \omega_t^b = \omega_0^b + \frac{\theta_-}{4\pi}, \quad \omega_0^- := \lim_{t \nearrow 0} \omega_t = \omega_0 + \frac{\theta_-}{2\pi}, \quad (3.8)$$

and that the operators \mathcal{D}_t^a and \mathcal{D}_t^b converge in the gap topology as $t \rightarrow 0^-$ to operators

$$\mathcal{D}_0^a : \text{Dom}(\mathcal{D}_0^a) \subset L^2(I_1) \oplus L^2(I_4) \rightarrow L^2(I_1) \oplus L^2(I_4),$$

$$\mathcal{D}_0^b : \text{Dom}(\mathcal{D}_0^b) \subset L^2(I_2) \oplus L^2(I_3) \rightarrow L^2(I_2) \oplus L^2(I_3),$$

where $\text{Dom}(\mathcal{D}_0^a)$ consists of functions $(u_1, u_4) \in L^{1,2}(I_1) \oplus L^{1,2}(I_4)$ such that

$$u_4(\partial_- I_4) = e^{-i\theta_-/2} u_1(\partial_+ I_1), \quad u_4(\partial_+ I_4) = u_1(\partial_- I_1),$$

$\text{Dom}(\mathcal{D}_0^b)$ consists of functions $(u_2, u_3) \in L^{1,2}(I_3) \oplus L^{1,2}(I_3)$ such that

$$u_2(\partial_- I_2) = e^{-i\theta_-/4\pi} u_3(\partial_+ I_3), \quad u_2(\partial_+ I_2) = u_1(\partial_- I_3),$$

where θ_- is depicted in Figure 3, and

$$\mathcal{D}_0^a(u_1, u_4) = \left(-i \frac{du_1}{ds} + \frac{1}{2}h_0 u_1, -i \frac{du_4}{ds} + \frac{1}{2}h_0 u_4 \right),$$

$$\mathcal{D}_0^b(u_2, u_3) = \left(-i \frac{du_2}{ds} + \frac{1}{2}h_0 u_2, -i \frac{du_3}{ds} + \frac{1}{2}h_0 u_3 \right).$$

The direct sum $\mathcal{D}_0^- = \mathcal{D}_0^a \oplus \mathcal{D}_0^b$ is the closed densely defined linear operator on $L^2(I)$ with domain of quadruples $\mathbf{u} = (u_1, \dots, u_4) \in L^{1,2}(I, \mathbb{C})$ satisfying the boundary condition

$$\partial_- \mathbf{u} = \mathbf{T}_- \partial_+ \mathbf{u},$$

where

$$\mathbf{T}_- : \mathbb{C}^4 \cong L^2(\partial_+ I) \rightarrow L^2(\partial_+ I) \cong \mathbb{C}^4,$$

is the transmission operator given by the unitary 4×4 matrix

$$\mathbf{T}_- = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & e^{-i\theta_-/2} & 0 \\ 0 & 1 & 0 & 0 \\ e^{-i\theta_-/2} & 0 & 0 & 0 \end{bmatrix}, \quad \text{and} \quad \mathcal{D}_0^- \begin{bmatrix} u_1 \\ \vdots \\ u_4 \end{bmatrix} = \left(-i \frac{d}{ds} + \frac{1}{2}h_0 \right) \begin{bmatrix} u_1 \\ \vdots \\ u_4 \end{bmatrix}.$$

Then

$$\xi_- = \xi_-^a + \xi_-^b,$$

where for $\bullet = a, b$ we have

$$\xi_-^\bullet = \frac{1}{2}(1 - 2\rho_-^\bullet), \quad \rho_-^\bullet = \omega_0^\bullet + \frac{\theta_-}{4\pi} - \left[\omega_0^\bullet + \frac{\theta_-}{4\pi} \right]. \quad (3.9)$$

Combining (3.6) and (3.8) with the equality $\theta_+ + \theta_- = \pi$ we deduce

$$\omega_0^+ - \omega_0^- = \lim_{t \searrow 0} \omega_t - \lim_{t \nearrow 0} \omega_t = -\frac{1}{2}. \quad (3.10)$$

To prove (3.5) we use the index formula (2.8). We have

$$\begin{aligned}
 i_{APS}(\bar{\partial}) &= -\frac{1}{2} - \omega_1 + \omega_{-1} - \xi_{D_1} + \xi_{D_{-1}}. \\
 &\stackrel{(3.10)}{=} \omega_0^+ - \omega_0^- - \omega_1 + \omega_{-1} - \xi_{D_1} + \xi_{D_{-1}} \\
 &= (\omega_0^+ + \xi^+) - (\omega_1 + \xi_{D_1}) - (\omega_0^- + \xi^-) + (\omega_{-1} + \xi_{D_{-1}}) - (\xi^+ - \xi^-) \\
 &\stackrel{(1.12)}{=} -\lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t; \varepsilon < t \leq 1) - \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t, -1 \leq t < -\varepsilon) - (\xi_+ - \xi_-).
 \end{aligned}$$

□

Remark 3.3 (Twisted Dolbeault operators). (a) Here the outline of an analytic argument proving (A₂). Using (3.1) we deduce that this component has a parametrization compatible with the orientation given by

$$y_t = -(\zeta_t + mx^2)^{1/2}, \quad |x| < d_t \quad (3.11)$$

where $\zeta_t = \frac{t}{\beta}$, $m = \frac{\alpha}{\beta}$ and d_t is such that the length of this arc is $2c/\lambda_t$. Observe that there exists $d_* > 0$ such that $\lim_{t \rightarrow 0^+} dt = d_*$. We have

$$dy_t = -mx(\zeta_t + mx^2)^{-1/2} dx.$$

Set

$$\begin{aligned}
 y_t' &:= \frac{dy_t}{dx} = -mx(\zeta_t + mx^2)^{-1/2}, \\
 y_t'' &:= \frac{d^2y_t}{dx^2} = -m(\zeta_t + mx^2)^{-1/2} + m^2x^2(\zeta_t + mx^2)^{-3/2} = -\frac{m\zeta_t}{(\zeta_t + mx^2)^{3/2}}.
 \end{aligned}$$

The arclength is

$$d\sigma^2 = \left(1 + (y_t')^2\right) dx^2 = \underbrace{\left(1 + \frac{m^2x^2}{\zeta_t + mx^2}\right)}_{=:w(t,x)^2} dx^2.$$

The mean curvature h_t is found using the Frenet formulæ. More precisely $h_t(x) = \frac{y_t''}{w^3}$. Then

$$h_t d\sigma = h_t w dx = \frac{y_t''}{1 + (y_t')^2} dx = -\frac{m\zeta_t dx}{(\zeta_t + mx^2)^{1/2}(\zeta_t + mx^2 + m^2x^2)}.$$

We observe now that we can write $h_t d\sigma = \phi_t^*(\rho_\infty du)$, where ϕ_t is the rescaling map

$$x \mapsto u = t^{-1/2}x \quad \text{and} \quad \rho_\infty(u) = -\frac{m\zeta_1}{(\zeta_1 + mu^2)^{1/2}(\zeta_1 + mu^2 + m^2u^2)}.$$

This then allows us to conclude via a standard argument that the densities $h_t d\sigma$ converge very weakly as $t \rightarrow 0^+$ to a δ -measure concentrated at the origin.

(b) The results in Theorem 3.2 extend without difficulty to Dolbeault operators twisted by line bundles. More precisely, given a Hermitian line bundle L and a hermitian connection A on L , we can form a Dolbeault operator $\bar{\partial}_A : C^\infty(L) \rightarrow C^\infty(L \otimes K_\Sigma^{-1})$. Fortunately, all the line bundles on a the two-dimensional cobordism Σ are trivializable. We fix a trivialization so that the connection A can be identified with a purely imaginary 1-form

$$A = ia, \quad a \in \Omega^1(\Sigma).$$

Then

$$\bar{\partial}_A = \bar{\partial} + ia^{0,1}.$$

The restriction of $D_A^+ = \sqrt{2}\bar{\partial}_A$ to the cooriented curve C_t is

$$D_A(t) = -i\nabla_s^A + \frac{1}{2}h_t = -i\frac{d}{ds} + \frac{1}{2}h_t + a_t, \quad a_t := a\left(\frac{d}{ds}\right) \in \Omega^0(C_t).$$

As in the proof of Theorem 3.2, we only need to understand the behavior of a_t in the neighborhood $\mathcal{O} \cap C_t$. Suppose for simplicity $t > 0$ and we concentrate only on the component of $C_t \cap \mathcal{O}$ that lies in the lower half-plane of Figure 3. In the neighborhood \mathcal{O} we can write

$$a = pdx + qdy, \quad p, q \in C^\infty(\mathcal{O}).$$

Using the parametrization (3.11) we deduce that

$$a|_{C_t \cap \mathcal{O}} = \left(p - mqx(\zeta_t + mx^2)^{-1/2} \right) dx = a_t ds = a_t w dx$$

Hence, as $t \rightarrow 0^+$, the measure $a_t ds$ converges to the measure $(p - m^{1/2}(2H(x) - 1)) dx$

(c) One may ask what happens in the case of a cobordism corresponding to a local min/max of a Morse function. In this case Σ is a disk, the regular level sets C_t are circles and the singular level set is a point. Consider for example the case of a local minimum. Assume that the metric near the minimum p_0 is Euclidean, and in some Euclidean coordinates near p_0 we have $f = x^2 + y^2$. Then C_t is the Euclidean circle of radius $t^{1/2}$, and the function h_t is the constant function $h_t = t^{-1/2}$. Then $\omega_t = \frac{1}{2}$, $\xi_t = \frac{1}{2}$ and the Atiyah-Patodi-Singer index of $\bar{\partial}$ on the Euclidean disk of radius $t^{1/2}$ is 0. The operator D_t can be identified with the operator

$$-i\frac{d}{ds} + \frac{1}{2t^{1/2}}$$

with periodic boundary conditions on the interval $[0, 2\pi t^{1/2}]$. Using the rescaling trick in Remark 1.6 we see that this operator is conjugate to the operator $L_t = -t^{1/2}i\frac{d}{ds} + \frac{1}{2}$ on the interval $[0, 2\pi]$ with periodic boundary conditions. The switched graphs of these operators

$$\tilde{\Gamma}_{L_t} = \{(L_t u, u); u \in L^{1,2}(0, 2\pi; \mathbb{C}); u(0) = u(2\pi)\} \subset \mathbf{H} \oplus \mathbf{H}, \quad \mathbf{H} = L^2(0, 2\pi; \mathbb{C}),$$

converge in the gap topology to the subspace $\mathbf{H}_+ = \mathbf{H} \oplus 0 \subset \mathbf{H} \oplus \mathbf{H}$. This limit is not the switched graph of any operator. However, this limiting space forms a Fredholm pair with $\mathbf{H}_- = 0 \oplus \mathbf{H}$ and invoking the results in [5] we conclude that the limit

$$\lim_{\varepsilon \searrow 0} SF(L_t; \varepsilon \leq t \leq t_0)$$

exists and it is finite. □

4. The Kashiwara-Wall index

In this final section we would like to identify the correction term in the right hand side of (3.5) with a symplectic invariant that often appears in surgery formulæ. To this aim, we need to elaborate on the symplectic point of view first outlined in Remark 1.9.

Fix a finite dimensional complex hermitian space \mathbf{E} , let $n := \dim \mathbf{E}$, and set

$$\widehat{\mathbf{E}} := \mathbf{E} \oplus \mathbf{E}, \quad \mathbf{E}_+ := \mathbf{E} \oplus 0, \quad \mathbf{E}_- := 0 \oplus \mathbf{E},$$

and let $J : \widehat{\mathbf{E}} \rightarrow \widehat{\mathbf{E}}$ be the unitary operator given by the block decomposition

$$J = \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}.$$

We let Lag denote the space of hermitian lagrangians on $\widehat{\mathbf{E}}$, i.e., complex subspaces $L \subset \widehat{\mathbf{E}}$ such that $L^\perp = JL$. As explained in [5, 14] any such a lagrangian can be identified with the graph⁷ of a complex isometry $T : \mathbf{E}_+ \rightarrow \mathbf{E}_-$, or equivalently, with the group $U(\mathbf{E})$ of unitary operators on \mathbf{E} . In other words, the graph map

$$\Gamma : U(\mathbf{E}) \rightarrow \text{Lag}(\widehat{\mathbf{E}}), \quad U(\mathbf{E}) \mapsto \Gamma_T \subset \widehat{\mathbf{E}}$$

is a diffeomorphism. The involution $L \leftrightarrow JL$ on Lag corresponds via this diffeomorphism to the involution $T \leftrightarrow -T$ on $U(\mathbf{E})$.

We define a branch of the logarithm $\log : \mathbb{C}^* \rightarrow \mathbb{C}$ by requiring $\text{Im } \log \in [-\pi, \pi)$. Equivalently,

$$\log z = \int_{\gamma_z} \frac{d\zeta}{\zeta},$$

where $\gamma_z : [0, 1] \rightarrow \mathbb{C}$ is any smooth path from 1 to z such that

$$\forall t \in [0, 1], \quad \gamma_z(t) \notin (-\infty, 0].$$

In particular, $\log(-1) = \pi i$. Following [11, §6] we define

$$\begin{aligned} \tau : U(\mathbf{E}) \times U(\mathbf{E}) &\rightarrow \mathbb{R}, \quad \tau(T_0, T_1) = \frac{1}{2\pi i} \text{tr } \log(T_1^{-1}T_0) \\ &= \frac{1}{2\pi i} \sum_{\lambda \in \mathbb{C}^*} (\log \lambda) m_\lambda, \quad m_\lambda := \dim \ker(\lambda - T_1^{-1}T_0). \end{aligned}$$

Observe that

$$e^{2\pi i \tau(T_0, T_1)} = \frac{\det T_0}{\det T_1} \tag{4.1}$$

Note that

$$\tau(T_0, T_1) + \tau(T_1, T_0) = \dim \ker(T_0 + T_1). \tag{4.2}$$

Via the graph diffeomorphism we obtain a map

$$\mu = \tau \circ \Gamma : \text{Lag} \times \text{Lag} \rightarrow \mathbb{R}.$$

The equality (4.2) can be rewritten as

$$\tau(L_0, L_1) + \tau(L_1, L_0) = \dim(L_0 \cap JL_1) = \dim(JL_0 \cap L_1). \tag{4.3}$$

We want to relate the invariant τ to the eta invariant of a natural selfadjoint operator. We associate to each pair $L_0, L_1 \in \text{Lag}$ the selfadjoint operator

$$D_{L_0, L_1} : V(L_0, L_1) \subset L^2(I, \widehat{\mathbf{E}}) \rightarrow L^2(I, \widehat{\mathbf{E}}),$$

where

$$V(L_0, L_1) = \left\{ u \in L^{1,2}(I, \widehat{\mathbf{E}}); \quad u(0) \in L_0, \quad u(1) \in L_1 \right\}, \quad D_{L_0, L_1} u = J \frac{du}{dt}.$$

This is a selfadjoint operator with compact resolvent. We want to describe its spectrum, and in particular, prove that it has a well defined eta invariant. Let $T_0, T_1 : \mathbf{E}_+ \rightarrow \mathbf{E}_-$ denote the isometries associated to L_0 and respectively L_1 . Then $T_1^{-1}T_0$ is a unitary operator on \mathbf{E}_+ so its spectrum consists of complex numbers of norm 1.

⁷In [11] a lagrangian is identified with the graph of an isometry $\mathbf{E}_- \rightarrow \mathbf{E}_+$ which explains why our formulæ will look a bit different than the ones on [11]. Our choice is based on the conventions in [5] which seem to minimize the number of signs in the Schubert calculus on Lag .

Proposition 4.1. *For any $L_0, L_1 \in \text{Lag}$ we have*

$$\text{spec } D_{L_0, L_1} = \frac{1}{2i} \exp^{-1} \left(\text{spec}(T_1^{-1}T_0) \right). \quad (4.4)$$

In particular, the spectrum of D_{L_0, L_1} consists of finitely many arithmetic progressions with ratio π so that the eta invariant of D_{L_0, L_1} is well defined.

Proof. Observe first that any $u \in L^2(I, \widehat{\mathbf{E}})$ decomposes as a pair

$$u = (u_+, u_-), \quad u_{\pm} \in L^2(I, \mathbf{E}_{\pm}).$$

If $u \in V(L_0, L_1)$ is an eigenvector of D_{L_0, L_1} corresponding to an eigenvalue λ then u satisfies the boundary value problems

$$-i \frac{du_+}{dt} = \lambda u_+, \quad i \frac{du_-}{dt} = \lambda u_-, \quad (4.5a)$$

$$u_-(0) = T_0 u_+(0), \quad u_-(1) = T_1 u_+(1). \quad (4.5b)$$

The equalities (4.5a) imply that

$$u_+(1) = e^{i\lambda} u_+(0), \quad u_-(1) = e^{-i\lambda} u_-(0).$$

Using (4.5b) we deduce

$$e^{i\lambda} T_1 u_+(0) = u_-(1) = e^{-i\lambda} u_-(0) = e^{-i\lambda} T_0 u_+(0).$$

Hence

$$e^{2i\lambda} \in \text{spec}(T_1^{-1}T_0) \implies \lambda \in \frac{1}{2i} \exp^{-1} \left(\text{spec}(T_1^{-1}T_0) \right).$$

Running the above argument in reverse we deduce that any $\lambda \in \frac{1}{2i} \exp^{-1} \left(\text{spec}(T_1^{-1}T_0) \right)$ is an eigenvalue of D_{L_0, L_1} . \square

We let $\xi(L_0, L_1)$ denote the reduced eta invariant of D_{L_0, L_1} ,

$$\xi(L_0, L_1) = \frac{1}{2} \left(\dim \ker D_{L_0, L_1} + \eta_{D_{L_0, L_1}}(0) \right).$$

If $e^{i\theta_1}, \dots, e^{i\theta_n}$, $\theta_1, \dots, \theta_n \in (0, 2\pi]$, are the eigenvalues of $T_1^{-1}T_0$, then the spectrum of D_{L_0, L_1} is

$$\text{spec}(D(L_0, L_1)) = \bigcup_{k=1}^n \left\{ \frac{\theta_k}{2} + \pi\mathbb{Z} \right\}.$$

and we deduce as in Section 1 using (1.8) that

$$\eta_{D_{L_0, L_1}} = \sum_{\theta_k \in (0, 2\pi)} \left(1 - \frac{\theta_k}{\pi} \right),$$

and

$$\xi(L_0, L_1) = \frac{1}{2} \sum_{\theta_k \in (0, 2\pi)} \left(1 - \frac{\theta_k}{\pi} \right) + \frac{1}{2} \dim \ker D_{L_0, L_1}.$$

On the other hand

$$\frac{1}{2\pi i} \text{tr} \log(-T_1^{-1}T_0) = \frac{1}{2\pi} \sum_{\theta_k \in (0, 2\pi]} (\theta_k - \pi) = -\frac{1}{2} \sum_{\theta_k \in (0, 2\pi)} \left(1 - \frac{\theta_k}{\pi} \right) + \frac{1}{2} \dim \ker(T_0 - T_1).$$

Since $\ker D_{L_0, L_1} \cong \ker(T_0 - T_1) \cong L_0 \cap L_1$ we conclude

$$\tau(T_0, -T_1) = \tau(-T_0, T_1) = \tau(JL_0, L_1) = -\xi(L_0, L_1) + \dim(L_0 \cap L_1).$$

Using (4.2) we deduce

$$\xi(L_0, L_1) = \tau(T_1, -T_0) = \tau(L_1, JL_0) = \tau(JL_1, L_0). \quad (4.6)$$

Following [11] (see also [4]) we associate to each triplet of lagrangians L_0, L_1, L_2 the quantity

$$\omega(L_0, L_1, L_2) := \tau(L_1, L_0) + \tau(L_2, L_1) + \tau(L_0, L_2),$$

and we will refer to its as the (*hermitian*) *Kashiwara-Wall index* (or simply the *index*) of the triplet. Observe that ω is indeed an integer since (4.1) implies that

$$e^{2\pi i \omega(L_0, L_1, L_2)} = 1.$$

We set

$$d(L_0, L_1, L_2) := \dim(JL_0 \cap L_1) + \dim(JL_1 \cap L_2) + \dim(JL_2 \cap L_0).$$

Using (4.3) we deduce that for any permutation φ of $\{0, 1, 2\}$ with signature $\epsilon(\varphi) \in \{\pm 1\}$ we have

$$\omega(L_0, L_1, L_2) - \epsilon(\varphi)\omega(L_{\varphi(0)}, L_{\varphi(1)}, L_{\varphi(2)}) = d(L_0, L_1, L_2) \times \begin{cases} 0, & \varphi \text{ even} \\ 1, & \varphi \text{ odd.} \end{cases} \quad (4.7)$$

We want to apply the above facts to a special choice of $\widehat{\mathbf{E}}$. Let \mathbf{I} denote the disjoint union of the intervals I_1, \dots, I_4 introduced in Section 3. They were obtained by removing the points \bar{a}_0, p_0 and \bar{b}_0 from the critical level set C_0 ; Figure 2. We interpret \mathbf{I} as an oriented 1-dimensional with boundary and we let

$$\widehat{\mathbf{E}} := L^2(\partial \mathbf{I}), \quad \mathbf{E}_{\pm} = L^2(\partial_{\pm} \mathbf{I}).$$

The spaces \mathbf{E}_{\pm} have canonical bases and thus we can identify both of them with the standard Hermitian space $\mathbf{E} = \mathbb{C}^4$. Define $J : \widehat{\mathbf{E}} \rightarrow \widehat{\mathbf{E}}$ as before. We have a canonical differential operator

$$D_0 : C^{\infty}(\mathbf{I}, \mathbb{C}) \rightarrow C^{\infty}(\mathbf{I}, \mathbb{C}), \quad D_0 \begin{bmatrix} u_1 \\ \vdots \\ u_4 \end{bmatrix} = \begin{bmatrix} -i \frac{du_1}{dt} + \frac{1}{2} h_0|_{I_1} \\ \vdots \\ \vdots \\ -i \frac{du_4}{dt} + \frac{1}{2} h_0|_{I_4} \end{bmatrix},$$

We set

$$\omega_k := \frac{1}{4\pi} \int_{I_k} h_0 ds$$

so that

$$\omega_0 = \omega_1 + \dots + \omega_4, \quad \omega_0^a = \omega_1 + \omega_4, \quad \omega_0^b = \omega_2 + \omega_3.$$

We have a natural restriction map

$$r : C^{\infty}(\mathbf{I}, \mathbb{C}) \rightarrow L^2(\partial \mathbf{I}, \mathbb{C}) = \widehat{\mathbf{E}}$$

and we define the *Cauchy data space* of D_0 to be the subspace

$$\Lambda_0 := r(\ker D_0) \subset \widehat{\mathbf{E}}.$$

We can verify easily that Λ_0 is a Lagrangian subspace of $\widehat{\mathbf{E}}$ that is described by the isometry $\mathbf{T}_0 : \mathbf{E}_+ \rightarrow \mathbf{E}_-$ given by the diagonal matrix

$$\mathbf{T}_0 = \text{Diag}(e^{2\pi i \omega_1}, \dots, e^{2\pi i \omega_4}).$$

⁸In the remainder of this section we assume⁸ that the operators \mathcal{D}_0^{\pm} that appear in Theorem 3.2 are invertible.

⁸This assumption is satisfied for a generic choice of metric on Σ .

Proposition 4.2. *Let \mathcal{D}_0^\pm be the operators that appear in Theorem 3.2. Then*

$$\xi_{\mathcal{D}_0^\pm} = -\xi(\Gamma_{\mathcal{T}_\pm}, \Lambda_0) = \xi(\Lambda_0, \Gamma_{\mathcal{T}_\pm}) = \tau(\Gamma_{\mathcal{T}_\pm}, J\Lambda_0) \quad (4.8)$$

Proof. We need to find the spectra of $\mathbf{T}_0^{-1}\mathbf{T}_\pm$. We set $z_k = e^{-2\pi i\omega_k}$, $k = 1, \dots, 4$ and $\rho = e^{-i\theta_+/2}$, so that $e^{i\theta_-/2} = -i\rho$. Then

$$\mathbf{T}_0^*\mathbf{T}_+ = \begin{bmatrix} 0 & 0 & 0 & z_1 \\ z_2\rho & 0 & 0 & 0 \\ 0 & z_3 & 0 & 0 \\ 0 & 0 & z_4\rho & 0 \end{bmatrix}, \quad \mathbf{T}_0^*\mathbf{T}_- = \begin{bmatrix} 0 & 0 & 0 & z_1 \\ 0 & 0 & -iz_2\rho & 0 \\ 0 & z_3 & 0 & 0 \\ -iz_4\rho & 0 & 0 & 0 \end{bmatrix}.$$

The eigenvalues of $\mathbf{T}_0^*\mathbf{T}_+$ are the fourth order roots of

$$\zeta = \rho^2 z_1 \cdots z_4 = e^{i(\theta_+ - 2\pi\omega_0)}.$$

Hence

$$\exp^{-1}(\text{spec}(\mathbf{T}_0^*\mathbf{T}_+)) = \frac{i(\theta_+ - 2\pi\omega_0)}{4} + \frac{\pi i}{2}\mathbb{Z}$$

and using (4.4) we deduce

$$\text{spec}(D_{\Gamma_{\mathcal{T}_+}, \Lambda_0}) = \frac{\pi}{2} \left\{ \left(\frac{\theta_+}{2\pi} - \omega_0 \right) + \mathbb{Z} \right\}.$$

The eigenvalues of $\mathbf{T}_0^*\mathbf{T}_-$ are the square roots of

$$z_1 z_4 e^{i\theta_-/2} = e^{-i(\theta_-/2 + 2\pi\omega_0^a)} \quad \text{and} \quad z_1 z_4 e^{i\theta_-/2} = e^{-i(\theta_-/2 + 2\pi\omega_0^b)}.$$

Hence

$$\text{spec}(D_{\Gamma_{\mathcal{T}_+}, \Lambda_0}) = \left\{ -\pi \left(\frac{\theta_-}{4\pi} + \omega_0^a \right) + \pi\mathbb{Z} \right\} \cup \left\{ -\pi \left(\frac{\theta_-}{4\pi} + \omega_0^b \right) + \pi\mathbb{Z} \right\}.$$

The desired conclusion follows using (3.7), (3.9) and (1.8). \square

Theorem 4.3. *Under the same assumptions and notations as in Theorem 3.2 we have*

$$i_{APS}(\bar{\partial}) + \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t; \varepsilon < t \leq 1) + \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t, -1 \leq t < -\varepsilon) = -\omega(J\Lambda_0, \Gamma_{\mathcal{T}_+}, \Gamma_{\mathcal{T}_-}).$$

Proof. We have

$$i_{APS}(\bar{\partial}) + \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t; \varepsilon < t \leq 1) + \lim_{\varepsilon \rightarrow 0^+} SF(\mathcal{D}_t, -1 \leq t < -\varepsilon) \stackrel{(3.5)}{=} -(\xi_+ - \xi_-)$$

$$\stackrel{(4.8)}{=} -\tau(\Gamma_{\mathcal{T}_+}, \Lambda_0) - \tau(J\Lambda_0, \Gamma_{\mathcal{T}_-}) = -\omega(J\Lambda_0, \Gamma_{\mathcal{T}_+}, \Gamma_{\mathcal{T}_-}) + \tau(\Gamma_{\mathcal{T}_+}, \Gamma_{\mathcal{T}_-}).$$

To compute $\tau(\Gamma_{\mathcal{T}_+}, \Gamma_{\mathcal{T}_-}) = \tau(\mathbf{T}_+, \mathbf{T}_-)$ we need to compute the spectrum of $\mathbf{T}_-^*\mathbf{T}_+$. We set $\rho = e^{i\theta_+/2}$ so that $e^{-i\theta_-/2} = -i\rho$. We have

$$\mathbf{T}_-^*\mathbf{T}_+ = \begin{bmatrix} 0 & 0 & 0 & i\bar{\rho} \\ 0 & 0 & 1 & 0 \\ 0 & i\bar{\rho} & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 & 0 & 0 & 1 \\ \rho & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \rho & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & i & 0 \\ 0 & 1 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

From the second and fourth column we see that 1 is an eigenvalue of $T_-^* T_+$ with multiplicity 2. The other two eigenvalues are $\pm i$, namely the eigenvalues of the 2×2 minor

$$\begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}.$$

This shows that $\tau(T_+, T_-) = 0$. □

Let I denote the unit interval $[0, 1]$. We set $\partial_+ I = \{1\}$, $\partial_- I = \{0\}$. We identify \widehat{E} with the finite dimensional Hilbert space $L^2(\partial I, E)$ and the Hilbert spaces E_{\pm} with

References

- [1] V.I. Arnold: *The complex lagrangian grassmannian*, *Funct. Anal. and its Appl.*, **34**(2000), 208-210.
- [2] M.F. Atiyah, V.K. Patodi, I.M. Singer: *Spectral asymmetry and Riemannian geometry. III*, *Math. Proc. Camb. Phil. Soc.* **79**(1976), 71-99.
- [3] N. Berline, E. Getzler, M. Vergne: *Heat Kernels and Dirac Operators*, Springer Verlag, 1992.
- [4] S. Cappell, R. Lee, E.Y. Miller: *On the Maslov index*, *Comm. Pure Appl. Math.* **47**(1994), 121-186.
- [5] D.F. Cibotaru: *Localization formulæ in odd K-theory*, [arXiv: 0901.2563](https://arxiv.org/abs/0901.2563).
- [6] R. E. Edwards: *Functional analysis*, Dover Publications, 1995.
- [7] G.B. Folland: *Real Analysis. Modern Techniques and Their Applications*, 2nd Edition, John Wiley & Sons, 1999.
- [8] P.B. Gilkey: *On the index of geometrical operators for Riemannian manifolds with boundary*, *Adv. Math.*, **102**(1993), 129-183.
- [9] G. Grubb: *Heat operator trace expansions and index for general Atiyah-Patodi-Singer boundary problems*, *Comm. Partial Differential Equations*, **17**(1992), no.11-12, 2031-2077.
- [10] T. Kato: *Perturbation Theory for Linear Operators*, Springer Verlag, 1995.
- [11] P. Kirk, M. Lesch: *The η -invariant, Maslov index, and spectral flow for Dirac-type operators on manifolds with boundary*, *Forum Math.*, **16**(2004), 553-629.
- [12] P. Kronheimer, T. Mrowka: *Monopoles and Three-Manifolds*, Cambridge University Press, 2007.
- [13] L.I. Nicolaescu: *Lectures on the Geometry of Manifolds*, 2nd Edition World Sci. Pub. Co. 2009.
- [14] _____: *Schubert calculus on the grassmannian of hermitian lagrangian spaces*, [arXiv: 0708.2669](https://arxiv.org/abs/0708.2669).
- [15] J. Oprea: *Differential Geometry and Its Applications*, Prentice Hall, 2007.
- [16] R. Seeely, I.M. Singer: *Extending $\bar{\partial}$ to singular Riemann surfaces*, *J. Geom. Physics*, **5**(1988), 121-136.
- [17] E.T. Whittaker, G.N. Watson: *A Course of Modern Analysis*, Cambridge University Press, Fourth Edition, 1927.
- [18] K. Yosida: *Functional Analysis*, Springer Verlag, 1980.
- [19] C.T.C Wall: *Non-additivity of the signature*, *Invent. Math.*, **7**(1969), 269-274.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556-4618.

E-mail address: cibotaru.1@nd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556-4618.

E-mail address: nicolaescu.1@nd.edu