

Duality between Harmonic and Bergman spaces

Mei-Chi Shaw*

ABSTRACT. In this paper we study the duality of the harmonic spaces on the annulus $\Omega = \Omega_1 \setminus \overline{\Omega^-}$ between two pseudoconvex domains with $\Omega^- \subset\subset \Omega_1$ in \mathbb{C}^n and the Bergman spaces on Ω^- . We show that on the annulus Ω , the space of harmonic forms for the critical case on $(0, n-1)$ -forms is infinite dimensional and it is dual to the Bergman space on the pseudoconvex domain Ω^- . The duality is further identified explicitly by the Bochner-Martinelli transform, generalizing a result of Hörmander.

Introduction

Let Ω^- and Ω_1 be two bounded pseudoconvex domains in \mathbb{C}^n with $\Omega^- \subset\subset \Omega_1$. In this paper we study the duality of the harmonic spaces on the annulus $\Omega = \Omega_1 \setminus \overline{\Omega^-}$ and the Bergman spaces on Ω^- . This paper is inspired by a recent paper of Hörmander [**Hö 2**] where the null space of the $\bar{\partial}$ -Neumann operator on a spherical shell as well as on an ellipsoid in \mathbb{C}^n has been computed by explicit formula for the critical case for $(0, n-1)$ -forms.

The $\bar{\partial}$ -Neumann problem on the annulus has been studied in [**Sh 1**] on an annulus between two pseudoconvex domains in \mathbb{C}^n or in a hermitian Stein manifold. When the boundary is smooth, the closed range property and boundary regularity for $\bar{\partial}$ were established in the earlier work (see [**BS**] or [**Sh1**]) for $0 < q \leq n-1$ and $n \geq 2$. In the case when $0 < q < n-1$, the space of harmonic forms is trivial. In this paper, we will study the critical case when $q = n-1$ on the annulus Ω . In this case the space of harmonic forms is infinite dimensional. Our goal is to establish the duality between the harmonic forms in the critical degree with the Bergman spaces on the domain Ω^- .

In the first section, we recall the Hodge decomposition theorem on the annulus between two pseudoconvex domains. In the second section we establish the duality between the harmonic forms with coefficients in the Sobolev $W^1(\Omega)$ spaces with the Bergman spaces on Ω^- . We then refine the duality to duality between L^2 spaces in Section 3.

2010 *Mathematics Subject Classification.* Primary 32W05, 35N15, 58J32.

Key words and phrases. Cauchy-Riemann equations, Pseudoconvex domains, Harmonic spaces, Bergman spaces.

*partially supported by NSF grant.

1. L^2 theory for $\bar{\partial}$ on the annulus between two weakly pseudoconvex domains in \mathbb{C}^n

We recall the following L^2 existence and estimates for $\bar{\partial}$ in the annulus between two pseudoconvex domains (see Theorems 3.2 and 3.3 in Shaw [Sh4]).

THEOREM 1.1. *Let $\Omega \subset\subset \mathbb{C}^n$, $n \geq 3$, be the annulus domain $\Omega = \Omega_1 \setminus \bar{\Omega}^-$ between two pseudoconvex domains Ω_1 and Ω^- . We assume that $\Omega^- \subset\subset \Omega_1$ and Ω^- has C^2 boundary. For any $f \in L^2_{(p,q)}(\Omega)$, where $0 \leq p \leq n$ and $0 \leq q < n-1$, such that $\bar{\partial}f = 0$ in Ω , the following hold:*

- (1) *there exists $F \in W_{(p,q)}^{-1}(\Omega_1)$ such that $F|_{\Omega} = f$ and $\bar{\partial}F = 0$ in Ω_1 in the distribution sense.*
- (2) *there exists $u \in L^2_{(p,q-1)}(\Omega)$ satisfying $\bar{\partial}u = f$ in Ω .*

For $q = n-1$, there is an additional compatibility condition for the $\bar{\partial}$ -closed extension of $(p, n-1)$ -forms.

THEOREM 1.2. *Let $\Omega \subset\subset \mathbb{C}^n$ be the annulus domain $\Omega = \Omega_1 \setminus \bar{\Omega}^-$ between two pseudoconvex domains Ω_1 and Ω^- . We assume that $\Omega^- \subset\subset \Omega_1$ and Ω^- has C^2 boundary. For any $\bar{\partial}$ -closed $f \in L^2_{(p,n-1)}(\Omega)$, where $0 \leq p \leq n$, the following hold:*

- (1) *There exists $F \in W_{(p,n-1)}^{-1}(\Omega_1)$ such that $F|_{\Omega} = f$ and $\bar{\partial}F = 0$ in Ω_1 in the distribution sense.*
- (2) *The restriction of f to $b\Omega^-$ satisfies the compatibility condition*

$$(1.1) \quad \int_{b\Omega^-} f \wedge \phi = 0, \quad \phi \in W_{(n-p,0)}^1(\Omega^-) \cap \text{Ker}(\bar{\partial}).$$

- (3) *There exists $u \in L^2_{(p,n-2)}(\Omega)$ satisfying $\bar{\partial}u = f$ in Ω .*

Corollary 1.3. *Let Ω be the same as in Theorem 1.2. Then $\bar{\partial}$ has closed range in $L^2_{(p,n-1)}(\Omega)$ and the $\bar{\partial}$ -Neumann operator $N_{(p,n-1)}$ exists on $L^2_{(p,n-1)}(\Omega)$.*

Theorem 1.4 (Hodge Decomposition Theorem). *Let $\Omega \subset\subset \mathbb{C}^n$ be the annulus domain $\Omega = \Omega_1 \setminus \bar{\Omega}^-$ between two pseudoconvex domains Ω_1 and Ω^- . We assume that $\Omega^- \subset\subset \Omega_1$ and Ω^- has C^2 boundary. Then the $\bar{\partial}$ -Neumann operator $N_{(p,q)}$ exists on $L^2_{(p,q)}(\Omega)$ for $0 \leq p \leq n$ and $0 \leq q \leq n$. For any $f \in L^2_{(p,q)}(\Omega)$, we have*

$$\begin{aligned} f &= \bar{\partial}^* \bar{\partial} N_{(p,0)} f + H_{(p,0)} f, & q &= 0. \\ f &= \bar{\partial} \bar{\partial}^* N_{(p,q)} f + \bar{\partial}^* \bar{\partial} N_{(p,q)} f, & 1 &\leq q \leq n-2. \\ f &= \bar{\partial} \bar{\partial}^* N_{(p,n-1)} f + \bar{\partial}^* \bar{\partial} N_{(p,n-1)} f + H_{(p,n-1)} f, & q &= n-1. \\ f &= \bar{\partial} \bar{\partial}^* N_{(p,n)} f, & q &= n. \end{aligned}$$

We have used the notation $H_{(p,q)}$ to denote the projection operator from $L^2_{(p,q)}(\Omega)$ onto the harmonic space $\mathcal{H}_{(p,q)}(\Omega) = \ker(\square_{(p,q)})$.

For a proof of Theorem 1.4, see Theorem 3.5 in [Sh4].

Remark: All the results can be extended to annulus between pseudoconvex domains in a Stein manifold with trivial modification.

2. The duality between $H_{W^1}^{(p,n-1)}(\Omega)$ and $H^{(n-p,0)}(\Omega^-)$

For $k \geq 0$, we define the Dolbeault cohomology $H_{W^k}^{(p,q)}(\Omega)$ with $W^k(\Omega)$ -coefficients by

$$H_{W^k}^{(p,q)}(\Omega) = \frac{\{f \in W_{(p,q)}^k(\Omega) \mid \bar{\partial}f = 0\}}{\{f \in W_{(p,q)}^k(\Omega) \mid f = \bar{\partial}u, u \in W_{(p,q-1)}^k(\Omega)\}}.$$

For $k \in \mathbb{R}$, we define $H_{W^k}^{(p,0)}(\Omega)$ to be the space of $(p,0)$ -forms with holomorphic coefficients in $W^k(\Omega)$.

If Ω is the annulus between two pseudoconvex domains as in Theorem 1.2, we have that the space $\{f \in W_{(p,q)}^k(\Omega) \mid f = \bar{\partial}u, u \in W_{(p,q-1)}^k(\Omega)\}$ is closed. Furthermore, we have from Theorem 1.4:

$$\mathcal{H}_{(p,n-1)}(\Omega) \simeq H_{L^2}^{(p,n-1)}(\Omega).$$

We will use the notation $H^{(p,n-1)}(\Omega)$ for $H_{L^2}^{(p,n-1)}(\Omega)$ and $H_{W^1}(\Omega) = H_{W^1}^{(0,0)}(\Omega)$.

Theorem 2.1. *Let $\Omega \subset \subset \mathbb{C}^n$ be the annulus domain $\Omega = \Omega_1 \setminus \bar{\Omega}^-$ between two pseudoconvex domains Ω_1 and Ω^- with smooth boundary and $\Omega^- \subset \subset \Omega_1$, $n \geq 2$. For each $k \geq 0$ and $0 \leq p \leq n$, the space $H_{W^k}^{(p,q)}(\Omega) = \{0\}$, when $0 < q < n-1$ and the space $H_{W^k}^{(p,n-1)}(\Omega)$ is of infinite dimension. Furthermore, we have the isomorphism:*

$$H_{W^k}^{(p,n-1)}(\Omega) \simeq (H_{W^{-k+1}}^{(n-p,0)}(\Omega^-))'$$

where the right-hand side is the space of all bounded linear functionals on the space $H_{W^{-k+1}}^{(n-p,0)}(\Omega^-)$.

PROOF. First we assume that $k = 0$. Suppose that $f \in L_{(p,n-1)}^2(\Omega)$ and $\bar{\partial}f = 0$. We define a pairing between $H^{(p,n-1)}(\Omega)$ and $H_{W^1}^{(n-p,0)}(\Omega^-)$

$$l : H^{(p,n-1)}(\Omega) \times H_{W^1}^{(n-p,0)}(\Omega^-) \rightarrow \mathbb{C}$$

by

$$(2.1) \quad l([f], h) = \int_{b\Omega^-} f \wedge h, \quad h \in H_{W^1}^{(n-p,0)}(\Omega^-).$$

First we note that the pairing (2.1) is well-defined. It is well-known any holomorphic function or forms with $L^2(\Omega)$ coefficients has a well-defined trace in $W^{-\frac{1}{2}}(b\Omega)$ (see e.g. [LM]). For any f in $L_{(p,n-1)}^2(\Omega)$ with $\bar{\partial}f = 0$ and $\bar{\partial}^*f = 0$, we also have a well-defined trace in $W^{-\frac{1}{2}}(b\Omega)$ (see [Sh3] for details). Any function or form with $W^1(\Omega^-)$ coefficients has trace in $W^{\frac{1}{2}}(b\Omega^-)$ from the Sobolev Trace Theorem. Thus the pairing between f and ϕ in (2.1) is well-defined since

$$|\int_{b\Omega^-} f \wedge h| \leq \|f\|_{W^{-\frac{1}{2}}(b\Omega)} \|h\|_{W^{\frac{1}{2}}(b\Omega^-)} \leq \|f\|_{L^2(\Omega)} \|h\|_{W^1(\Omega^-)}.$$

We also note that the pairing in (2.1) is independent of the choice of the representation function $[f]$. Let \tilde{f} be another representation of $[f]$, then $\tilde{f} = f + \bar{\partial}u$ for some element of the form $\bar{\partial}u \in L_{(p,n-1)}^2(\Omega)$ with $u \in L_{(p,n-2)}^2(\Omega)$. Using Friedrichs'

lemma, there exists a sequence $\{u_\nu\}$ such that $u_\nu \in C_{(p,n-2)}^\infty(\bar{\Omega})$ such that $u_\nu \rightarrow u$ in $L_{(p,n-2)}^2(\Omega)$ and $\bar{\partial}u_\nu \rightarrow \bar{\partial}u$ in $L_{(p,n-1)}^2(\Omega)$. It follows from Stokes' Theorem that

$$\begin{aligned} \int_{b\Omega^-} \bar{\partial}u \wedge h &= \lim_{\nu \rightarrow \infty} \int_{b\Omega^-} \bar{\partial}u_\nu \wedge h \\ &= \lim_{\nu \rightarrow \infty} (-1)^{p+n} \int_{b\Omega^-} u_\nu \wedge \bar{\partial}h = 0, \quad h \in H_{W^1}^{(n-p,0)}(\Omega^-). \end{aligned}$$

Thus the pairing (2.1) is well-defined.

If we assume that f satisfies the condition

$$\int_{b\Omega^-} f \wedge \phi = 0, \quad \phi \in W_{(n-p,0)}^1(\Omega^-) \cap \text{Ker}(\bar{\partial}),$$

from Theorem 1.2, there exists a $\bar{\partial}$ -closed form $F \in W_{(p,n-2)}^{-1}(\Omega_1)$ which is equal to f on Ω and one can find a solution $u \in L_{(p,n-2)}^2(\Omega)$ satisfying $\bar{\partial}u = f$. This implies that $[f] = 0$. Thus there is a 1-1 map from $H^{(p,n-1)}(\Omega)$ to $H_{W^1}^{(n-p,0)}(\Omega^-)$.

On the other hand, suppose that f is a bounded linear functional on $H_{W^1}^{(n-p,0)}(\Omega^-)$. We will show that l can be represented by some $[f]$ in (2.1). Since we assume that Ω^- is pseudoconvex and has smooth boundary, one has the duality for holomorphic space $H^1(\Omega^-) = L^2(\Omega^-) \cap \text{Ker}(\bar{\partial})$ and $H^{-1}(\Omega^-)$ (see **[BB]**). If the $\bar{\partial}$ -Neumann operator is exact regular on $W^1(\Omega^-)$, we can use the duality between the usual L^2 spaces. Otherwise, one can use the exact regularity for the weighted $\bar{\partial}$ -Neumann operator with weights $t|z|^2$ for sufficiently large $t > 0$. The weight function can be viewed as the bundle metric $e^{-t|z|^2}$ for the trivial line bundle \mathbb{C} and the dual space will be equipped with the dual metric $e^{t|z|^2}$ for \mathbb{C} . In particular the pairing (2.1) is well-defined. For simplicity, we assume that the $\bar{\partial}$ -Neumann operator is exact regular. But all the arguments remain the same if we use weighted spaces with the dual weighted norms.

Thus l can be represented by $(n-p,0)$ -form g with distribution coefficients in $H^{-1}(\Omega^-) = \text{ker}(\bar{\partial}) \cap W^{-1}(\Omega^-)$. Extending g to be zero outside Ω^- , then $\star g$ is a (p,n) -form on Ω_1 , a top degree form which is always $\bar{\partial}$ -exact. The extension by zero of g results in a form which is in $W^{-1}(\Omega_1)$. This is due to the fact that holomorphic functions in $W^{-1}(\Omega^-)$ is also in the dual of $W^1(\Omega^-)$. We remark that for a general function or forms, this is not true. But when the functions or forms are harmonic, then the dual space of W_0^1 , denoted by W^{-1} , coincides with the dual space of W^1 for domains with smooth boundary. For detailed explanation of this subtle point, we refer the reader to the paper by Boas (see Appendix B in **[Boa]** where the dual space of W^1 is denoted by W_*^{-1}).

Thus we have that $\star g = \bar{\partial}U$ on Ω_1 for some $U \in L_{(p,n-1)}^2(\Omega_1)$. Let $f = U$ on Ω . It follows that $\bar{\partial}f = 0$ on Ω and the linear functional

$$l(h) = \int_{\Omega^-} \star g \wedge h = \int_{\Omega^-} \bar{\partial}U \wedge h = \int_{b\Omega^-} f \wedge h, \quad h \in H_{W^1}^{(n-p,0)}(\Omega^-).$$

Since $f \in L_{(p,n-1)}^2(\Omega)$, we have that the bounded linear functional l is represented by $[f] \in H^{(p,n-1)}(\Omega)$. This proves the theorem for $k = 0$.

Suppose that $k \geq 1$ and $f \in W_{(p,n-1)}^k(\Omega)$ and $\bar{\partial}f = 0$. We define a pairing between $H_{W^k}^{(p,n-1)}(\Omega)$ and $H_{W^{-k+1}}^{(n-p,0)}(\Omega^-)$ by

$$l : H_{W^k}^{(p,n-1)}(\Omega) \times H_{W^{-k+1}}^{(n-p,0)}(\Omega^-)$$

$$(2.2) \quad l([f], h) = \int_{b\Omega^-} f \wedge h.$$

It is easy to see that the pairing (2.2) is well-defined as before. If f satisfies the condition

$$\int_{b\Omega^-} f \wedge \phi = 0, \quad \phi \in W_{(n-p,0)}^{-k+1}(\Omega^-) \cap \text{Ker}(\bar{\partial}),$$

there exists a $\bar{\partial}$ -closed form $F \in W_{(p,n-1)}^{k-1}(\Omega_1)$ which is equal to f on Ω and one can find a solution $u \in W_{(p,n-2)}^k(\Omega)$ satisfying $\bar{\partial}u = f$. For a proof, see Corollary 2 in the recent paper by Chakrabarti-Shaw [CS2]. This implies that $[f] = 0$.

Thus there is a one to one map from $H_{W^k}^{(p,n-1)}(\Omega)$ to $H_{W^{-k+1}}^{(n-p,0)}(\Omega^-)'$. Any element in $H_{W^{-k+1}}^{(n-p,0)}(\Omega^-)'$ can be identified as a (p, n) -form $\star g$ with $H_{W^{k-1}}(\Omega^-)$ -coefficients. Thus repeating the same arguments as before, there exists $f \in W^k(\Omega)$ with $\bar{\partial}f = 0$ such that any bounded linear functional can be given by f in the equation (2.1). \square

We remark that in Theorem 2.1, the boundary of Ω is assumed to be C^∞ smooth in order to have the duality for all $k \geq 0$. For each fixed k , the duality result holds for sufficiently smooth (depending on k) domains Ω^- and Ω_1 . In particular, Theorem 2.1 holds for $k = 1$ for much less smooth domains Ω_1 and Ω^- . In the following, we will only assume that the boundary for Ω^- be Lipschitz, i.e., locally it is the graph of a Lipschitz function.

Let $\bar{\partial}_c : L_{(p,n-1)}^2(\Omega^-) \rightarrow L_{(p,n)}^2(\Omega^-)$ be the minimal closure of $\bar{\partial}$. By this we mean that $f \in \text{Dom}(\bar{\partial}_c)$ if and only if that there exists a sequence of smooth forms f_ν in $C_{(p,n-1)}^\infty(\Omega)$ compactly supported in Ω such that $f_\nu \rightarrow f$ and $\bar{\partial}f_\nu \rightarrow \bar{\partial}f$ in L^2 . Let ϑ be the dual of $\bar{\partial}_c$. Then ϑ is equal to the maximal closure of the operator

$$\vartheta : L_{(p,n)}^2(\Omega^-) \rightarrow L_{(p,n-1)}^2(\Omega^-).$$

We set

$$\square_{(p,n)}^c(\Omega^-) = \bar{\partial}_c \vartheta : L_{(p,n)}^2(\Omega^-) \rightarrow L_{(p,n)}^2(\Omega^-).$$

The kernel of $\square_{(p,n)}^c(\Omega^-)$ is denoted by $\mathcal{H}_c^{(p,n)}(\Omega^-)$, the space of harmonic forms of degree (p, n) with compact support.

Theorem 2.2. *Let $\Omega \subset \subset \mathbb{C}^n$ be the annulus domain $\Omega = \Omega_1 \setminus \bar{\Omega}^-$ between two bounded pseudoconvex domains Ω_1 and Ω^- with $\Omega^- \subset \subset \Omega_1$, $n \geq 2$. We assume that Ω_1 has C^2 boundary and Ω^- has Lipschitz boundary. The space $H_{W^1}^{(p,n-1)}(\Omega)$ is of infinite dimension and we have the isomorphism:*

$$H_{W^1}^{(p,n-1)}(\Omega) \simeq \mathcal{H}_c^{(p,n)}(\Omega^-) \simeq H^{(n-p,0)}(\Omega^-).$$

PROOF. From the closed range property for $\bar{\partial}$ on Ω^- and its L^2 dual for all degrees, it follows (see [CS2]) that range of $\bar{\partial}_c$ is also closed for all degrees. In particular, we have

$$L^2_{(p,n)}(\Omega^-) = \text{Range}(\bar{\partial}_c) \oplus \text{Ker}(\vartheta).$$

Here we only need the boundary Ω^- to be Lipschitz smooth (see [CS2] for details).

This gives that

$$(2.3) \quad \mathcal{H}_c^{(p,n)}(\Omega^-) = \text{Ker}(\vartheta).$$

Using star operator, one has that

$$(2.4) \quad \text{Ker}(\vartheta) \simeq H^{(n-p,0)}(\Omega^-).$$

From the extension of the $\bar{\partial}$ -closed forms from the annulus to Ω_1 as in the proof of Theorem 1.2, we will show that $H_{W^1}^{(p,n-1)}(\Omega)$ is isomorphic to the perp of $\text{Range}(\bar{\partial}_c)$. To see this, let $f \in W^1_{(p,n-1)}(\Omega)$ and let $\bar{\partial}f = 0$ in Ω . We extend f to be a form $\tilde{f} \in W^1_{(p,n-1)}(\Omega_1)$. The equation

$$(2.5) \quad \bar{\partial}_c u = \bar{\partial} \tilde{f}$$

for some $u \in L^2_{(p,n-1)}(\Omega^-)$ if and only if

$$(2.6) \quad \int_{b\Omega^-} f \wedge \phi = 0, \quad \phi \in L^2_{(n-p,0)}(\Omega^-) \cap \text{Ker}(\bar{\partial}).$$

For a proof of the equivalence of (2.5) and (2.6), see the proof Proposition 5 in [CS2].

In this case, f can be extended to be $\bar{\partial}$ -closed form F where

$$F = \begin{cases} f, & z \in \Omega, \\ \tilde{f} - u, & z \in \Omega^-. \end{cases}$$

It follows that $\bar{\partial}F = 0$ in Ω_1 and $F = f$ on Ω . The form F is in $L^2_{(p,n-1)}(\Omega_1)$ but F is in $W^1(\Omega)$ since $F = f$ on Ω . Since we assume that the boundary Ω_1 is C^2 , we can find a solution (see [Ha]) $F = \bar{\partial}U$ for some $U \in W^1_{(p,n-2)}(\Omega_1)$. In fact we can use the solution $U = F + \bar{\partial}_t^* N_t F$ by the weighted $\bar{\partial}$ -Neumann operator N_t on Ω_1 . Then U is in W^1 near the boundary Ω_1 from the boundary regularity for the weighted $\bar{\partial}$ -Neumann operator. Since the weighted $\bar{\partial}$ -Neumann operator N_t is elliptic in the interior of Ω_1 , $N_t F$ is in $W^2(\Omega, \text{loc})$. Thus the solution U is in $W^1(\Omega, \text{loc})$. Thus U is in $W^1_{(p,n-2)}(\Omega_1)$. This shows that for any $[f] \in H_{W^1}^{(p,n-1)}(\Omega)$, $[f] = 0$ if and only if (2.5) or (2.6) is satisfied for any representation $f \in W^1_{(p,n-1)}(\Omega)$. Repeating the arguments as in Theorem 2.1 and using (2.3) and (2.4), we have proved the theorem. \square

3. The isomorphism between $H^{(p,n-1)}(\Omega)$ and $H^{(n-p,0)}(\Omega^-)$

In this section we will further establish the isomorphism between the spaces $H^{(p,n-1)}(\Omega)$ and $H^{(n-p,0)}(\Omega^-)$.

Theorem 3.1. *Let $\Omega \subset \subset \mathbb{C}^n$ be the annulus domain $\Omega = \Omega_1 \setminus \overline{\Omega^-}$ between two pseudoconvex domains Ω_1 and Ω^- and $\Omega^- \subset \subset \Omega_1$, $n \geq 2$. We assume that the boundary of Ω^- is C^2 smooth. Then we have the isomorphism:*

$$(3.1) \quad H^{(p,n-1)}(\Omega) \simeq H^{(n-p,0)}(\Omega^-).$$

Furthermore, if we assume that Ω has C^2 smooth boundary, then we have the isomorphism:

$$(3.2) \quad H_{W^1}^{(p,n-1)}(\Omega) \simeq H^{(p,n-1)}(\Omega).$$

PROOF. It follows from Theorem 2.2 that

$$H_{W^1}^{(p,n-1)}(\Omega) \simeq H^{(n-p,0)}(\Omega^-).$$

Thus it suffices to prove (3.1).

Let $h \in H^{(n-p,0)}(\Omega^-)$. We will associate h with a $\bar{\partial}$ -closed form h^+ in Ω as follows:

Let ρ be a normalized C^2 defining function for Ω^- . Since h has holomorphic coefficients in $L^2(\Omega^-)$, it is well known that h has $W^{-\frac{1}{2}}$ boundary values on $b\Omega^-$. Let $h_1 = h \wedge \bar{\partial}\rho$. The form $\star h_1$ is a $(p, n-1)$ -form on Ω^- and it has boundary value in $W^{-\frac{1}{2}}(b\Omega^-)$. We denote the restriction of $\star h_1$ to $b\Omega^-$ by

$$(3.3) \quad h_b = \star h_1|_{b\Omega^-}.$$

Let $h^+ = B^+ h_b$ and $h^- = B^- h_b$ be the Bochner transform of h_b defined by

$$(3.4) \quad h^+ = B^+ h_b = \int_{b\Omega^-} B(\zeta, z) \wedge h_b, \quad z \in \Omega,$$

$$(3.5) \quad h^- = B^- h_b = \int_{b\Omega^-} B(\zeta, z) \wedge h_b, \quad z \in \Omega^-.$$

We have the jump formula:

$$h_b = B^+ h_b - B^- h_b = h^+ - h^-$$

in terms of distributions. Also each h^+ and h^- are $\bar{\partial}$ -closed and L^2 on Ω and Ω^- respectively.

We define a map $l^+ : H^{(n-p,0)}(\Omega^-) \rightarrow H^{(p,n-1)}(\Omega)$ by

$$l^+ h = [h^+], \quad h \in H^{(n-p,0)}(\Omega^-)$$

where h^+ is defined by (3.4). Since h^+ is $\bar{\partial}$ -closed on Ω and has L^2 coefficients, the map l^+ is well-defined.

We next show that l^+ is one to one. If $l^+(h) = [h^+] = 0$ for some $h \in H^{(n-p,0)}(\Omega^-)$, we will show that $h = 0$. Since $[h^+] = 0$, this implies that h^+ can be represented by a $\bar{\partial}$ -exact form and there exists $u^+ \in L^2_{(p,n-2)}(\Omega)$ such that

$$(3.6) \quad h^+ = \bar{\partial}u^+.$$

Let h^- be defined by (3.5). Since Ω^- is pseudoconvex, we have

$$(3.7) \quad h^- = \bar{\partial}u^-$$

for some $u^- \in L^2_{(p,n-2)}(\Omega^-)$. It follows from (3.6) and (3.7) that that for each $g \in H^{(n-p,0)}_{W^1}(\Omega^-)$,

$$\int_{b\Omega^-} h_b \wedge g = \int_{b\Omega^-} (h^+ - h^-) \wedge g = \int_{b\Omega^-} (\bar{\partial}u^+ - \bar{\partial}u^-) \wedge g = 0.$$

This implies that h_b is a linear functional vanishing on $H^{(n-p,0)}_{W^1}(\Omega^-)$. But from the regularity for the weighted $\bar{\partial}$ -Neumann operator for $W^1_{(n-p,1)}(\Omega)$ (since we assume that Ω^- is C^2), we have that the space $H^{(n-p,0)}_{W^1}(\Omega^-)$ is dense in $H^{(n-p,0)}_{L^2}(\Omega^-)$. Since the functional vanishes on a dense subspace, h_b must be zero. This proves that $h = 0$ if $l^+h = [h^+] = 0$. Thus l^+ is one to one.

To show that l^+ is onto, take an element $F \in L^2_{(p,n-1)}(\Omega)$ such that $\bar{\partial}F = 0$. For simplicity, we assume that $p = n$. We will construct a holomorphic function h in $L^2(\Omega)$ such that $l^+h = F$. Note that from Theorem 1.4, any element $[F]$ can be represented by a harmonic form and we may assume that F is in $\mathcal{H}_{(n,n-1)}(\Omega)$. This implies that $\bar{\partial}F = \bar{\partial}^*F = 0$. It follows that F has boundary value with $W^{-\frac{1}{2}}$ -coefficients. Choose a special orthonormal frame field basis $w_1, \dots, w_n = \sqrt{2}\bar{\partial}\rho$ for $(1,0)$ -forms. Then near the boundary F written in the special orthonormal frame fields as

$$F = \sum_{i=1}^n F_i(dV \lrcorner \bar{w}_i)$$

where $dV = w_1 \wedge \bar{w}_1 \dots w_n \wedge \bar{w}_n$ is the volume element. Using

$$\star \bar{w}_n = \star w_n = \frac{1}{2} \star d\rho = \frac{1}{2} d\sigma$$

on $b\Omega^-$, we have

$$F|_{b\Omega^-} = F_n dV \lrcorner \bar{w}_n|_{b\Omega^-} = \frac{1}{2} F_n d\sigma,$$

where $d\sigma$ is the surface element on $b\Omega^-$.

We claim that \bar{F}_n is a CR distribution on $b\Omega^-$. To see this, note that $\bar{\partial}\star F = 0$ since $\partial F = 0$. Restricted to the boundary, this implies that $\bar{\partial}\bar{F}_n \wedge \bar{w}_n = 0$ on $b\Omega^-$. Thus \bar{F}_n is a CR distribution in $W^{-\frac{1}{2}}(b\Omega^-)$. Let $h = \bar{F}_n$ be the holomorphic extension of \bar{F}_n from the boundary to Ω^- . Then h is an L^2 holomorphic function on Ω^- . Let $h^+ = l^+h$ be the $\bar{\partial}$ -closed form in $L^2_{(n,n-1)}(\Omega)$. It remains to show that $[h^+] = [F]$. This follows from $h^+ - h^- = h_b = \star(\bar{F}_n \wedge \bar{\partial}\rho) = F_n dV \lrcorner \bar{w}_n$ on $b\Omega^-$. If we define

$$G = \begin{cases} F - h^+, & z \in \Omega \\ h^-, & z \in \Omega^- \end{cases}$$

Then G is an L^2 $\bar{\partial}$ -closed form in Ω_1 . Thus we have $G = \bar{\partial}U$ for some $U \in L^2_{(n,n-2)}(\Omega_1)$ is $\bar{\partial}$ -exact on Ω_1 . Thus $[F] = [h^+]$ in $H^{(n,n-1)}(\Omega)$. This proves that l^+ is onto. The theorem is proved.

Corollary 3.2. *Let Ω be the same as Theorem 3.1. Each element f in the harmonic space $\mathcal{H}_{(p,n-1)}(\Omega)$ can be represented by some h^+ , where h is a harmonic form in $L^2_{(n-p,0)}(\Omega^-)$. We have the following representation for the harmonic space*

$$\mathcal{H}_{(p,n-1)}(\Omega) = \{h^+ \mid h \in L^2_{(n-p,0)}(\Omega^-), \bar{\partial}h = 0.\}$$

where h^+ is defined by (3.4).

PROOF. We have proved that for every $f \in \mathcal{H}_{(p,n-1)}(\Omega)$, we can write $f = \star h \partial \rho = \bar{h} \star \bar{\partial} \rho$ for some holomorphic h in $L^2_{(n-p,0)}(\Omega^-)$. On the other hand, any $h \in \mathcal{H}_{(n-p,0)}(\Omega^-)$, the associated $B^+(h_b) = h^+$ is in $L^2_{(p,n-1)}(\Omega)$. The form h^+ is automatically $\bar{\partial}$ -closed. To see that it is in the domain of $\bar{\partial}^*$ and $\bar{\partial}^* \bar{h}^+ = 0$, we repeat the arguments before and the corollary is proved.

Remarks:

- (1) All the results can be extended to any annulus between two pseudoconvex domains in a Stein manifold with trivial modification. It can also be applied to an annulus between two pseudoconvex domains in complex manifolds if one has the existence and the W^1 regularity of the $\bar{\partial}$ -Neumann operator on the pseudoconvex domain Ω^- . We refer the reader to some related results in [HI] (see also [CaS] and [CS1] and [CS2]).
- (2) If we assume that the boundary is C^∞ smooth, we can also have the isomorphism between $H^{(p,n-1)}(\Omega)$ and $H_{W^k}^{(p,n-1)}(\Omega)$ for all k following the same proof.

4. The null space for the $\bar{\partial}$ -Neumann operator between balls

When the domain $\Omega = \{z \in \mathbb{C}^n \mid 0 < R_0 < |z| < R_1\}$ is the annulus between two balls centered at 0, the harmonic space $\mathcal{H}_{(0,n-1)}$ has been computed explicitly in Hörmander (see equation (2.3) in [Hör2]). He proved that any $(0, n-1)$ -form f is in the null space of the $\bar{\partial}$ -Neumann operator if and only if

$$(4.1) \quad f = \sum_1^n (-1)^j \frac{\bar{z}_j}{|z|^{2n}} h \left(\frac{\bar{z}}{|z|^2} \right) d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{j-1} \wedge d\bar{z}_{j+1} \wedge \cdots \wedge d\bar{z}_n,$$

where h is a holomorphic function in $L^2(\Omega^*)$, $\Omega^* = \{z \in \mathbb{C}^n \mid |z| < \frac{1}{R_0}\}$. It is easy to check that $\bar{\partial} f = \partial f = 0$.

To see that f is in the domain of $\bar{\partial}^*$, we note that $f = \frac{1}{|z|^{2n}} h \left(\frac{\bar{z}}{|z|^2} \right) \star \bar{\partial} \rho$ where $\rho = |z|^2$. Thus $f \in \text{Dom}(\bar{\partial}^*)$. We will show that the representation by the Bochner transform stated the harmonic forms from Corollary 3.2 agrees with the formula (4.1). For simplicity, we will assume that the inner ball is the unit ball, i.e., $R_0 = 1$ and h is a holomorphic $(n, 0)$ form in $L^2_{(n,0)}(B_1)$. Let $\rho(z) = |z|^2 - 1$. Then $\bar{h}(\bar{z})$ is holomorphic with L^2 coefficients. The Bochner transform described in Section 3 is given by

$$(4.2) \quad \begin{aligned} B^+ h_b &= \int_{\zeta \in bB_1} B_0(\zeta, z) \wedge h(\bar{\zeta}) \wedge \star \bar{\partial} \rho(\zeta) \\ &= \int_{\zeta \in bB_1} B_0(\zeta, z) \wedge h\left(\frac{\bar{\zeta}}{|\zeta|^2}\right) \wedge \sum_j \frac{\bar{\zeta}_j}{|\zeta|^{2n}} [d\bar{\zeta}_j] \end{aligned}$$

where $[d\bar{\zeta}_j]$ denote the $(n, n-1)$ -form $dV \lrcorner d\bar{\zeta}_j$. It is easy to see that

$$\bar{\partial} \left(\sum_j \frac{\bar{\zeta}_j}{|\zeta|^{2n}} [d\bar{\zeta}_j] \right) = 0, \quad \zeta \neq 0.$$

We also have

$$\begin{aligned} \bar{\partial} \left(h \left(\frac{\bar{\zeta}}{|\zeta|^2} \right) \wedge \sum_j \frac{\bar{\zeta}_j}{|\zeta|^{2n}} [d\bar{\zeta}_j] \right) &= \sum_k \sum_j \frac{\partial h}{\partial w_k} \left(\frac{\delta_{jk}}{|\zeta|^2} - \frac{\bar{\zeta}_k \zeta_j}{|\zeta|^4} \right) \frac{\bar{\zeta}_j}{|\zeta|^{2n}} dV \\ &= \sum_k \frac{\partial h}{\partial w_k} \left(\frac{\bar{\zeta}_k}{|\zeta|^2} - \frac{\bar{\zeta}_k}{|\zeta|^2} \right) dV = 0. \end{aligned}$$

Applying Stokes's Theorem to (4.2), we see that

$$B^+ h_b = \sum \frac{\bar{z}_j}{|z|^{2n}} h \left(\frac{\bar{z}}{|z|^2} \right) dV \lrcorner d\bar{z}_j.$$

We mention that in [Hör2], it is also proved that for any $f \in \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*) \cap \mathcal{H}_{(n,n-1)}(\Omega)^\perp$,

$$(4.3) \quad \max(n-2, 1) \|f\|^2 \leq R_1^2 \|\bar{\partial}f\|^2 + \|\bar{\partial}^*f\|^2.$$

Notice that the constant in (4.3) is independent of the inner diameter R_0 . It is not known if one can have such estimates on the more general annulus between two pseudoconvex domains.

References

- [BB] Bell, S. R. and Boas, H. P., *Regularity of the Bergman projection and duality of holomorphic function spaces*, Math. Ann. **267** (1984), 473-478.
- [Boa] Boas, H. P., *The Szegő projection: Sobolev estimates in regular domains*, Trans. Amer. Math. Soc., **300** (1987), 109-132.
- [BS] Boas, H. P. and Shaw, M.-C., *Sobolev Estimates for the Lewy Operator on Weakly pseudo-convex boundaries*, Math. Annalen **274** (1986), 221-231.
- [BSt] Boas, H. P., and Straube, E. J., *Sobolev estimates for the $\bar{\partial}$ -Neumann operator on domains in \mathbb{C}^n admitting a defining function that is plurisubharmonic on the boundary*, Math. Zeit., **206** (1991), 81-88.
- [CaS] J. Cao and M.-C. Shaw, *The $\bar{\partial}$ -Cauchy problem and nonexistence of Lipschitz Levi-flat hypersurfaces in $\mathbb{C}P^n$ with $n \geq 3$* , Math. Zeit. **256** (2007), 175-192.
- [CS1] D. Chakrabarti and M.-C. Shaw, *The Cauchy-Riemann equations on product domains*, To appear in Math. Annalen (on line July 27, 2010).
- [CS2] D. Chakrabarti and M.-C. Shaw, *L^2 Serre duality on domains in complex manifolds with applications*, To appear in Trans. Amer. Math. Society.
- [CS] Chen, S.-C. and Shaw, M.-C., *Partial Differential Equations in Several Complex Variables*, American Math. Society-International Press, Studies in Advanced Mathematics, Volume 19, Providence, R.I., 2001.
- [Fo] Folland, G. B., *The tangential Cauchy-Riemann complex on spheres*, Trans. Amer. Math. Society **171** (1972), 83-133.
- [FK] Folland, G. B. and Kohn, J. J., *The Neumann Problem for the Cauchy-Riemann Complex*, Ann. Math. Studies 75, Princeton University Press, Princeton, N.J., 1972.
- [Gr] Grisvard, P., *Elliptic Problems in Nonsmooth Domains*, Pitman, Boston, 1985.
- [Ha] Harrington, P.S., *Sobolev Estimates for the Cauchy-Riemann Complex on C^1 Pseudoconvex Domains*, Math. Z. **262** (2009), 199-217.
- [HI] Henkin, G. M. and Iordan, A., *Regularity of $\bar{\partial}$ on pseudoconcave compacts and applications*, (see also Erratum: Asian J. Math., vol **7**, (2003) No. 1, pp. 147-148), Asian J. Math. **4** (2000), 855-884.
- [Hör1] Hörmander, L., *L^2 estimates and existence theorems for the $\bar{\partial}$ operator*, Acta Math. **113** (1965), 89-152.
- [Hör2] Hörmander, L., *The null space of the $\bar{\partial}$ -Neumann operator*, Ann. Inst. Fourier (Grenoble) **54** (2004), 1305-1369.

- [Hor] Hortmann, M., *Über die Lösbarkeit der $\bar{\partial}$ -Gleichung mit Hilfe von L^p , C^k , und D' -stetigen Integraloperatoren*, Math. Ann. **223** (1976), 139-156.
- [Ko1] Kohn, J.J., *Harmonic integrals on strongly pseudoconvex manifolds, I*, Ann. of Math. **78** (1963), 112-148.
- [Ko2] Kohn, J. J., *Global regularity for $\bar{\partial}$ on weakly pseudoconvex manifolds*, Trans. Amer. Math. Soc., **181** (1973), 273–292.
- [Ko3] Kohn, J.J., *The range of the tangential Cauchy-Riemann operator*, Duke Math. Journ. **53** (1986), 525-545.
- [KoR] Kohn, J. J., and Rossi, H., *On the extension of holomorphic functions from the boundary of a complex manifold*, Ann. Math., **81** (1965), 451-472.
- [LM] Lions, J.-L., and Magenes, E., *Non-Homogeneous Boundary Value Problems and Applications, Volume I*, Springer-Verlag, New York, 1972.
- [MS] Michel, J., and Shaw, M.-C., *Subelliptic estimates for the $\bar{\partial}$ -Neumann operator on piecewise smooth strictly pseudoconvex domains*, Duke Math. J., **93** (1998), 115–128.
- [Ra] Range, R. M., *Holomorphic Functions and Integral Representations in Several Complex Variables*, Graduate Texts in Math., Vol.108, Springer-Verlag, N.Y., 1986.
- [Sh1] Shaw, M.-C., *Global solvability and regularity for $\bar{\partial}$ on an annulus between two weakly pseudoconvex domains*, Trans. Amer. Math. Soc., **291** (1985), 255-267.
- [Sh2] Shaw, M.-C., *L^2 estimates and existence theorems for the tangential Cauchy-Riemann complex.*, Invent. Math. **82** (1985), 133-150.
- [Sh3] Shaw, M.-C., *L^2 estimates and existence theorems for $\bar{\partial}_b$ on Lipschitz boundaries*, Math. Zeit. **244** (2003), 91-123.
- [Sh4] M.-C. Shaw, *The closed range property for $\bar{\partial}$ on domains with pseudoconcave boundary*, Proceedings for the Fribourg conference, Trends in Mathematics (2010), 307-320.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
USA

E-mail address: Shaw.1@nd.edu