

HAMILTONICITY OF THE CARTESIAN PRODUCT OF TWO DIRECTED CYCLES MINUS A SUBGROUP

VICTORIA BARONE, MATTHEW MAUNTEL, AND MICAH MILLER

ABSTRACT. Let $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ be the product of two directed cycles minus a subgroup. Also, let $A = \frac{a}{c}$ and $B = \frac{b}{d}$. We show that this digraph is Hamiltonian if and only if there exist positive integers m and n such that $Am + Bn = AB - 1$ and $\gcd(dm, cn) = 1$.

1. INTRODUCTION

Improving on the work of [3] Curran [2] used the theory of torus knots to prove that the Cartesian product $\mathbb{Z}_a \times \mathbb{Z}_b$ of two directed cycles is Hamiltonian if and only if there exists a pair of relatively prime positive integers m and n such that $am + bn = ab$. Penn and Witte [5] used the same method to prove a similar result for digraphs of the form $\mathbb{Z}_a \times \mathbb{Z}_b - \{(0, 0)\}$, the digraph $\mathbb{Z}_a \times \mathbb{Z}_b$ with one vertex removed. This digraph $\mathbb{Z}_a \times \mathbb{Z}_b - \{(0, 0)\}$ is Hamiltonian if and only if there exists a pair of relatively prime positive integers m and n such that $am + bn = ab - 1$.

Note that removing one vertex from $\mathbb{Z}_a \times \mathbb{Z}_b$ as in [5] is equivalent to removing the subgroup $\mathbb{Z}_1 \times \mathbb{Z}_1$. In this paper, we consider a natural generalization by removing a subgroup of the form $\mathbb{Z}_c \times \mathbb{Z}_d$ and find necessary and sufficient conditions for the digraph $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ to be Hamiltonian. Here, \mathbb{Z}_c is a subgroup of \mathbb{Z}_a and \mathbb{Z}_d is a subgroup of \mathbb{Z}_b . See Figure 1 for a picture of the digraph $(\mathbb{Z}_8 \times \mathbb{Z}_6) - (\mathbb{Z}_2 \times \mathbb{Z}_2)$.

2. BACKGROUND

We recall some definitions and results that will be useful. We refer the reader to [1] for the basic language of digraphs. We are considering digraphs of the form $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$, where \mathbb{Z}_c and \mathbb{Z}_d are subgroups of \mathbb{Z}_a and \mathbb{Z}_b , respectively. Thus, we start with $\mathbb{Z}_a \times \mathbb{Z}_b$, remove vertices that belong to $\mathbb{Z}_c \times \mathbb{Z}_d$, and consider the induced subgraph on the remaining vertices.

Definition 2.1. Let H be a Hamiltonian circuit on $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. Then a vertex (x, y) **travels by** $(1, 0)$ if the out-edge of (x, y) is an in-edge of $(x + 1, y)$ in H . Similarly, a vertex (x, y) **travels by** $(0, 1)$ if the out-edge of (x, y) is an in-edge of $(x, y + 1)$ in H .

Note that in a Hamiltonian circuit, each vertex travels by either $(1, 0)$ or by $(0, 1)$.

Definition 2.2. Let $g = \gcd(a, b)$. For any integer p , let $\langle p \rangle$ be the subset of $\mathbb{Z}_a \times \mathbb{Z}_b$ consisting of pairs (x, y) such that $x + y = p$ modulo g .

The authors thank Daniel C. Isaksen and Frank Connolly for their guidance through the course of the REU at the University of Notre Dame in the summer of 2002. The authors were supported by an NSF REU grant number DMS-0139018.

The subset $\langle 0 \rangle$ is the subgroup of $\mathbb{Z}_a \times \mathbb{Z}_b$ generated by $(1, -1)$, and $\langle p \rangle$ is the coset $(p, 0) + \langle 0 \rangle$. The subgroup $\langle 0 \rangle$ has index g , so there are exactly g distinct cosets.

The following lemma shows why such cosets are useful.

Lemma 2.3. *Let H be a spanning circuit of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. Suppose that (x, y) , $(x + 1, y)$, and $(x + 1, y - 1)$ all belong to $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. Then (x, y) travels by $(1, 0)$ if and only if $(x + 1, y - 1)$ travels by $(1, 0)$. Also, (x, y) travels by $(0, 1)$ if and only if $(x + 1, y - 1)$ travels by $(0, 1)$.*

Curran and Witte [2] give a proof this lemma.

We recall the following result from [5].

Theorem 2.4. *The digraph $\mathbb{Z}_a \times \mathbb{Z}_b - \{(0, 0)\}$ is Hamiltonian if and only if there exist relatively prime positive integers m and n such that $am + bn = ab - 1$.*

In the previous theorem, the numbers m and n have useful geometric interpretations. If we embed $\mathbb{Z}_a \times \mathbb{Z}_b$ in the torus in the obvious way, then a Hamiltonian circuit is an embedded loop, and its knot class is equal to (m, n) . See [6] for more details on knot classes.

3. THE MAIN THEOREM

We now come to our main result.

Theorem 3.1. *Let $A = \frac{a}{c}$ and $B = \frac{b}{d}$. Then $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ is Hamiltonian if and only if there exist positive integers m and n such that*

- (1) $Am + Bn = AB - 1$, and
- (2) $\gcd(dm, cn) = 1$.

In order to prove this theorem, we need the following facts about Hamiltonian digraphs.

Lemma 3.2. *If $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ is Hamiltonian, then $\gcd(A, B) = 1$.*

Proof. Suppose for contradiction that the coset $\langle -1 \rangle$ contains no vertex of $\mathbb{Z}_c \times \mathbb{Z}_d$. The vertex $(-1, 0)$ must travel by $(0, 1)$, so a repeated application of Lemma 2.3 implies that every vertex in $\langle -1 \rangle$ must also travel by $(0, 1)$. This is impossible because the vertex $(0, -1)$ certainly travels by $(1, 0)$. By contradiction, $\langle -1 \rangle$ contains a vertex $(\alpha a, \beta b)$ for some integers α and β .

From the definition of a coset we find that $\alpha A + \beta B = -1 \pmod{g}$. There exists an integer γ such that $\alpha A + \beta B = -1 + g\gamma$. Now $\gcd(A, B)$ divides g , so $\gcd(A, B)$ divides $\alpha A + \beta B - g\gamma = -1$. \square

Corollary 3.3. *If $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ has a Hamiltonian circuit, then it is unique.*

Proof. We will show that every coset contains a vertex $(\alpha A, \beta B)$ for some α and β in \mathbb{Z} . The repeated application of Lemma 2.3 then gives the result.

From the proof of Lemma 3.2, we know that $\alpha A + \beta B = -1 \pmod{g}$, for some integers α and β . Multiplying this equation by $-p$ we get $(-\alpha p)A + (-\beta p)B = p \pmod{g}$. From the definition of a coset, we know $((-\alpha p)A, (-\beta p)B)$ is in $\langle p \rangle$. \square

The following corollary tells us that any Hamiltonian circuit of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ must be periodic.

Corollary 3.4. *Suppose $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ has a Hamiltonian circuit. Let α and β be integers. Then (x, y) and $(x + \alpha A, y + \beta B)$ travel in the same direction.*

Proof. Let H be a Hamiltonian circuit of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. Then $\phi(x, y) = (x + \alpha A, y + \beta B)$ is an automorphism of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. So $\phi(H)$ is a Hamiltonian circuit of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ as well. But the Hamiltonian circuit is unique (by Corollary 3.3), so $\phi(H) = H$. Thus, any vertex v travels in the same direction as the vertex $\phi(v)$. \square

Now we are ready to prove Theorem 3.1.

Proof. First suppose that $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ is Hamiltonian. Let H be the Hamiltonian circuit, and let (m_0, n_0) be its knot class. We know that m_0 and n_0 are relatively prime positive integers [6]. We also know that, in H , there are exactly am_0 vertices that travel by $(1, 0)$ and exactly bn_0 vertices that travel by $(0, 1)$, and that H contains each vertex in $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. By counting vertices, $am_0 + bn_0 = ab - cd$. We divide this equation by cd and get $A\frac{m_0}{d} + B\frac{n_0}{c} = AB - 1$.

Consider the sets of vertices $U = \{(0, y) \mid y \in \mathbb{Z}_b\}$ and $V = \{(x, 0) \mid x \in \mathbb{Z}_a\}$. By Corollary 3.4, the number of vertices in U travelling by $(1, 0)$ is a multiple of d . Since the knot class of H is (m_0, n_0) , there are exactly m_0 vertices in U that travel by $(1, 0)$. Therefore, d divides m_0 . The same argument shows that c divides n_0 .

Now let $m = \frac{m_0}{d}$ and $n = \frac{n_0}{c}$. We know that m and n are integers. Since $m_0 = md$ and $n_0 = nc$, we know md and nc are relatively prime. This proves conditions (1) and (2).

Now suppose that conditions (1) and (2) hold. By Theorem 2.4, our assumptions imply that there is a Hamiltonian circuit H' on $\mathbb{Z}_A \times \mathbb{Z}_B - \mathbb{Z}_1 \times \mathbb{Z}_1$. Note that the knot class of H' is (m, n) . We construct a spanning subgraph H of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ by requiring each vertex (x, y) in $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$ to travel in the same direction as the vertex (\bar{x}, \bar{y}) in $\mathbb{Z}_A \times \mathbb{Z}_B - \mathbb{Z}_1 \times \mathbb{Z}_1$. Here, \bar{x} and \bar{y} represent the residue classes of x and y , modulo A and modulo B , respectively. By construction, we know the knot class of H is (dm, cn) . The subgraph H is connected because we assumed that $\gcd(dm, cn) = 1$ [6]. Therefore, H is a Hamiltonian circuit of $(\mathbb{Z}_a \times \mathbb{Z}_b) - (\mathbb{Z}_c \times \mathbb{Z}_d)$. \square

Example 3.5. We give two examples illustrating the theorem. First, consider $(\mathbb{Z}_8 \times \mathbb{Z}_6) - (\mathbb{Z}_2 \times \mathbb{Z}_2)$ as shown in Figure 1. This digraph is not Hamiltonian because condition (2) of Theorem 3.1 can never be satisfied. On the other hand, $(\mathbb{Z}_8 \times \mathbb{Z}_6) - (\mathbb{Z}_1 \times \mathbb{Z}_2)$ is Hamiltonian; take $m = 1$ and $n = 5$.

4. QUESTIONS

We end with a few open questions related to our theorem. The first question extends our problem to larger dimensions.

Question 4.1. *When does the digraph $\mathbb{Z}_{a_1} \times \mathbb{Z}_{a_2} \times \cdots \times \mathbb{Z}_{a_n} - \mathbb{Z}_{c_1} \times \mathbb{Z}_{c_2} \times \cdots \times \mathbb{Z}_{c_n}$ have a Hamiltonian circuit?*

Not every subgroup of $\mathbb{Z}_a \times \mathbb{Z}_b$ is of the form $\mathbb{Z}_c \times \mathbb{Z}_d$. This leads to our next question.

Question 4.2. *Let A be any subgroup of $\mathbb{Z}_a \times \mathbb{Z}_b$. When does the digraph $\mathbb{Z}_a \times \mathbb{Z}_b - A$ have a Hamiltonian circuit?*

Instead of just removing a subgroup, it is also possible to remove more than one coset of a subgroup.

Question 4.3. *Choose a positive number r . When is it possible to remove r distinct cosets of $\mathbb{Z}_c \times \mathbb{Z}_d$ from $\mathbb{Z}_a \times \mathbb{Z}_b$ and obtain a Hamiltonian graph?*

The digraph $\mathbb{Z}_a \times \mathbb{Z}_b$ is said to be hyperhamiltonian if there is a spanning circuit which passes through one vertex exactly twice and all others exactly once. The digraph $\mathbb{Z}_a \times \mathbb{Z}_b$ is hyperhamiltonian if and only if there exist positive integers m and n such that $am + bn = ab + 1$ and $\gcd(m, n) = 1$ or 2 [4]. This idea can be generalized in the following way.

Question 4.4. *When is there a spanning circuit of $\mathbb{Z}_a \times \mathbb{Z}_b$ that passes through each vertex in the subgroup $\mathbb{Z}_c \times \mathbb{Z}_d$ exactly twice and all others exactly once?*

REFERENCES

- [1] G. Chartrand and L. Lesniak, *Graphs and Digraphs*, Chapman and Hall, New York (1996).
- [2] S. J. Curran and D. Witte, Hamilton paths in cartesian products of directed cycles. *Annals Discrete Math.* 27 (1985) 35-74.
- [3] W. T. Trotter, Jr. and P. Erdős, When the Cartesian product of directed cycles is Hamiltonian. *J. Graph Theory.* 2 (1978), 137-142.
- [4] J. Gallian and D. Witte, When the cartesian product of two directed cycles is hyperhamiltonian. *J. Graph Theory.* 11 (1987) 21-24.
- [5] L. E. Penn and D. Witte, When the cartesian product of two directed cycles is hypohamiltonian. *J. Graph Theory.* 7 (1983) 441-443.
- [6] D. Rolfsen, *Knots and Links*, Publish or Perish, Berkeley, CA (1976).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556