

1.(6pts) Let D be the region in the plane bounded by the circle $x^2 + y^2 = 4$ with density $\rho(x, y) = x^2 + y^2$. Which of the following statements is true:

- (a) The center of mass of D is the origin.
- (b) The center of mass of D is $(1, 1)$
- (c) The center of mass of D is $(0.5, -0.5)$
- (d) The center of mass of D is $(-0.5, 0.5)$
- (e) The center of mass of D is $(2, -2)$

2.(6pts) Calculate the volume enclosed by the paraboloid $z = x^2 + y^2$ and the plane $z = 1$.

- (a) $\frac{\pi}{2}$
- (b) π
- (c) $\frac{1}{2}$
- (d) 2π
- (e) 1

3.(6pts) Let S be the solid inside both the cone $z = \sqrt{x^2 + y^2}$ and the sphere $x^2 + y^2 + z^2 = 1$.

Write the iterated integral $\iiint_S z \, dV$ in spherical coordinates.

- (a) $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^1 \rho^3 \cos \phi \sin \phi \, d\rho \, d\phi \, d\theta$
- (b) $\int_0^{2\pi} \int_0^{\frac{\pi}{2}} \int_0^1 \rho^3 \cos \phi \sin \phi \, d\rho \, d\phi \, d\theta$
- (c) $\int_0^{2\pi} \int_0^{\pi} \int_0^1 \rho^3 \cos \phi \sin \phi \, d\rho \, d\phi \, d\theta$
- (d) $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^1 \rho^2 \cos \phi \sin \phi \, d\rho \, d\phi \, d\theta$
- (e) $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^1 \rho \cos \phi \, d\rho \, d\phi \, d\theta$

4.(6pts) Let a thin wire in space be in the shape of the helix C given by $\mathbf{r}(t) = \langle \cos 3t, \sin 3t, 3\sqrt{3}t \rangle$, $0 \leq t \leq 2\pi$. Let the linear density at (x, y, z) be given by $\mu(x, y, z) = \frac{z}{x^2 + y^2}$. Compute the mass of the wire.

- (a) $36\sqrt{3} \pi^2$
- (b) $216\sqrt{3} \pi^2$
- (c) $36\sqrt{3} \pi$
- (d) $6\sqrt{3} \pi^2$
- (e) 12π

5.(6pts) Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$, where $\mathbf{F} = -2xy\mathbf{i} + 4y\mathbf{j} + \mathbf{k}$ and $\mathbf{r} = t\mathbf{i} + t^2\mathbf{j} + \mathbf{k}$, $0 \leq t \leq 2$.

- (a) 24
- (b) 12
- (c) 48
- (d) 18
- (e) 32

6.(6pts) Let $\mathbf{F} = \nabla f$ where $f(x, y) = x^2 \cos(y) + e^{\sin(y)}$. Evaluate the integral $\int_C \mathbf{F} \cdot d\mathbf{r}$ where C is the semicircle $x^2 + y^2 = 1, x \geq 0$ and the orientation is counterclockwise.

- (a) $e^{\sin(1)} - e^{\sin(-1)}$
- (b) $e^{\sin(-1)} - e^{\sin(1)}$
- (c) -2
- (d) 2
- (e) $\cos(1) + \cos(-1)$

7.(6pts) Rewrite the iterated integral

$$\int_0^3 \int_0^{2-\frac{2z}{3}} \int_0^{6-2z-3y} \frac{e^z}{(z+2)(6-z)} dx dy dz$$

as an iterated integral $dy dx dz$. In other words, switch $dx dy$ to $dy dx$.

(a) $\int_0^3 \int_0^{6-2z} \int_0^{\frac{6-2z-x}{3}} \frac{e^z}{(z+2)(6-z)} dy dx dz$

(b) $\int_0^3 \int_0^{2-\frac{2z}{3}} \int_0^{6-2z-3x} \frac{e^z}{(z+2)(6-z)} dy dx dz$

(c) $\int_0^3 \int_0^{6-2z} \int_0^{6-2z-3x} \frac{e^z}{(z+2)(6-z)} dy dx dz$

(d) $\int_0^3 \int_0^{6-2z-3x} \int_0^{6-2z} \frac{e^z}{(z+2)(6-z)} dy dx dz$

(e) $\int_0^3 \int_0^{6-2y-3x} \int_0^{6-2x} \frac{e^z}{(z+2)(6-z)} dy dx dz$

8.(10pts) Let S be the “ice-cream cone” bounded below by $z = \sqrt{3(x^2 + y^2)}$ and above by $x^2 + y^2 + z^2 = 4$. Use spherical coordinates to express the volume of S as an integral. **Do not evaluate the integral.**

9.(10pts) Consider the double integral $\iint_D y dA$ where D is the region bounded by $y + 2x = 2$, $y + 2x = 6$, $y - 2x = -2$, and $y - 2x = 2$. Let T^{-1} be the transformation defined by $u = y + 2x$ and $w = y - 2x$. Describe the region R such that $T(R) = D$ and write $\iint_D y dA$

as a double integral over R . Note $x = \frac{1}{4}(u - w)$ and $y = \frac{1}{2}(u + w)$.

Do not evaluate the integral.

10. (10pts) Find a potential function for the field $\langle 2xe^{x^2y} + 2x^3ye^{x^2y} + 2xe^{x^2} - x^4e^{x^2y} + 2y \rangle$

~~11.(10pts) Use Green's Theorem to write down a double integral over the shaded region in Figure 12 which is equal to the line integral~~

$$\int_C (y + (\sin x)e^{-zx}) dx + (x^2 + (y \cos y)e^{y^2} + 2 - 3x) dy$$

along the boundary of the region.




Figure 12.

1. Solution. The region D and the density $\rho(x, y)$ are symmetric about the x -axis, hence the center of mass is located in the x -axis. Similarly, D and $\rho(x, y)$ are symmetric about the y -axis, so the center of mass is located in the y -axis. Therefore, the center of mass of D is the origin.

2. Solution. Using cylindrical co-ordinates, the paraboloid is $z = r^2$. The surfaces intersect when $r^2 = 1$, ie, $r = 1$. It follows that the volume is

$$\int_0^{2\pi} \int_0^1 \int_{r^2}^1 r \, dz \, dr \, d\phi = 2\pi \int_0^1 (r - r^3) \, dr = 2\pi \left(\frac{r^2}{2} - \frac{r^4}{4} \right) \Big|_0^1 = \frac{\pi}{2}$$

3. Solution. The cone $z = \sqrt{x^2 + y^2}$ is the same as $\phi = \frac{\pi}{4}$ in spherical coordinates and the sphere $x^2 + y^2 + z^2 = 1$ is $\rho = 1$. The limits of integration are therefore $0 \leq \rho \leq 1$, $0 \leq \phi \leq \frac{\pi}{4}$ and $0 \leq \theta \leq 2\pi$. The iterated integral is therefore

$$\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^1 \rho^3 \cos \phi \sin \phi \, d\rho \, d\phi \, d\theta$$

4. Solution. Recall

$$\int_C \mu(x, y, z) ds = \int_0^{2\pi} f(\mathbf{r}(t)) |\mathbf{r}'(t)| dt$$

First, we find $\mathbf{r}'(t)$:

$$\mathbf{r}'(t) = \langle -3 \sin 3t, 3 \cos 3t, 3\sqrt{3} \rangle$$

then

$$|\mathbf{r}'(t)| = \sqrt{(-3 \sin 3t)^2 + (3 \cos 3t)^2 + (3\sqrt{3})^2} = \sqrt{9 + 27} = \sqrt{36} = 6$$

So, the mass is

$$\begin{aligned} \int_C \mu(x, y, z) ds &= \int_0^{2\pi} \mu(\mathbf{r}(t)) |\mathbf{r}'(t)| dt \\ &= \int_0^{2\pi} \frac{3\sqrt{3} t}{(\cos 3t)^2 + (\sin 3t)^2} (6) dt \\ &= \int_0^{2\pi} 18\sqrt{3} t dt \\ &= 9\sqrt{3} t^2 \Big|_0^{2\pi} = 36\sqrt{3}\pi^2 \end{aligned}$$


5. Solution. Since $\mathbf{F}(\mathbf{r}(t)) = \langle -2t^3, 4t^2, 1 \rangle$ and $\mathbf{r}' = \langle 1, 2t, 0 \rangle$. We get that $\mathbf{F}(\mathbf{r}(t)) \bullet \mathbf{r}'(t) = 6t^3$. Thus $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^2 6t^3 dt = 24$.

6. Solution. Using the Fundamental Theorem of Line Integrals, we see that

$$\int_C \mathbf{F} \cdot d\mathbf{r} = f(0, 1) - f(0, -1) = (0 + e^{\sin(1)}) - (0 + e^{\sin(-1)}) = e^{\sin(1)} - e^{\sin(-1)}$$

.

7. Solution. The outer integral is the same in both cases so start by drawing the plane region determined by $\int_0^{2-\frac{2z}{3}} \int_0^{6-2z-3y} \dots dx dy$ for an arbitrary z between 0 and 1.




The integral in the other order has the outside integral running from $x = 0$ to $x = 6 - 2z$. In the inner integral, y starts at 0 and runs to $\frac{6 - 2z - x}{3}$.

8. Solution. The sphere $x^2 + y^2 + z^2 = 4$ is $\rho = 2$ and the cone $z = \sqrt{3(x^2 + y^2)}$ transforms as $\phi = \frac{\pi}{6}$. The volume is then given by

$$\iiint_S dV = \int_0^{2\pi} \int_0^{\frac{\pi}{6}} \int_0^2 \rho^2 \sin \phi d\rho d\phi d\theta$$

9. **Solution.** First let draw the region D .




In terms of u and w , $y + 2x = 2$ is equivalent to $u = 2$.

In terms of u and w , $y + 2x = 6$ is equivalent to $u = 6$.

In terms of u and w , $y - 2x = 2$ is equivalent to $w = 2$.

In terms of u and w , $y - 2x = -2$ is equivalent to $w = -2$.

The region R is the rectangle with corners (u, w) equal to $(2, 2)$, $(6, 2)$, $(2, -2)$, $(6, -2)$. Here is a picture.



Next So $\frac{\partial(x, y)}{\partial(u, w)} = \det \begin{vmatrix} \frac{\partial \frac{1}{4}(u-w)}{\partial u} & \frac{\partial \frac{1}{4}(u-w)}{\partial w} \\ \frac{\partial \frac{1}{2}(u+w)}{\partial u} & \frac{\partial \frac{1}{2}(u+w)}{\partial w} \end{vmatrix} = \det \begin{vmatrix} \frac{1}{4} & -\frac{1}{4} \\ \frac{1}{2} & \frac{1}{2} \end{vmatrix} = \frac{1}{8} - \left(-\frac{1}{8}\right) = \frac{1}{4}$.

Then

$$\iint_D y \, dA = \iint_R \left(\frac{u+w}{2} \cdot \frac{1}{4} \right) dA$$

10. Solution.

$$\frac{\partial p}{\partial y} = x^4 e^{x^2 y} + 2y$$

so

$$p = x^2 e^{x^2 y} + y^2 + h(x)$$

$$\frac{\partial p}{\partial x} = \frac{\partial x^2 e^{x^2 y} + y^2 + h(x)}{\partial x} = 2x e^{x^2 y} + x^2 (2xy) e^{x^2 y} + h'(x) = 2x e^{x^2 y} + 2x^3 y e^{x^2 y} + 2x e^{x^2}$$

so

$$\begin{aligned} h'(x) &= 2x e^{x^2} \\ p &= x^2 e^{x^2 y} + y^2 + e^{x^2} \end{aligned}$$

OR

Start with

$$\frac{\partial p}{\partial x} = 2x e^{x^2 y} + 2x^3 y e^{x^2 y} + 2x e^{x^2}$$

This is harder to integrate but it can be done. An anti-derivative of the first term is $\frac{e^{x^2 y}}{y}$.

An anti-derivative of the third term is e^{x^2} . To find an anti-derivative for the second term, use the substitution $t = x^2$, so $dt = 2x dx$ and $\int 2x^3 y e^{x^2 y} dx = \int y t e^{yt} dt$. This is a parts:

$u = t$ $dv = e^{yt} dt$ so $du = dt$ and $v = \frac{e^{yt}}{y}$.

$$\text{Hence } \int 2x^3 y e^{x^2 y} dx = \int y t e^{yt} dt = y t \frac{e^{yt}}{y} - y \int \frac{e^{yt}}{y} dt = t e^{yt} - \frac{e^{yt}}{y} = x^2 e^{x^2 y} - \frac{e^{x^2 y}}{y}.$$

Therefore

$$p = \frac{e^{x^2 y}}{y} + x^2 e^{x^2 y} - \frac{e^{x^2 y}}{y} + e^{x^2} + h(y) = x^2 e^{x^2 y} + e^{x^2} + h(y)$$

Then

$$\frac{\partial p}{\partial y} = x^4 e^{x^2 y} + h'(y) = x^4 e^{x^2 y} + 2y$$

so $h'(y) = 2y$ and $h(y) = y^2$ so

$$p = x^2 e^{x^2 y} + e^{x^2} + y^2$$

is a potential function.

11. Solution. Let D be the shaded region. $M = y^2 + (\sin x)e^x - 2x$ and $\frac{\partial M}{\partial y} = 2y$. $N = x^2 + (y \cos y)e^{y^2} + 2 - 3x$ and $\frac{\partial N}{\partial x} = 2x - 3$. Hence $\int_{\partial D} (y^2 + (\sin x)e^x - 2x)dx + (x^2 + (y \cos y)e^{y^2} + 2 - 3x)dy = \iint_D (2x - 3 + 2y) dA$.

Now $\int_{\partial D} (y^2 + (\sin x)e^x - 2x)dx + (x^2 + (y \cos y)e^{y^2} + 2 - 3x)dy$ is the line integral around the boundary curve oriented so that the region is on your left. This is the given orientation on C so $\int_C (y^2 + (\sin x)e^x - 2x)dx + (x^2 + (y \cos y)e^{y^2} + 2 - 3x)dy = \iint_D (2x - 3 + 2y) dA$.
