

VANISHING HOMOLOGY OVER NILPOTENT GROUPS¹

WILLIAM G. DWYER

ABSTRACT. Let π be a nilpotent group and let M be a π -module. Under certain finiteness assumptions we prove that the twisted homology groups $H_i(\pi, M)$ vanish for all positive i whenever $H_0(\pi, M) = 0$.

The purpose of this note is to prove the following vanishing theorem:

(1) **Theorem.** *Let π be a finitely generated nilpotent group, and let M be a π -module which is finitely generated over $\mathbf{Z}[\pi]$. Assume that $H_0(\pi, M) = 0$. Then $H_i(\pi, M) = 0$ for all $i \geq 0$.*

In the statement of this theorem, the π -module M is, as usual, an abelian group equipped with a left π -action, $\mathbf{Z}[\pi]$ is the integral group ring of π , and $H_i(\pi, M)$, $i \geq 0$, are the twisted homology groups defined in [4].

With a little care, the proof below will also yield the following more general result:

(2) **Theorem.** *Let π be as in (1), and let $\{M_s\}_{s \geq 0}$ be a tower of π -modules, each of which is finitely generated over $\mathbf{Z}[\pi]$. Then if $\{H_0(\pi, M_s)\}_{s \geq 0}$ is protrivial, all the other towers $\{H_i(\pi, M_s)\}_{s \geq 0}$ for $i > 0$ are protrivial too.*

For a definition of the terms in this statement, and a discussion of the basic properties of towers, see [1].

Several topological applications of these theorems will be examined in forthcoming papers [2], [3]. Our statements are parallel to the results of [5], although very different in detail.

The author does not know in what sense these theorems are "best possible." There are examples to show that the finite generation condition on M and the nilpotency condition on π are unavoidable, but the finite generation condition on π may well be redundant. In fact, a simple proof of (1)

Received by the editors December 3, 1973.

AMS (MOS) subject classifications (1970). Primary 18H10; Secondary 20E15, 20D15, 20K40.

Key words and phrases. Nilpotent group, twisted group homology.

¹Research supported by NSF GP-38590.

for arbitrary abelian π appears at the end of this note, but the author cannot see how to generalize it.

Preliminaries. In the discussion below, a π -module M such that $H_0(\pi, M)$ vanishes is called *perfect*; a module M such that $H_i(\pi, M)$ vanishes for all $i \geq 0$ is called *acyclic*. Three observations will be used repeatedly:

(3) *Any quotient module of a perfect π -module is perfect.*

This follows at once from the fact that $H_0(\pi, -)$ is right exact.

(4) *If M' is a submodule of M , M is perfect, and M/M' is acyclic, then M' is perfect.*

This follows at once from the long exact homology sequence of $0 \rightarrow M' \rightarrow M \rightarrow M/M' \rightarrow 0$.

(5) *The integral group ring of a finitely generated nilpotent group is (left and right) noetherian.*

This is the backbone of the argument below. A proof is indicated in [6].

Now the proof of (1) proceeds by induction on the number of central cyclic extensions needed to construct π . Consequently, we can assume that σ is a cyclic subgroup of the center of π and that the obvious inductive theorem is known for π/σ -modules. Let M be a finitely generated perfect $\mathbb{Z}[\pi]$ -module. We let s be a generator of σ , and T the endomorphism of M given by $m \mapsto m - s \cdot m$ for all $m \in M$. The letter p will denote the order of σ , which can be assumed either prime or infinite; if p is infinity, then by convention every element of every abelian group is said to be of order p .

Special cases. In this paragraph we will prove that M is acyclic if it has one of the following three special forms:

Type I. T is injective on M .

Type II. T is the zero map $M \rightarrow M$, and M has no elements of order p . (This type is trivial if $p = \infty$.)

Type III. T is the zero map $M \rightarrow M$, and every element of M has order p .

If M falls into one of these three classes, we will compute the E^2 -term of the Lyndon spectral sequence $[M]$

$$E_{p,q}^2 = H_p(\pi/\sigma, H_q(\sigma, M)) \Rightarrow H_{p+q}(\pi, M)$$

and show that it vanishes. Actually, the computation below will only show that

$$H_0(\pi/\sigma, H_j(\sigma, M)) = E_{0,j}^2 = 0 \quad \text{for all } j.$$

The induction hypothesis, together with the easily proven fact that each $H_j(\sigma, M)$ is a finitely generated $\mathbb{Z}[\pi/\sigma]$ -module, will then give $H_i(\pi/\sigma, H_j(\sigma, M)) = E_{i,j}^2 = 0$ for all $i, j \geq 0$.

The computation depends on explicit knowledge of the homology groups $H_i(\sigma, M)$, which are well known to be given as follows [4]:

Case I. σ is finite of order p . Let N be the endomorphism of M given by $1 + s + s^2 + \dots + s^{p-1}$.

$$H_0(\sigma, M) = M/\text{image}(T),$$

$$H_{2i}(\sigma, M) = \text{kernel}(N)/\text{image}(T), \quad i \geq 1,$$

$$H_{2i+1}(\sigma, M) = \text{kernel}(T)/\text{image}(N), \quad i > 0.$$

Case II. σ is infinite cyclic.

$$H_0(\sigma, M) = M/\text{image}(T), \quad H_1(\sigma, M) = \text{kernel}(T), \quad H_i(\sigma, M) = 0, \quad i > 1.$$

The computation now breaks into three parts, according to the structure of M . Recall that we are given $H_0(\pi/\sigma, H_0(\sigma, M)) \simeq H_0(\pi, M) = 0$.

Type I. In this case $H_{2i+1}(\sigma, M) = 0$ for all $i \geq 0$. If σ is infinite cyclic, $H_{2i}(\sigma, M)$, $i \geq 1$, is zero, so there is nothing more to show. Otherwise, if σ is finite, each $H_{2i}(\sigma, M)$, $i \geq 1$, is a sub- π/σ -module of $H_0(\sigma, M)$, and so, by the argument of (4), $H_0(\pi/\sigma, H_{2i}(\sigma, M)) = 0$.

Type II. If σ is infinite, there is nothing to show. If σ is finite, then $H_{2i}(\sigma, M) = 0$ for $i \geq 1$ and $H_{2i+1}(\sigma, M)$, $i \geq 0$, is a quotient π/σ -module of $H_0(\sigma, M)$, and so, by (3), $H_0(\pi/\sigma, H_{2i+1}(\sigma, M)) = 0$.

Type III. If σ is finite, then $H_i(\sigma, M) \simeq M \simeq H_0(\sigma, M)$ for all $i > 0$, so that $H_0(\pi/\sigma, H_i(\sigma, M)) = 0$. If σ is infinite, then $H_1(\sigma, M) \simeq M \simeq H_0(\sigma, M)$, and $H_i(\sigma, M) = 0$ for $i > 1$, so the same argument works.

The general case. Let M be an arbitrary perfect finitely generated $\mathbb{Z}[\pi]$ -module. In order to show that M is acyclic, it is enough to show that there is a finite π -filtration of M ,

$$0 = F_0 \subseteq F_1 \subseteq F_2 \subseteq \dots \subseteq F_k = M$$

such that each filtration quotient F_{i+1}/F_i is of Type I, II, or III. Indeed, if such a filtration exists, then we show by descending induction on i that

M/F_i is acyclic for all $0 \leq i \leq k$. The induction starts with the fact that $M/F_k = M/M = 0$ is acyclic. Now suppose that M/F_i is acyclic, $i > 0$. There is a short exact sequence

$$0 \rightarrow F_i/F_{i-1} \rightarrow M/F_{i-1} \rightarrow M/F_i \rightarrow 0.$$

By (3) M/F_{i-1} , as a quotient of M , is a perfect π -module. The long exact homology sequence of this short exact sequence, and the fact that M/F_i is acyclic, show that F_i/F_{i-1} is perfect. Since this π -module is of Type I, II, or III, it must be acyclic. Another look at the long exact homology sequence verifies that M/F_{i-1} is acyclic.

Constructing such a filtration of M is not hard. Let T be as above, and define F_i ($i \geq 0$) by

$$F_0 = \{0\} \subset M, \quad F_i = \text{kernel}\{T^i: M \rightarrow M\}, \quad i \geq 1.$$

Since σ is in the center of π , $\{F_i | i \geq 0\}$ is a family of π -equivariant submodules of M . By the noetherian condition (5), this family must have a maximal element, so, for some $K \geq 0$, $F_K = F_{K+1}$. Then M/F_K is of Type I. If σ is infinite cyclic, each F_{i+1}/F_i is already of Type III, and we are done. Otherwise, if σ is finite of order p , it is enough to show that each F_{i+1}/F_i can be filtered (in a π -equivariant way) so that the filtration quotients are of Type II or III. To do this, pick $0 \leq i \leq K-1$, and define G_j ($j \geq 0$) by

$$G_0 = F_i, \quad G_j = \{x \in F_{i+1} | p^j x \in F_i\}.$$

Again by the noetherian condition, there is some $J \geq 0$ such that $G_J = G_{J+1}$. Clearly F_{i+1}/G_J is of Type II, and each G_{j+1}/G_j is of Type III. This completes the proof.

A simple proof of the abelian case. We give a conceptual proof for arbitrary abelian π that any perfect π -module M which is finitely generated over $\mathbf{Z}[\pi]$ is acyclic. It is enough to assume that M has a single generator m over $\mathbf{Z}[\pi]$; induction on the number of generators, using (4), then gives the general case.

Let $I \subseteq \mathbf{Z}[\pi]$ be the augmentation ideal—the kernel of the natural epimorphism $\mathbf{Z}[\pi] \rightarrow \mathbf{Z}$. By hypothesis, $H_0(\pi, M) = \mathbf{Z} \otimes_{\mathbf{Z}[\pi]} M = M/I \cdot M = 0$, so there must be some $r \in I$ such that $r \cdot m = m$. Since π is abelian, left multiplication by r commutes with the action of π and so must induce the identity map $M \rightarrow M$, and therefore the identity map $H_*(\pi, M) \rightarrow H_*(\pi, M)$.

However, again since π is abelian, left multiplication by r is easily seen to induce a natural transformation of the functor $H_*(\pi, -)$ into itself; this natural transformation is evidently zero on $H_0(\pi, -)$ and so, by the basic theorems about derived functors, must be identically zero. Consequently, the identity map $H_*(\pi, M) \rightarrow H_*(\pi, M)$ coincides with the zero map.

REFERENCES

1. A. K. Bousfield and D. M. Kan, *Homotopy limits, completions, and localizations*, Springer, New York, 1972.
2. W. Dwyer, *Generalized convergence of the Eilenberg-Moore spectral sequence* (in preparation).
3. E. Dror, *Homology circles* (in preparation).
4. S. Mac Lane, *Homology*, Die Grundlehren der math. Wissenschaften, Band 114, Academic Press, New York; Springer-Verlag, Berlin, 1963. MR 28 # 122.
5. T. Nakayama, *On modules of trivial cohomology over a finite group*, Illinois J. Math. 1 (1957), 36-43. MR 18, 793.
6. D. Passman, *Infinite group rings*, Dekker, New York, 1971, p. 136.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CONNECTICUT
06520